

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 March 2006 (23.03.2006)

PCT

(10) International Publication Number
WO 2006/031291 A2

(51) International Patent Classification:

C12N 9/36 (2006.01) C12N 1/21 (2006.01)
C07H 21/04 (2006.01) C12N 15/75 (2006.01)

(21) International Application Number:

PCT/US2005/025831

(22) International Filing Date: 21 July 2005 (21.07.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

10/939,262 10 September 2004 (10.09.2004) US

(71) Applicants and

(72) Inventors: FANG, Fang [US/US]; 11124 Corte Plano Verano, San Diego, CA 92130 (US). MALAKHOV, Michael [RU/US]; 11530 Camino La Bar, San Diego, CA 92130 (US).

(74) Agents: PRESTON, David et al.; David R Preston & Associates, 12625 High Bluff Drive, Suite 205, San Diego, CA 92130 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A NOVEL CLASS OF THERAPEUTIC PROTEIN BASED MOLECULES



(57) Abstract: The present invention provides new compositions and methods for preventing and treating pathogen infection. In particular, the present invention provides compounds having an anchoring domain that anchors the compound to the surface of a target cell, and a therapeutic domain that can act extracellularly to prevent infection of a target cell by a pathogen, such as a virus. The present invention also comprises therapeutic compositions having sialidase activity, including protein-based compounds having sialidase catalytic domains. Compounds of the invention can be used for treating or preventing pathogen infection, and for treating and reducing allergic and inflammatory responses. The invention also provides compositions and methods for enhancing transduction of target cells by recombinant viruses. Such compositions and methods can be used

in gene therapy.

A NOVEL CLASS OF THERAPEUTIC PROTEIN BASED MOLECULES

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims benefit of priority to United States patent application serial number 10/939,262 filed 10 September 2004, entitled, "Novel Class of Therapeutic Protein Based Molecules," which is herein incorporated by reference in its entirety.

10 The present application also incorporates by reference in their entirety the following patent applications: United States application 10/718,986, filed 21 November 2003, entitled "Broad spectrum anti-viral therapeutics and prophylaxis", United States provisional patent application serial number 60/428,535, filed 22 November 2002, entitled "Broad spectrum anti-viral therapeutics and prophylaxis", United States provisional patent application serial number 60/464,217, filed 19 April 2003, entitled
15 "Class of broad spectrum anti-viral protein", United States provisional patent application serial number 60/561,749, filed 13 April 13 2004, entitled "Anti-microbial therapeutics and prophylaxis", United States provisional patent application serial number 60/580,084, filed 16 June 16, 2004, entitled "Class of broad spectrum anti-microbial agents".

20

BACKGROUND OF THE INVENTION

25 The invention relates to therapeutic compositions that can be used to prevent and treat infection of human and animal subjects by a pathogen, and specifically to protein-based therapeutic compositions that can be used for the prevention and treatment of viral or bacterial infections. The invention also relates to therapeutic protein-based compositions that can be used to prevent or ameliorate allergic and inflammatory responses. The invention also relates to protein-based compositions for increasing transduction efficiency of a recombinant virus, such as a recombinant virus used for gene
30 therapy.

Influenza is a highly infectious acute respiratory disease that has plagued the human race since ancient times. It is characterized by recurrent annual epidemics and periodic major worldwide pandemics. Because of the high disease-related morbidity and mortality, direct and indirect social economic impacts of influenza are enormous. Yearly epidemics cause approximately 300,000 hospitalizations and 25,000 deaths in the United States alone. Four pandemics occurred in the last century; together they caused tens of millions of deaths. Mathematical models based on earlier pandemic experiences have estimated that 89,000-207,000 deaths, 18-42 million outpatient visits and 20-47 million additional illnesses will occur during the next pandemic (Meltzer, MI, Cox, NJ and Fukuda, K. (1999) *Emerg Infect Dis* 5:659-671).

Influenza is typically caused by infection of two types of viruses, *Influenza virus A* and *Influenza virus B* (the third type *Influenza virus C* only causes minor common cold like symptoms). They belong to the *orthomyxoviridae* family of RNA viruses. Both type A and type B viruses have 8 segmented negative-strand RNA genomes enclosed in a lipid envelope derived from the host cell. The viral envelope is covered with spikes that are composed of three types of proteins: hemagglutinin (HA) which attaches virus to host cell receptors and mediates fusion of viral and cellular membranes; neuraminidase (NA) which facilitates the release of the new viruses from host cells; and a small number of M2 proteins which serve as ion channels.

Infections by influenza type A and B viruses are typically initiated at the mucosal surface of the upper respiratory tract. Viral replication is primarily limited to the upper respiratory tract but can extend to the lower respiratory tract and cause bronchopneumonia that can be fatal.

Influenza viral protein hemagglutinin (HA) is the major viral envelope protein. It plays an essential role in viral infection. The importance of HA is evidenced by the fact that it is the major target for protective neutralizing antibodies produced by the host immune response (Hayden, FG. (1996) In *Antiviral drug resistance* (ed. D. D. Richman), pp. 59-77. Chichester, UK: John Wiley & Sons Ltd.). It is now clear that HA has two different functions in viral infection. First, HA is responsible for the attachment of the virus to sialic acid cell receptors. Second, HA mediates viral entry into target cells by triggering fusion of the viral envelope with cellular membranes.

HA is synthesized as a precursor protein, HA0, which is transferred through the Golgi apparatus to the cell surface as a trimeric molecular complex. HA0 is further cleaved to generate the C terminus HA1 (residue 328 of HA0) and the N terminus of HA2. It is generally believed that the cleavage occurs at the cell surface or on released
5 viruses. The cleavage of HA0 into HA1/HA2 is not required for HA binding to sialic acid receptor; however, it is believed to be necessary for viral infectivity (Klenk, HD and Rott, R. (1988) *Adv Vir Res.* 34:247-281; Kido, H, Niwa, Y, Beppu, Y and Towatari, T. (1996) *Advan Enzyme Regul* 36:325-347; Skehel, JJ and Wiley, DC. (2000) *Annu Rev Biochem* 69:531-569; Zambon, M. (2001) *Rev Med Virol* 11:227-241.)

10 Currently, influenza is controlled by vaccination and anti-viral compounds. Inactivated influenza vaccines are now in worldwide use, especially in high-risk groups. The vaccine viruses are grown in fertile hen's eggs, inactivated by chemical means and purified. The vaccines are usually trivalent, containing representative influenza A viruses (H1N1 and H3N2) and influenza B strains. The vaccine strains need to be regularly
15 updated in order to maintain efficacy; this effort is coordinated by the World Health Organization (WHO). During inter-pandemic periods, it usually takes 8 months before the updated influenza vaccines are ready for the market (Wood, J. (2001) *Phil Trans R Soc Lond B* 356:1953-1960). However, historically, pandemics spread to most continents within 6 months, and future pandemics are expected to spread even faster with increased
20 international travel (Gust, ID, Hampson, AW., and Lavanchy, D. (2001) *Rev Med Virol* 11:59-70). Therefore it is inevitable that an effective vaccine will be unavailable or in very short supply during the first waves of future pandemics.

Anti-viral compounds have become the mainstay for treating inter-pandemic diseases. Currently, they are also the only potential alternative for controlling pandemics
25 during the initial period when vaccines are not available. Two classes of antiviral compounds are currently on the market: the M2 inhibitors, such as amantadine and rimantadine; and the NA inhibitors, which include oseltamivir (Tamiflu) and zanamivir (Relenza). Both classes of molecules have proven efficacy in prevention and treatment of influenza. However, side effects and the risk of generating drug-resistant viruses remain
30 the top two concerns for using them widely as chemoprophylaxis (Hayden, FG. (1996) In *Antiviral drug resistance* (ed. D. D. Richman), pp. 59-77. Chichester, UK: John Wiley

& Sons Ltd.). Most importantly, future pandemic strains, either evolved naturally or artificially created by genetic engineering in bio-warfare, may be resistant to all the available anti-viral compounds, and this will have devastating consequences globally.

In summary, currently available vaccination and anti-viral compounds are limited
5 by some fundamental shortcomings. Novel therapeutic and prophylactic modalities are needed to address future influenza pandemics.

Respiratory tract infections (RTIs) are the most common, and potentially most severe, types of infectious diseases. Clinically, RTIs include sinusitis, otitis,
10 laryngitis, bronchitis and pneumonia. Based on numerous etiology and epidemiology studies, it is clear that although many microorganisms have the potential to cause RTIs, only a handful of pathogens are responsible for vast majority of the cases. Such pathogens include *S. pneumoniae*, *M. pneumoniae*, *H. influenzae*, *M. catarrhalis*, influenza A & B, and parainfluenza virus. Besides causing CAP and AECB, several of
15 the bacterial pathogens, such as *S. pneumoniae* and *H. influenzae*, are also the common cause of acute sinusitis, otitis media, as well as invasive infections leading to sepsis, meningitis, etc. Therefore these microorganisms are of the highest clinical importance.

One common feature of all respiratory pathogenic bacteria is that they establish commensal colonization on the mucosal surface of the upper airway; such colonization
20 precedes an infection and is prerequisite for infections. The bacterial colonization in a neonate occurs shortly after birth. During lifetime, the upper airway, specifically the nasopharynx and oropharynx, remains a dynamic ecological reservoir of microbial species with bacteria being acquired, eliminated and re-acquired continually. In most cases the bacterial flora in the pharynx is harmless. However, when the condition of the
25 host is altered, some microorganisms may invade adjacent tissues or bloodstream to cause diseases. In addition to serving as the port of entry for mucosal and invasive infections by both bacteria and viruses, the nasopharynx is also the major source of spreading the pathogenic microorganisms between individuals, as well as the reservoir where antibiotic-resistant bacteria are selected (Garcia-Rodriguez and Martinez, J
30 Antimicrob Chemother, (2002) 50(Suppl S2), 59-73; Soriano and Rodriguez-Cerrato, J Antimicrob Chemother, (2002) 50 Suppl S2, 51-58). It is well established clinically

that individuals who are prone to RTIs tend to be persistent and recurrent carriers of the pathogenic bacteria (Garcia-Rodriguez and Martinez, J Antimicrob Chemother, (2002) 50(Suppl S2), 59-73; Mbaki et al., Tohoku J Exp. Med., (1987) 153(2), 111-121).

5 *Helicobacter pylori* is a human pathogen implicated in gastritis and peptic ulcer. The bacterium resides in the human stomach and binds to epithelial cells of the gastric antrum. It has been demonstrated that the bacterial adhesion is mediated by binding of *Helicobacter pylori* adhesin I and II to sialic acids on the epithelial surface.

10 Siglecs (sialic acid binding Ig-like lectins) are members of the immunoglobulin (Ig) superfamily that bind to sialic acid and are mainly expressed by cells of the hematopoietic system. At least 11 siglecs have been discovered and they seem to exclusively recognize cell surface sialic acid as the ligand. It is believed that the binding of siglecs to sialic acid mediates cell-cell adhesion and interactions (Crocker and Varki, Trends Immunol., (2001) 22(6), 337-342; Angata and Brinkman-Van der Linden, 15 Biochim. Biophys. Acta, (2002) 1572(2-3), 294-316). Siglec-8 (SAF-2) is an adhesion molecule that is highly restricted to the surface of eosinophils, basophils, and mast cells, which are the central effector cells in allergic conditions including allergic rhinitis, asthma and eczema. Siglec-8 is considered to be responsible for mediating the recruitment of the three allergic cell types to the airway, the lungs and other sites of 20 allergy. Siglec-1 (sialoadhesion) and siglec-2 (CD22) are the adhesion molecules on macrophages and B cells, both types of cells play central roles in immune reactions that lead to inflammation.

25 Recombinant viruses, in particular adeno-associated virus (AAV), can be used to transfer the wild type cystic fibrosis transmembrane conductance regulator (CFTR) gene into the epithelial cells to correct the genetic defect that causes cystic fibrosis (Flotte and Carter, Methods Enzymol., (1998) 292, 717-732). Clinical trials with AAV vectors have shown efficient and safe delivery of the CFTR gene into epithelial cells with low levels of gene transfer (Wagner et al., Lancet, (1998) 351(9117), 1702-1703). Compared 30 to adenoviral vectors, AAV offers more stable gene expression and diminished cellular immunity. However, the transduction efficiency of AAV in vivo is rather low in the lung

(Wagner et al., Lancet, (1998) 351(9117), 1702-1703). A method that can improve transduction efficiency of AAV in vivo is needed to achieve full therapeutic potential of gene therapy for cystic fibrosis. It has been shown that negatively charged carbohydrates, such as sialic acid, inhibit the transduction efficiency of AAV vector to the well-differentiated airway epithelium, and treatment of the airway epithelium by glycosidases, including a neuraminidase, and endoglycosidase H, enhances transduction efficiency of the AAV vector (Bals et al., J Virol., (1999) 73(7), 6085-6088).

10

BRIEF SUMMARY OF THE INVENTION

15

The present invention recognizes that current therapeutics for preventing and treating infection by pathogens are often difficult to provide in a timely manner, can have undesirable side effects, and can lead to drug-resistant pathogen strains. The present invention also recognizes that the current approach to treat allergy and inflammation has limited efficacy and is associated with side effects. In addition, the present invention also recognizes that the current approach to administer recombinant viruses yield low transduction efficiency and unsatisfactory efficacy of the gene therapy.

20

The present invention provides new compositions and methods for preventing and treating pathogen infection. In particular, the present invention provides compounds that can act extracellularly to prevent infection of a cell by a pathogen. Some preferred embodiments of the present invention are therapeutic compounds having an anchoring domain that anchors the compound to the surface of a target cell, and a therapeutic domain that can act extracellularly to prevent infection of the target cell by a pathogen, such as a virus or bacterium.

25

30

In one aspect, the invention provides a protein-based composition for preventing or treating infection by a pathogen. The composition comprises a compound that comprises at least one therapeutic domain comprising a peptide or protein, where the therapeutic domain has at least one extracellular activity that can prevent the infection of a target cell by a pathogen, and at least one anchoring domain that can bind at or near the membrane of a target cell.

In some embodiments of this aspect of the present invention, the at least one therapeutic domain comprises an inhibitory activity that prevents or impedes the infection of a target cell by a pathogen. In a preferred embodiment, the inhibitory activity inhibits the activity of a protease that can process a viral protein necessary for infection of a target cell. In a particularly preferred embodiment, the compound comprises a therapeutic domain that can inhibit the processing of the HA protein of influenza virus, and the anchoring domain can bind the compound at the surface of a respiratory epithelial cell.

In some embodiments of the present invention, at least one therapeutic domain comprises a catalytic activity. In a preferred embodiment, the catalytic activity removes a moiety from the surface of a target cell that is necessary for infection of the target cell. In a particularly preferred embodiment, the therapeutic domain is a sialidase that can digest sialic acid moieties on the surface of epithelial target cells, and the anchoring domain is a GAG-binding domain of a human protein that can bind heparin or heparan sulfate moieties at the surface of an epithelial cell.

In another aspect, the present invention includes pharmaceutical compositions for treating or preventing pathogen infection in a subject. Pharmaceutical compositions comprise a compound of the present invention comprising at least one therapeutic domain and at least one anchoring domain. The pharmaceutical composition can also comprise solutions, stabilizers, fillers and the like. In some preferred embodiments, the pharmaceutical composition is formulated as an inhalant. In some preferred embodiments, the pharmaceutical composition is formulated as a nasal spray.

Another aspect of the present invention is a pharmaceutical composition comprising at least one sialidase. The sialidase can be isolated from any source, such as, for example, a bacterial or mammalian source, or can be a recombinant protein that is substantially homologous to a naturally occurring sialidase. A pharmaceutical composition comprising a sialidase can be formulated for nasal, tracheal, bronchial, oral, or topical administration, or can be formulated as an injectable solution or as eyedrops. A pharmaceutical composition comprising a sialidase can be used to treat or prevent pathogen infection, to treat or prevent allergy or inflammatory response, or to enhance the transduction efficiency of a recombinant virus for gene therapy.

Yet another aspect of the present invention is a sialidase catalytic domain protein. In this aspect, proteins that comprise the catalytic domain of a sialidase but comprise less than the entire sialidase the catalytic domain sequence is derived from are considered sialidase catalytic domain proteins. Sialidase catalytic domain proteins can comprise
5 other protein sequences, such as but not limited to functional domains derived from other proteins. A pharmaceutical composition comprising a sialidase can be formulated for nasal, tracheal, bronchial, oral, or topical administration, or can be formulated as an injectable solution or as eyedrops. A pharmaceutical composition comprising a sialidase can be used to treat or prevent pathogen infection, to treat or prevent allergy or
10 inflammatory response, or to enhance the transduction efficiency of a recombinant virus for gene therapy.

In yet another aspect, the present invention includes a method for treating or preventing infection by a pathogen. In preferred embodiments, the method comprises administering a sialidase activity, such as a sialidase or a sialidase catalytic domain
15 protein, including a sialidase catalytic domain fusion protein, to a subject to prevent or treat an infection. A pathogen can be, for example, a viral or bacterial pathogen. The method includes applying a pharmaceutically effective amount of a compound of the present invention to at least one target cell of a subject. Preferably, the pharmaceutical composition can be applied by the use of a spray, inhalant, or topical formulation.

20 The present invention also provides new compositions and methods for treating allergy and inflammation. In particular, the present invention provides compounds that can act extracellularly to prevent or inhibit adhesion and function of inflammatory cells. Some preferred embodiments of compounds for treating allergy or inflammation comprise at least one therapeutic domain that has the said extracellular activity and an at
25 least one anchoring domain that anchors the compound to the surface of a target cell. In some preferred embodiments, the method comprises administering a sialidase activity, such as a sialidase or a sialidase catalytic domain protein, including a sialidase catalytic domain fusion protein to a subject to prevent or treat an allergic or inflammatory response. The allergic or inflammatory response can be asthma, allergic rhinitis, skin
30 conditions such as eczema, or response to plant or animal toxins. The method includes applying a pharmaceutically effective amount of a compound of the present invention to

at least one target cell of a subject. Preferably, the pharmaceutical composition can applied by the use of a spray, inhalant, or topical formulation.

The present invention also provides new compositions and methods for improving efficiency of gene transfer by recombinant viral vectors during gene therapy. In particular, the present invention provides compounds that can act extracellularly to
5 reduce the physical or chemical barrier that hinders transduction by gene therapy vectors, such as AAV vector. Some preferred compounds of the present invention for improving efficiency of gene transfer by recombinant viral vectors comprise at least one therapeutic domain that has an extracellular activity and an at least one anchoring domain that
10 anchors the compound to the surface of a target cell. In some preferred embodiments, the method comprises administering a sialidase activity, such as a sialidase or a sialidase catalytic domain protein, including a sialidase catalytic domain fusion protein to a subject to facilitate transduction of a target cell by a recombinant viral vector. The method includes applying an effective amount of a compound of the present invention along with
15 a recombinant viral vector to at least one target cell. A pharmaceutical composition of the present invention can applied by the use of a spray, inhalant, or topical formulation.

BRIEF DESCRIPTION OF SEVERAL VIEWS OF THE DRAWINGS

20 **Figure 1** is a schematic depiction of the primary amino acid structure of aprotinin.

Figure 2 shows GAG-binding sequences of four human genes: PF4, human platelet factor 4; IL8, human interleukin 8; AT III, human antithrombin III; ApoE, human apolipoprotein E; AAMP, human angio-associated migratory cell protein.

25 **Figure 3** is a sequence comparison between human sialidases NEU2 and NEU4.

Figure 4 is a table comparing substrate specificity of bacterial and fungal sialidases.

30 **Figure 5** depicts the nucleotide and amino acid sequences of Construct #1 encoding His6-AvCD. NcoI and HindIII sites used for cloning into pTrec99a are shown in bold.

Figure 6 depicts the nucleotide and amino acid sequences of Construct #2 encoding AR-AvCD. NcoI and HindIII sites used for cloning into pTrc99a are shown in bold.

- 5 **Figure 7 depicts the nucleotide and amino acid sequences of Construct #3 encoding AR-G₄S-AvCD. NcoI and HindIII sites used for cloning into pTrc99a are shown in bold.**

Figure 8 is a graph of data from an experiment showing that the AR-tag enhances the removal of $\alpha(2,6)$ -linked sialic acid from MDCK cells. The Y axis shows the percentage of $\alpha(2,6)$ -linked sialic acid remaining on the surface of MDCK cells after treatment with various dilutions of recombinant AvCD (Construct #1) (diamonds) or recombinant AR-AvCD (Construct #2) (squares).

10

Figure 9 is a graph depicting the protection against influenza viruses conferred by treating MDCK cells with recombinant AR-AvCD protein made from Construct #2 or the isolated sialidase of *A. ureafaciens*. The challenge viral strains are: A/WS/33 (H1N1); A/PR/8 (H1N1); A/Japan/305/57 (H2N2); A/Victoria/504/2000 (H3N2); A/HongKong/8/68 (H3N2); B/Lee/40; 7. B/Maryland/1/59; and Turkey/Wis/66 (H9N2).

15

Figure 10 is a graph showing the level of inhibition of influenza virus amplification by the recombinant AR-AvCD sialidase and the recombinant AR-G₄S-AvCD sialidase. The challenge viral strains are: A/PR/8 (H1N1); A/WS/33 (H1N1); A/Japan/305/57 (H2N2); A/HongKong/8/68 (H3N2); B/Lee/40; 7. B/Maryland/1/59; and Turkey/Wis/66 (H9N2).

20

Figure 11 provides graphs showing that topical administration of recombinant AR-AvCD sialidase fusion protein reduces the inflammatory responses of ferrets infected with an influenza A (H1N1) virus. (A) The total number of inflammatory cells from nasal wash samples obtained from infected animals at the indicated times after infection. (B) The protein concentration was determined in cell-free nasal wash samples of infected ferrets. Infected ferrets were vehicle-treated (squares) or were treated with recombinant AR-AvCD sialidase fusion protein made from Construct #2 (triangles). Uninfected animals

25

30

were also treated with recombinant AR-AvCD sialidase fusion protein (diamonds). Statistically significant values are labeled with * ($p < 0.05$) and ** ($p < 0.01$).

5 **Figure 12** is a table depicting inhibition of viral replication, cell protection EC₅₀'s, and selective indexes for two sialidase catalytic domain fusion proteins of the present invention. All EC₅₀'s are in mU/ml.

10 **Figure 13** is a table depicting viral replication in the respiratory tract of ferrets treated with a sialidase catalytic domain fusion proteins of the present invention and ferrets treated with a control vehicle.

DETAILED DESCRIPTION OF THE INVENTION

15 Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Generally, the nomenclature used herein and the manufacture or laboratory procedures described below are well known and commonly employed in the art. Conventional methods are used for these procedures, such as those provided in the art and various general references. Where a term is provided in the singular, the inventors also contemplate the plural of that term. Where there are discrepancies in terms and definitions used in references that are incorporated by reference, the terms used in this application shall have the definitions given herein. As employed throughout the disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

20
25

A "pathogen" can be any virus or microorganism that can infect a cell, a tissue or an organism. A pathogen can be a virus, bacterium, or protozoan.

A "target cell" is any cell that can be infected by a pathogen or any cell that can interact with inflammatory cells, or a host cell that is the intended destination for an exogenous gene transferred by a recombinant virus.

5 A "recombinant virus" or a "recombinant viral vector", a "gene therapy viral vector" or a "gene therapy vector" is defined as a genetically engineered virus that comprises one or more exogenous genes. When a target cell is transduced by a recombinant virus, the exogenous gene(s) is transferred to the target cell. Genes transferred to a target cell can be expressed in the cell to provide the intended therapeutic effects. Currently, most commonly used gene therapy viral vectors are based on four
10 types of viruses: retrovirus (including lentivirus), adeno-virus, adeno-associated virus (AAV) and herpes simplex virus type 1.

"Inflammatory cells" are the cells that carry out or participate in inflammatory responses of the immune system. Inflammatory cells include include B lymphocytes, T lymphocytes, macrophages, basophils, eosinophils, mast cells, NK cells and monocytes.

15 An "extracellular activity that can prevent the infection of a target cell by a pathogen" is any activity that can block or impede infection of a target cell by a pathogen by acting at or near the exterior surface of a target cell. An extracellular activity that can prevent the infection of a target cell by a pathogen, can be an activity such as, but not limited to, a catalytic activity or an inhibitory activity. For example, a catalytic activity
20 can be an enzymatic activity that degrades one or more entities (such as but not limited to ligands, receptors, or enzymes) on a pathogen, on a target cell, or in the vicinity of a target cell, in which the one or more entities contribute to the infection process. A catalytic activity can also modify one or more entities on a pathogen, on a target cell, or in the vicinity of a target cell, such that the infection-promoting property of the entity is
25 reduced. An inhibitory activity can be an activity that, for example, binds to a receptor or ligand and prevents the receptor or ligand from binding a moiety, where the binding is necessary for or promotes the infection process. An inhibitory activity can also be an inhibitor of an enzyme or receptor that prevents the enzyme or receptor from performing a function that is necessary for or promotes the infection process. The exterior of a target
30 cell includes the target cell membrane itself, as well as the extracellular milieu surrounding the target cell, including extracellular matrix, intracellular spaces, and

luminal spaces. For epithelial cells, the exterior of a target cell also includes the apical or luminal surface of the cell membrane that form luminal linings, and the extracellular milieu near the luminal surface. An "extracellular activity that can prevent the infection of a target cell by a pathogen" can be any type of chemical entity, including a protein, polypeptide, peptide, nucleic acid, peptide nucleic acid, nucleic acid analogue, nucleotide, nucleotide analogue, small organic molecule, polymer, lipids, steroid, fatty acid, carbohydrate, and the like, including combinations of any of these. Preferably, however, the activity comprises a peptide or protein or coupled to a peptide or protein.

An "extracellular activity that can improve transduction efficiency, or gene transfer efficiency, by a recombinant virus" is any activity that reduces or eliminates physical or chemical barriers that impedes host cell entry by a recombinant virus by acting at or near the exterior surface of a target cell. An extracellular activity that can improve transduction efficiency, or gene transfer efficiency, by a recombinant virus can be an activity such as, but not limited to, a catalytic activity or an inhibitory activity. For example, a catalytic activity can be an enzymatic activity that degrades one or more entities (such as but not limited to ligands, receptors, or enzymes) on a pathogen, on a target cell, or in the vicinity of a target cell, in which the one or more entities contribute to the infection process. A catalytic activity can also modify one or more entities on a pathogen, on a target cell, or in the vicinity of a target cell, such that the infection-promoting property of the entity is reduced. An inhibitory activity can be an activity that, for example, binds to a receptor or ligand and prevents the receptor or ligand from binding a moiety, where the binding is necessary for or promotes the infection process. An inhibitory activity can also be an inhibitor of an enzyme or receptor that prevents the enzyme or receptor from performing a function that is necessary for or promotes the infection process. The exterior of a target cell includes the target cell membrane itself, as well as the extracellular milieu surrounding the target cell, including extracellular matrix, intracellular spaces, and luminal spaces. For epithelial cells, the exterior of a target cell also includes the apical or luminal surface of the cell membrane that form luminal linings, and the extracellular milieu near the luminal surface. An "extracellular activity that can prevent the infection of a target cell by a pathogen" can be any type of chemical entity, including a protein, polypeptide, peptide, nucleic acid, peptide nucleic acid,

nucleic acid analogue, nucleotide, nucleotide analogue, small organic molecule, polymer, lipids, steroid, fatty acid, carbohydrate, and the like, including combinations of any of these. Preferably, however, the activity comprises a peptide or protein or coupled to a peptide or protein.

5

An “extracellular activity that can inhibit adhesion or function of inflammatory cells” is any activity that can prevent inflammatory cells from contacting the target cell and affecting the normal physiological status of the target cell.

10 A “domain that can anchor said at least one therapeutic domain to the membrane of a target cell”, also called an “extracellular anchoring domain” or simply, “anchoring domain” refers to a chemical entity that can stably bind a moiety that is at or on the exterior of a cell surface or is in close proximity to the surface of a cell. An extracellular anchoring domain can be reversibly or irreversibly linked to one or more moieties, such as, preferably, one or more therapeutic domains, and thereby cause the one or more
15 attached therapeutic moieties to be retained at or in close proximity to the exterior surface of a eukaryotic cell. Preferably, an extracellular anchoring domain binds at least one molecule on the surface of a target cell or at least one molecule found in close association with the surface of a target cell. For example, an extracellular anchoring domain can bind a molecule covalently or noncovalently associated with the cell membrane of a target
20 cell, or can bind a molecule present in the extracellular matrix surrounding a target cell. An extracellular anchoring domain preferably is a peptide, polypeptide, or protein, and can also comprise any additional type of chemical entity, including one or more additional proteins, polypeptides, or peptides, a nucleic acid, peptide nucleic acid, nucleic acid analogue, nucleotide, nucleotide analogue, small organic molecule, polymer, lipids,
25 steroid, fatty acid, carbohydrate, or a combination of any of these.

As used herein, a protein or peptide sequences is “substantially homologous” to a reference sequence when it is either identical to a reference sequence, or comprises one or more amino acid deletions, one or more additional amino acids, or more one or more conservative amino acid substitutions, and retains the same or essentially the same
30 activity as the reference sequence. Conservative substitutions may be defined as exchanges within one of the following five groups:

- I. Small, aliphatic, nonpolar or slightly polar residues: Ala, Ser, Thr, Pro, Gly
- II. Polar, negatively charged residues and their amides: Asp, Asn, Glu, Gln
- III. Polar, positively charged residues: His, Arg, Lys
- 5 IV. Large, aliphatic nonpolar residues: Met, Leu, Ile, Val, Cys
- V. Large aromatic residues: Phe, Try, Trp

Within the foregoing groups, the following substitution are considered to be "highly conservative": Asp/Glu, His/Arg/Lys, Phe/Tyr/Trp, and Met/Leu/Ile/Val. Semi-conservative substitutions are defined to be exchanges between two of groups (I)-(IV) above which are limited to supergroup (A), comprising (I), (II), and (III) above, or to supergroup (B), comprising (IV) and (V) above. In addition, where hydrophobic amino acids are specified in the application, they refer to the amino acids Ala, Gly, Pro, Met, Leu, Ile, Val, Cys, Phe, and Trp, whereas hydrophilic amino acids refer to Ser, Thr, Asp, Asn, Glu, Gln, His, Arg, Lys, and Tyr.

15 A "sialidase" is an enzyme that can remove a sialic acid residue from a substrate molecule. The sialidases (N-acylneuraminosylglycohydrolases, EC 3.2.1.18) are a group of enzymes that hydrolytically remove sialic acid residues from sialo-glycoconjugates. Sialic acids are alpha-keto acids with 9-carbon backbones that are usually found at the outermost positions of the oligosaccharide chains that are attached to glycoproteins and glycolipids. One of the major types of sialic acids is N-acetylneuraminic acid (Neu5Ac), which is the biosynthetic precursor for most of the other types. The substrate molecule can be, as nonlimiting examples, an oligosaccharide, a polysaccharide, a glycoprotein, a ganglioside, or a synthetic molecule. For example, a sialidase can cleave bonds having alpha(2,3)-Gal, alpha(2,6)-Gal, or alpha(2,8)-Gal linkages between a sialic acid residue and the remainder of a substrate molecule. A sialidase can also cleave any or all of the linkages between the sialic acid residue and the remainder of the substrate molecule. Two major linkages between Neu5Ac and the penultimate galactose residues of carbohydrate side chains are found in nature, Neu5Ac alpha (2,3)-Gal and Neu5Ac alpha (2,6)-Gal. Both Neu5Ac alpha (2,3)-Gal and Neu5Ac alpha (2,6)-Gal molecules can be recognized by influenza viruses as the receptor, although human viruses seem to prefer Neu5Ac alpha (2,6)-Gal, avian and equine viruses predominantly recognize Neu5Ac alpha (2,3)-

Gal. A sialidase can be a naturally-occurring sialidase, an engineered sialidase (such as, but not limited to a sialidase whose amino acid sequence is based on the sequence of a naturally-occurring sialidase, including a sequence that is substantially homologous to the sequence of a naturally-occurring sialidase). As used herein, "sialidase" can also mean
5 the active portion of a naturally-occurring sialidase, or a peptide or protein that comprises sequences based on the active portion of a naturally-occurring sialidase.

A "fusion protein" is a protein comprising amino acid sequences from at least two different sources. A fusion protein can comprise amino acid sequence that is derived from
10 a naturally occurring protein or is substantially homologous to all or a portion of a naturally occurring protein, and in addition can comprise from one to a very large number of amino acids that are derived from or substantially homologous to all or a portion of a different naturally occurring protein. In the alternative, a fusion protein can comprise amino acid sequence that is derived from a naturally occurring protein or is substantially
15 homologous to all or a portion of a naturally occurring protein, and in addition can comprise from one to a very large number of amino acids that are synthetic sequences.

A "sialidase catalytic domain protein" is a protein that comprises the catalytic domain of a sialidase, or an amino acid sequence that is substantially homologous to the
20 catalytic domain of a sialidase, but does not comprises the entire amino acid sequence of the sialidase the catalytic domain is derived from, wherein the sialidase catalytic domain protein retains substantially the same activity as the intact sialidase the catalytic domain is derived from. A sialidase catalytic domain protein can comprise amino acid sequences that are not derived from a sialidase, but this is not required. A sialidase catalytic domain
25 protein can comprise amino acid sequences that are derived from or substantially homologous to amino acid sequences of one or more other known proteins, or can comprise one or more amino acids that are not derived from or substantially homologous to amino acid sequences of other known proteins.

30

I. Composition for preventing or treating infection by a pathogen

The present invention includes peptide or protein-based compounds that comprise at least one domain that can anchor at least one therapeutic domain to the membrane of a eukaryotic cell and at least one therapeutic domain having an extracellular activity that can prevent the infection of a cell by a pathogen. By "peptide or protein-based" compounds, it is meant that the two major domains of the compound have an amino acid framework, in which the amino acids are joined by peptide bonds. A peptide or protein-based compound can also have other chemical compounds or groups attached to the amino acid framework or backbone, including moieties that contribute to the anchoring activity of the anchoring domain, or moieties that contribute to the infection-preventing activity or the therapeutic domain. For example, the protein-based therapeutics of the present invention can comprise compounds and molecules such as but not limited to: carbohydrates, fatty acids, lipids, steroids, nucleotides, nucleotide analogues, nucleic acid molecules, nucleic acid analogues, peptide nucleic acid molecules, small organic molecules, or even polymers. The protein-based therapeutics of the present invention can also comprise modified or non-naturally occurring amino acids. Non-amino acid portions of the compounds can serve any purpose, including but not limited to: facilitating the purification of the compound, improving the solubility or distribution or the compound (such as in a therapeutic formulation), linking domains of the compound or linking chemical moieties to the compound, contributing to the two-dimensional or three-dimensional structure of the compound, increasing the overall size of the compound, increasing the stability of the compound, and contributing to the anchoring activity or therapeutic activity of the compound.

The peptide or protein-based compounds of the present invention can also include protein or peptide sequences in addition to those that comprise anchoring domains or therapeutic domains. The additional protein sequences can serve any purpose, including but not limited to any of the purposes outlined above (facilitating the purification of the compound, improving the solubility or distribution or the compound, linking domains of the compound or linking chemical moieties to the compound, contributing to the two-dimensional or three-dimensional structure of the compound, increasing the overall size

of the compound, increasing the stability of the compound, or contributing to the anchoring activity or therapeutic activity of the compound). Preferably any additional protein or amino acid sequences are part of a single polypeptide or protein chain that includes the anchoring domain or domains and therapeutic domain or domains, but any
5 feasible arrangement of protein sequences is within the scope of the present invention.

The anchoring domain and therapeutic domain can be arranged in any appropriate way that allows the compound to bind at or near a target cell membrane such that the therapeutic domain can exhibit an extracellular activity that prevents or impedes infection of the target cell by a pathogen. The compound will preferably have at least one protein
10 or peptide-based anchoring domain and at least one peptide or protein-based therapeutic domain. In this case, the domains can be arranged linearly along the peptide backbone in any order. The anchoring domain can be N-terminal to the therapeutic domain, or can be C-terminal to the therapeutic domain. It is also possible to have one or more therapeutic domains flanked by at least one anchoring domain on each end. Alternatively, one or
15 more anchoring domains can be flanked by at least one therapeutic domain on each end. Chemical, or preferably, peptide, linkers can optionally be used to join some or all of the domains of a compound.

It is also possible to have the domains in a nonlinear, branched arrangement. For example, the therapeutic domain can be attached to a derivatized side chain of an amino
20 acid that is part of a polypeptide chain that also includes, or is linked to, the anchoring domain.

A compound of the present invention can have more than one anchoring domain. In cases in which a compound has more than one anchoring domain, the anchoring domains can be the same or different. A compound of the present invention can have
25 more than one therapeutic domain. In cases in which a compound has more than one therapeutic domain, the therapeutic domains can be the same or different. Where a compound comprises multiple anchoring domains, the anchoring domains can be arranged in tandem (with or without linkers) or on alternate sides of other domains, such as therapeutic domains. Where a compound comprises multiple therapeutic domains, the
30 therapeutic domains can be arranged in tandem (with or without linkers) or on alternate sides of other domains, such as, but not limited to, anchoring domains.

A peptide or protein-based compound of the present invention can be made by any appropriate way, including purifying naturally occurring proteins, optionally proteolytically cleaving the proteins to obtain the desired functional domains, and conjugating the functional domains to other functional domains. Peptides can also be chemically synthesized, and optionally chemically conjugated to other peptides or chemical moieties. Preferably, however, a peptide or protein-based compound of the present invention is made by engineering a nucleic acid construct to encode at least one anchoring domain and at least one therapeutic domain together (with or without nucleic acid linkers) in a continuous polypeptide. The nucleic acid constructs, preferably having appropriate expression sequences, can be transfected into prokaryotic or eukaryotic cells, and the therapeutic protein-based compound can be expressed by the cells and purified. Any desired chemical moieties can optionally be conjugated to the peptide or protein-based compound after purification. In some cases, cell lines can be chosen for expressing the protein-based therapeutic for their ability to perform desirable post-translational modifications (such as, but not limited to glycosylation).

A great variety of constructs can be designed and their protein products tested for desirable activities (such as, for example, binding activity of an anchoring domain, or a binding, catalytic, or inhibitory activity of a therapeutic domain). The protein products of nucleic acid constructs can also be tested for their efficacy in preventing or impeding infection of a target cell by a pathogen. *In vitro* and *in vivo* tests for the infectivity of pathogens are known in the art, such as those described in the Examples for the infectivity of influenza virus.

Anchoring Domain

As used herein, an "extracellular anchoring domain" or "anchoring domain" is any moiety that can stably bind an entity that is at or on the exterior surface of a target cell or is in close proximity to the exterior surface of a target cell. An anchoring domain serves to retain a compound of the present invention at or near the external surface of a target cell.

An extracellular anchoring domain preferably binds 1) a molecule expressed on the surface of a target cell, or a moiety, domain, or epitope of a molecule expressed on

the surface of a target cell, 2) a chemical entity attached to a molecule expressed on the surface of a target cell, or 3) a molecule of the extracellular matrix surrounding a target cell.

5 An anchoring domain is preferably a peptide or protein domain (including a modified or derivatized peptide or protein domain), or comprises a moiety coupled to a peptide or protein. A moiety coupled to a peptide or protein can be any type of molecule that can contribute to the binding of the anchoring domain to an entity at or near the target cell surface, and is preferably an organic molecule, such as, for example, nucleic acid, peptide nucleic acid, nucleic acid analogue, nucleotide, nucleotide analogue, small
10 organic molecule, polymer, lipids, steroid, fatty acid, carbohydrate, or any combination of any of these.

A molecule, complex, domain, or epitope that is bound by an anchoring domain may or may not be specific for the target cell. For example, an anchoring domain may bind an epitope present on molecules on or in close proximity to the target cell and that
15 occur at sites other than the vicinity of the target cell as well. In many cases, however, localized delivery of a therapeutic compound of the present invention will restrict its occurrence primarily to the surface of target cells. In other cases, a molecule, complex, moiety, domain, or epitope bound by an anchoring domain may be specific to a target tissue or target cell type.

20 Target tissue or target cell type includes the sites in an animal or human body where a pathogen invades or amplifies. For example, a target cell can be an endothelial cell that can be infected by a pathogen. A composition of the present invention can comprise an anchoring domain that can bind a cell surface epitope, for example, that is specific for the endothelial cell type. In another example, a target cell can be an epithelial
25 cell and a composition of the present invention can bind an epitope present on the cell surface of many epithelial cell types, or present in the extracellular matrix of different types of epithelial cells. In this case localized delivery of the composition can restrict its localization to the site of the epithelial cells that are targets of the pathogen.

A compound for preventing or treating infection by a pathogen can comprise an
30 anchoring domain that can bind at or near the surface of epithelial cells. For example, heparan sulfate, closely related to heparin, is a type of glycosaminoglycan (GAG) that is

ubiquitously present on cell membranes, including the surface of respiratory epithelium. Many proteins specifically bind to heparin/heparan sulfate, and the GAG-binding sequences in these proteins have been identified (Meyer, FA, King, M and Gelman, RA. (1975) *Biochimica et Biophysica Acta* 392: 223-232; Schauer, S. ed., pp233. Sialic Acids Chemistry, Metabolism and Function. Springer-Verlag, 1982). For example, the GAG-binding sequences of human platelet factor 4 (PF4) (SEQ ID NO:2), human interleukin 8 (IL8) (SEQ ID NO:3), human antithrombin III (AT III) (SEQ ID NO:4), human apoprotein E (ApoE) (SEQ ID NO:5), human angio-associated migratory cell protein (AAMP) (SEQ ID NO:6), or human amphiregulin (SEQ ID NO:7) (Figure 2) have been shown to have very high affinity (in the nanomolar range) towards heparin (Lee, MK and Lander, AD. (1991) *Proc Natl Acad Sci USA* 88:2768-2772; Goger, B, Halden, Y, Rek, A, Mosl, R, Pye, D, Gallagher, J and Kungl, AJ. (2002) *Biochem.* 41:1640-1646; Witt, DP and Lander AD (1994) *Curr Bio* 4:394-400; Weisgraber, KH, Rall, SC, Mahley, RW, Milne, RW and Marcel, Y. (1986) *J Bio Chem* 261:2068-2076). The GAG-binding sequences of these proteins are distinct from their receptor-binding sequences, so they will not induce the biological activities associated with the full-length proteins or the receptor-binding domains. These sequences, or other sequences that have been identified or are identified in the future as heparin/heparan sulfate binding sequences, or sequences substantially homologous to identified heparin/heparan sulfate binding sequences that have heparin/heparan sulfate binding activity, can be used as epithelium-anchoring-domains in compounds of the present invention that can be used to prevent or treat, for example, respiratory epithelium-infecting viruses such as, but not limited to, influenza virus.

An anchoring domain can bind a moiety that is specific to the target cell type of a particular species or can bind a moiety that is found in the target cell type of more than one species. In cases where the anchoring domain can bind moieties that are present at the surface of target cells of more than one species, and a virus or pathogen can infect more than one species, a therapeutic compound can have utility for more than one species (providing that the therapeutic domain is also effective across the relevant species.) For example, in the case of therapeutic compounds that can be used against influenza virus, a therapeutic compound of the present invention that has an anchoring domain that binds

heparin/heparan sulfate, the compound can be used in mammals (including humans) as well as avians.

Therapeutic Domain

5 A compound of the present invention includes at least one therapeutic domain that has an extracellular activity that can prevent or impede the infection of a cell by a pathogen, can modulate the immune response of a subject, or can improve transduction efficiency of a recombinant virus. The therapeutic activity can be, as nonlimiting examples, a binding activity, a catalytic activity, or an inhibitory activity. In some
10 embodiments of the present invention, the therapeutic activity acts to modify or inhibit a function of the pathogen that contributes to infectivity of the cell by the pathogen. In other embodiments, a therapeutic domain can modify or inhibit a function of the target cell or target organism.

 For example, the therapeutic domain can bind a receptor on a target cell that is
15 necessary for binding of the pathogen to a target cell. In this way the therapeutic moiety can block binding of the pathogen to a target cell and prevent infection. In an alternative, a therapeutic domain can bind a molecule or epitope on a pathogen to prevent an interaction of the molecule or epitope with a target cell that is necessary for infection. A therapeutic domain can also have a catalytic activity that can degrade a molecule or
20 epitope of the pathogen or host that allows for or promotes infection of a target cell by a host. In yet other embodiments, a therapeutic domain can be an inhibitor of an activity that is necessary for target cell infection by a pathogen. The inhibited activity can be an activity of the host organism or of the pathogen.

 The therapeutic domain preferably acts extracellularly, meaning that its infection-preventing, inflammatory response-modulating, or transduction-enhancing activity takes
25 place at the target cell surface or in the immediate area surrounding the target cell, including sites within the extracellular matrix, intracellular spaces, or luminal spaces of tissues.

 A therapeutic domain is preferably a peptide or protein domain (including a
30 modified or derivatized peptide or protein domain), or comprises a moiety coupled to a peptide or protein. A moiety coupled to a peptide or protein can be any type of molecule

that can prevent or impede the infection of a target cell by a pathogen, and is preferably an organic molecule, such as, for example, nucleic acid, peptide nucleic acid, nucleic acid analogue, nucleotide, nucleotide analogue, small organic molecule, polymer, lipids, steroid, fatty acid, carbohydrate, or any combination of any of these.

5 A therapeutic domain can be a synthetic peptide or polypeptide, or can comprise a synthetic molecule that can be conjugated to a peptide or polypeptide, can be a naturally-occurring peptide or protein, or a domain of naturally-occurring protein. A therapeutic domain can also be a peptide or protein that is substantially homologous to a naturally-occurring peptide or protein.

10 A therapeutic domain can have utility in a particular species, or can prevent or impede pathogen infection in more than one species. For example, therapeutic domains that inhibit pathogen functions can in general be used in a range of species that can be infected by the host, while therapeutic domains that interrupt host-pathogen interactions by interfering with a property of the host may or may not be species-specific. In many
15 cases, anchoring domains and therapeutic domains can be effective in more than one species, so that compounds of the present invention can be used to advance human and animal health, while reducing propagation and spread of the virus through animal hosts. For example, when the therapeutic domain is a sialidase, a sialidase that can cleave more than one type of linkage between a sialic acid residue and the remainder of a substrate
20 molecule, in particular, a sialidase that can cleave both alpha(2, 6)-Gal and alpha (2, 3)-Gal linkages, can protect humans from infections by a broad-spectrum of influenza viruses, including viruses that are naturally hosted in different species such as birds, pigs or horses.

25 *Linkers*

 A compound of the present invention can optionally include one or more linkers that can join domains of the compound. Linkers can be used to provide optimal spacing or folding of the domains of a compound. The domains of a compound joined by linkers can be therapeutic domains, anchoring domains, or any other domains or moieties of the
30 compound that provide additional functions such as enhancing compound stability, facilitating purification, etc. A linker used to join domains of compounds of the present

invention can be a chemical linker or an amino acid or peptide linker. Where a compound comprises more than one linker, the linkers can be the same or different. Where a compound comprises more than one linker, the linkers can be of the same or different lengths.

5 Many chemical linkers of various compositions, polarity, reactivity, length, flexibility, and cleavability are known in the art of organic chemistry. Preferred linkers of the present invention include amino acid or peptide linkers. Peptide linkers are well known in the art. Preferably linkers are between one and one hundred amino acids in length, and more preferably between one and thirty amino acids in length, although
10 length is not a limitation in the linkers of the compounds of the present invention. Preferably linkers comprise amino acid sequences that do not interfere with the conformation and activity of peptides or proteins encoded by monomers of the present invention. Some preferred linkers of the present invention are those that include the amino acid glycine. For example, linkers having the sequence:
15 (GGGGS (SEQ ID NO:10))_n, where *n* is a whole number between 1 and 20, or more preferably between 1 and 12, can be used to link domains of therapeutic compounds of the present invention.

The present invention also comprises nucleic acid molecules that encode protein-based compounds of the present invention that comprise at least one therapeutic domain
20 and at least one anchoring domain. The nucleic acid molecules can have codons optimized for expression in particular cell types, such as, for example *E. coli* or human cells. The nucleic acid molecules or the present invention that encode protein-based compounds of the present invention that comprise at least one therapeutic domain and at least one anchoring domain can also comprise other nucleic acid sequences, including but
25 not limited to sequences that enhance gene expression. The nucleic acid molecules can be in vectors, such as but not limited to expression vectors.

Composition comprising at least one anchoring domain and at least one protease inhibitor

In some aspects of the present invention, a therapeutic domain that has an
30 extracellular activity that can prevent the infection of a cell by a pathogen is a protease

inhibitor. The protease inhibitor can be any type of chemical entity, such as, for example, a carbohydrate or polymer, but is preferably a protein or peptide that inhibits the activity of an enzyme. Preferably, the protease inhibitor inhibits the activity of an enzyme that at least partially processes at least one pathogen or host cell protein, where the processing of the pathogen or host cell protein is necessary for pathogen infectivity. The enzyme that can process a viral protein necessary for pathogen infectivity can be a pathogen enzyme, or an enzyme that originates from the host organism. Preferably, the processing enzyme acts at or near the target cell surface, so that a compound of the present invention that is anchored at or near the surface of a target cell can effectively inhibit the activity of the enzyme.

Compounds of the present invention that comprise protease inhibitory domains can be used to inhibit infection by any pathogen that requires a protease in its life cycle, in which the protease is active at or near the surface of the host cell. These protein-based compositions can have, for example, one of the following structures:

(Anchoring Domain)*n*-linker-(Protease Inhibitor)*n* (*n*=1,2, 3 or more)
or :
(Protease Inhibitor)*n*-linker-(Anchoring Domain)*n* (*n*=1,2,3 or more)

The protease inhibitor can be a monomeric form of a peptide or polypeptide or can be multiple copies of the same polypeptide that are either linked directly or with spacing sequence in between. Alternatively, different polypeptide-based protease inhibitors can be linked with each other, such as, for example, aprotinin linked with soybean protease inhibitor as protease inhibiting functional domains. The polypeptides or peptides can be linked directly or via a spacer composed of peptide linker sequence. The anchoring domain can be any peptide or polypeptide that can bind at or near the surface of target cells.

The protease inhibitor can be a naturally occurring protease inhibitor (or an active portion thereof) or can be an engineered protease inhibitor. A peptide protease inhibitor used in a compound of the present invention can have a sequence substantially homologous to a naturally occurring protease inhibitor, having one or more deletions,

additions, or substitutions while retaining the activity, or substantially retaining the same activity, of the naturally occurring protease inhibitor.

In one preferred embodiment of the present invention, a therapeutic compound of the present invention is for the prevention and treatment of influenza in humans, and the
5 therapeutic domain is a protein or peptide protease inhibitor that can inhibit a serine protease that can cleave the influenza virus hemagglutinin precursor protein HA0 into HA1 and HA2.

A number of serine protease inhibitors have been shown to reduce HA cleavage and influenza virus activation in cultured cells, in chicken embryos and in lungs of
10 infected mice. They include many of the commonly used trypsin inhibitors, such as: aprotinin (Zhirmov OP, Ikizler MR and Wright PF. (2002) *J Virol* 76:8682-8689), leupeptin (Zhirmov OP, Ikizler MR and Wright PF. (2002) *J Virol* 76:8682-8689; Tashiro M, Klenk HD and Rott R.(1987) *J Gen Virol* 68:2039-2043), soybean protease inhibitor (Barbey-Morel CL, Oeltmann TN, Edwards KM and Wright PF. (1987) *J Infect Dis*
15 155:667-672), e-aminocaproic acid (Zhirmov OP, Ovchartenko AV and Bukrinskaya AG. 1982. *Arch Virol* 73:263-272) and n-p-tosyl-L-lysine chloromethylketone (TLCK) (Barbey-Morel CL, Oeltmann TN, Edwards KM and Wright PF. (1987) *J Infect Dis* 155:667-672). Among these, aerosol inhalation of aprotinin has shown definitive therapeutic effects against influenza and parainfluenza bronchopneumonia in mice
20 (Zhirmov OP, Ovcharenko AV and Bukrinskaya AG. (1984) *J Gen Virol* 65:191-196; Zhirmov OP, Ovcharenko AV and Bukrinskaya AG. (1985) *J Gen Virol* 66:1633-1638; Zhirmov OP. (1987) *J Med Virol* 21:161-167; Ovcharenko AV and Zhirmov OP. (1994) *Antiviral Res* 23:107-118) as well as in human (Zhirmov OP. (1983) *Problems Virol.* 4:9-12 (in Russian)).

25 Aprotinin (SEQ ID NO: 1; Figure 1) is a 58 amino acid polypeptide inhibitor (also called Trasylol or bovine pancreatic trypsin inhibitor (BPTI)). A compound of the present invention can have one or more aprotinin domains; for example, a therapeutic composition of the present invention can have from one to six aprotinin polypeptides, more preferably from one to three aprotinin polypeptides. A compound of the present
30 invention can also have a therapeutic domain comprising a polypeptide or peptide having substantial homology to the amino acid sequence of aprotinin.

A compound for preventing or treating influenza that comprises a protease inhibitor preferably comprises an anchoring domain that can bind at or near the surface of epithelial cells. In some preferred embodiments, the epithelium anchoring domain is a GAG-binding sequence from a human protein, such as, for example, the GAG-binding
5 sequence of human platelet factor 4 (PF4) (SEQ ID NO:2), human interleukin 8 (IL8) (SEQ ID NO:3), human antithrombin III (AT III) (SEQ ID NO:4), human apoprotein E (ApoE) (SEQ ID NO:5), human angio-associated migratory cell protein (AAMP) (SEQ ID NO:6), or human amphiregulin (SEQ ID NO:7) (Figure 2). A compound of the present invention can also have an anchoring domain comprising a polypeptide or peptide
10 having substantial homology to the amino acid sequences of the GAG-binding domains listed in SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, and SEQ ID NO:7.

Clinically, a drug comprising aprotinin and an epithelial anchoring domain can be administered by aerosol inhalation to cover the entire respiratory tract to prevent and treat
15 bronchopneumonia caused by influenza viruses, or any other virus, such as parainfluenza virus, that requires serine proteases in its life cycle. Alternatively, an aprotinin/epithelial anchoring domain fusion protein can be administered as nasal spray to treat uncomplicated early stage influenza cases or other infections by respiratory viruses. In addition, an aprotinin/epithelial anchoring domain fusion protein can be used as a
20 prophylaxis for influenza or other viral infections before an infection occurs.

Composition comprising at least one anchoring domain and at least one catalytic activity

In some aspects of the present invention, a therapeutic domain that has an extracellular activity that can prevent the infection of a cell by a pathogen is a catalytic
25 activity. The enzymatic activity can be a catalytic activity that removes, degrades or modifies a host molecule or complex or a pathogen molecule or complex that contributes to the infectivity of the pathogen. Preferably the host molecule or complex or pathogen molecule or complex that is removed, degraded, or modified by the enzymatic activity of a compound of the present invention is on, at, or near the surface of a target cell, so that a
30 compound of the present invention that is anchored to the surface of a target cell can effectively inhibit the host or pathogen molecule or complex.

For example, a therapeutic domain can have a catalytic activity that can digest a molecule or epitope of the pathogen or target cell that is required for host-pathogen binding, and subsequent entry of the pathogen into the target cell. Receptors on target cells that allow for the entry of viruses into cells can be the target of an enzymatic activity of a compound of the present invention.

Compounds of the present invention that comprise catalytic domains can be used to inhibit infection by any pathogen that uses a receptor to gain entry to a target cell, as long as removal of the receptor does not impair the organism. These protein-based compositions can have, for example, one of the following structures:

10

(Anchoring Domain)_n-[linker]-(Enzymatic Activity)_n (n=1,2, 3 or more)

or :

(Enzymatic Activity)_n (n=1,2, 3 or more)-[linker]-(Anchoring Domain)_n,

15

where the linkers are optional.

The enzymatic activity can be a monomeric form of a peptide or polypeptide or can be multiple copies of the same polypeptide that are either linked directly or with spacing sequence in between. The polypeptides or peptides can be linked directly or via a spacer composed of peptide linker sequence. The anchoring domain can be any peptide or polypeptide that can bind to or near the surface of target cells.

In one preferred embodiment of the present invention, a therapeutic domain comprises a sialidase that can eliminate or greatly reduce the level of sialic acid on the surface of epithelial cells. Sialic acid is a receptor for influenza viruses. Thus, treating the surface of respiratory epithelial cells with a sialidase can prevent influenza infections or interrupt early infections. The therapeutic domain can comprise a complete sialidase protein, or an active portion thereof. . Sialic acid is a receptor for influenza viruses, and at least one of the receptors for parainfluenza virus, some coronavirus and rotavirus, *Streptococcus pneumoniae*, *Mycoplasma pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis*, *Pseudomonas aeruginosa*, and *Helicobacter pylori*. Thus, treating the surface of respiratory epithelial cells with a sialidase can prevent influenza or other viral

infections or interrupt early infections, as well as prevent or reduce colonization of bacteria such as *Streptococcus pneumoniae*, *Mycoplasma pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis*, and *Pseudomonas aeruginosa*. Treating the gastrointestinal epithelial cells with a sialidase can prevent or reduce colonization of

5 *Helicobacter pylori* in the stomach.

Sialic acid also mediates cell adhesion and interactions between inflammatory cells and target cells. Therefore, treating the surface of respiratory epithelial cells with a sialidase can prevent the recruitment of inflammatory cells to the airway surface, and therefore can treat allergic reactions including asthma and allergic rhinitis.

10 Since sialic acid serves as a barrier that hinder cell entry by a gene therapy vector, treating the target cells with a sialidase can increase transduction efficiency, and therefore improve efficacy of the gene therapy.

Preferred sialidases are the large bacterial sialidases that can degrade the receptor sialic acids Neu5Ac alpha(2,6)-Gal and Neu5Ac alpha(2,3)-Gal. For example, the

15 bacterial sialidase enzymes from *Clostridium perfringens* (Genbank Accession Number X87369), *Actinomyces viscosus* (Genbank Accession Number X62276), *Arthrobacter ureafaciens*, or *Micromonospora viridifaciens* (Genbank Accession Number D01045) can be used. Therapeutic domains of compounds of the present invention can comprise all or a portion of the amino acid sequence of a large bacterial sialidase or can comprise amino

20 acid sequences that are substantially homologous to all or a portion of the amino acid sequence of a large bacterial sialidase. In one preferred embodiment, a therapeutic domain comprises a sialidase encoded by *Actinomyces viscosus*, such as that of **SEQ ID NO:12**, or such as sialidase sequence substantially homologous to **SEQ ID NO:12**. In yet another preferred embodiment, a therapeutic domain comprises the catalytic domain of

25 the *Actinomyces viscosus* sialidase extending from amino acids 274-666 of **SEQ ID NO:12**, or a substantially homologous sequence.

Other preferred sialidases are the human sialidases such as those encoded by the genes NEU2 (**SEQ ID NO:8**; Genbank Accession Number Y16535; Monti, E, Preti, Rossi, E., Ballabio, A and Borsani G. (1999) *Genomics* 57:137-143) and NEU4 (**SEQ ID**

30 **NO:9**; Genbank Accession Number NM080741; Monti, E, Preti, A, Venerando, B and Borsani, G. (2002) *Neurochem Res* 27:646-663) (**Figure 3**). Therapeutic domains of

compounds of the present invention can comprise all or a portion of the amino acid sequences of a human sialidase or can comprise amino acid sequences that are substantially homologous to all or a portion of the amino acid sequences of a human sialidase. Preferably, where a therapeutic domain comprises a portion of the amino acid sequences of a naturally occurring sialidase, or sequences substantially homologous to a portion of the amino acid sequences of a naturally occurring sialidase, the portion comprises essentially the same activity as the human sialidase.

A compound for preventing or treating influenza that comprises an enzymatic domain preferably comprises an anchoring domain that can bind at or near the surface of epithelial cells. In some preferred embodiments, the epithelium-anchoring domain is a GAG-binding sequence from a human protein, such as, for example, the GAG-binding amino acid sequences of human platelet factor 4 (PF4) (SEQ ID NO:2), human interleukin 8 (IL8) (SEQ ID NO:3), human antithrombin III (AT III) (SEQ ID NO:4), human apoprotein E (ApoE) (SEQ ID NO:5), human angio-associated migratory cell protein (AAMP) (SEQ ID NO:6), and human amphiregulin (SEQ ID NO:7) (Figure 2). An epithelial anchoring domain can also be substantially homologous to a naturally occurring GAG-binding sequence, such as those listed in Figure 2.

It is also within the scope of the present invention to use compounds comprising a human sialidase, or comprising a sialidase with substantial homology to a sialidase, in the absence of an anchoring domain, in the treatment or prevention of pathogen infections, such as but not limited to influenza, paramyxovirus, coronavirus, rotavirus, and *Pseudomonas aeruginosa* infections or bacterial infections; in the treatment or prevention of allergic and inflammatory responses, and to improve the transduction efficiency of a recombinant virus.

The present invention recognizes that such infections may be prevented or abated by the use of sialidases, such as, but not limited to, the *A. viscosus* sialidase or human sialidases such as NEU2 and NEU4. The sialidases can optionally be adapted, by genetic or chemical engineering, or by pharmaceutical formulation, to improve their half life or retention at the respiratory epithelium.

Because influenza viruses primarily infect the upper respiratory tract, removing the receptor sialic acid locally in the nasal cavity and nasopharynx area can prevent infections or interrupt early infections. The sialidase can be delivered to the upper respiratory tract as a nasal spray, and it can be used either in therapeutic mode during
5 early stage of influenza (or other infection) or in prophylactic mode before the infection occurs. Alternatively, it can be delivered to the lower respiratory tract as an inhalant to treat influenza and to prevent influenza complications, such as bronchopneumonia.

10 II. Therapeutic Composition Comprising at least one Sialidase Activity

The present invention includes a therapeutic composition that comprises at least one sialidase activity. The sialidase activity can be a sialidase isolated from any source, such as, for example, a bacterial or mammalian source, or can be a recombinant protein that is substantially homologous to at least a portion of a naturally occurring sialidase.
15 Preferred sialidases are the large bacterial sialidases that can degrade the receptor sialic acids Neu5Ac alpha(2,6)-Gal and Neu5Ac alpha(2,3)-Gal. For example, the bacterial sialidase enzymes from *Clostridium perfringens* (Genbank Accession Number X87369), *Actinomyces viscosus* (Genbank Accession Number L06898), *Arthrobacter ureafaciens*, or *Micromonospora viridifaciens* (Genbank Accession Number D01045) or substantially
20 homologous proteins can be used.

For example, therapeutic compounds of the present invention can comprise a large bacterial sialidase or can comprise a protein with the amino acid sequence of a large bacterial sialidase or can comprise amino acid sequences that are substantially homologous to the amino acid sequence of a large bacterial sialidase. A preferred
25 pharmaceutical composition of the present invention comprises the *A. viscosus* sialidase (SEQ ID NO:12), or comprises a protein substantially homologous to the *A. viscosus* sialidase.

Other preferred sialidases are the human sialidases such as those encoded by the genes NEU2 (SEQ ID NO:8; Genbank Accession Number Y16535; Monti, E, Preti, Rossi, E., Ballabio, A and Borsani G. (1999) *Genomics* 57:137-143) and NEU4 (SEQ ID
30 NO:9; Genbank Accession Number NM080741; Monti, E, Preti, A, Venerando, B and

Borsani, G. (2002) *Neurochem Res* 27:646-663) (Figure 3). Therapeutic domains of compounds of the present invention can comprise a human sialidase protein that is substantially homologous to the amino acid sequences of a human sialidase or can comprise amino acid sequences that are substantially homologous to all or a portion of the amino acid sequences of a human sialidase. Preferably, where a therapeutic domain comprises a portion of the amino acid sequences of a naturally occurring sialidase, or sequences substantially homologous to a portion of the amino acid sequences of a naturally occurring sialidase, the portion comprises essentially the same activity as the human sialidase.

10 A pharmaceutical composition comprising a sialidase can include other compounds, including but not limited to other proteins, that can also have therapeutic activity. A pharmaceutical composition comprising a sialidase can include other compounds that can enhance the stability, solubility, packaging, delivery, consistency, taste, or fragrance of the composition.

15 A pharmaceutical composition comprising a sialidase can be formulated for nasal, tracheal, bronchial, oral, or topical administration, or can be formulated as an injectable solution or as eyedrops. A pharmaceutical composition comprising a sialidase can be used to treat or prevent pathogen infection, to treat or prevent allergy or inflammatory response, or to enhance the transduction efficiency of a recombinant virus for gene
20 therapy.

III. Sialidase Catalytic Domain Proteins

25 The present invention also includes sialidase catalytic domain proteins. As used herein a "sialidase catalytic domain protein" comprises a catalytic domain of a sialidase but does not comprise the entire amino acid sequence of the sialidase from which the catalytic domain is derived. A sialidase catalytic domain protein has sialidase activity. Preferably, a sialidase catalytic domain protein comprises at least 10%, at least 20%, at
30 least 50%, at least 70% of the activity of the sialidase from which the catalytic domain sequence is derived. More preferably, a sialidase catalytic domain protein comprises at

least 90% of the activity of the sialidase from which the catalytic domain sequence is derived.

A sialidase catalytic domain protein can include other amino acid sequences, such as but not limited to additional sialidase sequences, sequences derived from other proteins, or sequences that are not derived from sequences of naturally-occurring proteins. Additional amino acid sequences can perform any of a number of functions, including contributing other activities to the catalytic domain protein, enhancing the expression, processing, folding, or stability of the sialidase catalytic domain protein, or even providing a desirable size or spacing of the protein.

10 A preferred sialidase catalytic domain protein is a protein that comprises the catalytic domain of the *A. viscosus* sialidase. Preferably, an *A. viscosus* sialidase catalytic domain protein comprises amino acids 270-666 of the *A. viscosus* sialidase sequence (SEQ ID NO:12). Preferably, an *A. viscosus* sialidase catalytic domain protein comprises an amino acid sequence that begins at any of the amino acids from amino acid 270 to
15 amino acid 290 of the *A. viscosus* sialidase sequence (SEQ ID NO:12) and ends at any of the amino acids from amino acid 665 to amino acid 901 of said *A. viscosus* sialidase sequence (SEQ ID NO:12), and lacks any *A. viscosus* sialidase protein sequence extending from amino acid 1 to amino acid 269. (As used herein "lacks any *A. viscosus* sialidase protein sequence extending from amino acid 1 to amino acid 269" means lacks
20 any stretch of four or more consecutive amino acids as they appear in the designated protein or amino acid sequence.)

In some preferred embodiments, an *A. viscosus* sialidase catalytic domain protein comprises amino acids 274-681 of the *A. viscosus* sialidase sequence (SEQ ID NO:12) and lacks other *A. viscosus* sialidase sequence. In some preferred embodiments, an *A. viscosus* sialidase catalytic domain protein comprises amino acids 274-666 of the *A. viscosus* sialidase sequence (SEQ ID NO:12) and lacks any other *A. viscosus* sialidase sequence. In some preferred embodiments, an *A. viscosus* sialidase catalytic domain protein comprises amino acids 290-666 of the *A. viscosus* sialidase sequence (SEQ ID NO:12) and lacks any other *A. viscosus* sialidase sequence. In yet other preferred
25
30 embodiments, an *A. viscosus* sialidase catalytic domain protein comprises amino acids

290-681 of the *A. viscosus* sialidase sequence (SEQ ID NO:12) and lacks any other *A. viscosus* sialidase sequence.

The present invention also comprises nucleic acid molecules that encode protein-based compounds of the present invention that comprise a catalytic domain of a sialidase.

- 5 The nucleic acid molecules can have codons optimized for expression in particular cell types, such as, for example *E. coli* or human cells. The nucleic acid molecules or the present invention that encode protein-based compounds of the present invention that comprise at least one catalytic domain of a sialidase can also comprise other nucleic acid sequences, including but not limited to sequences that enhance gene expression. The
- 10 nucleic acid molecules can be in vectors, such as but not limited to expression vectors.

Fusion Proteins

- Sialidase catalytic domain proteins can be fusion proteins, in which the fusion protein comprises at least one sialidase catalytic domain and at least one other protein
- 15 domain, including but not limited to: a purification domain, a protein tag, a protein stability domain, a solubility domain, a protein size-increasing domain, a protein folding domain, a protein localization domain, an anchoring domain, an N-terminal domain, a C-terminal domain, a catalytic activity domain, a binding domain, or a catalytic activity-enhancing domain. Preferably, the at least one other protein domain is derived from
- 20 another source, such as, but not limited to, sequences from another protein. The at least one other protein domain need not be based on any known protein sequence, but can be engineered and empirically tested to perform any function in the fusion protein.

- Purification domains can include, as nonlimiting examples, one or more of a his tag, a calmodulin binding domain, a maltose binding protein domain, a streptavidin
- 25 domain, a streptavidin binding domain, an intein domain, or a chitin binding domain. Protein tags can comprise sequences that can be used for antibody detection of proteins, such as, for example, the myc tag, the hemagglutinin tag, or the FLAG tag. Protein domains that enhance protein expression, modification, folding, stability, size, or localization can be based on sequences of known proteins or engineered. Other protein
- 30 domains can have binding or catalytic activity or enhance the catalytic activity of the sialidase catalytic domain.

Preferred fusion proteins of the present invention comprise at least one sialidase catalytic domain and at least one anchoring domain. Preferred anchoring domains include GAG-binding domains, such as the GAG-binding domain or human amphiregulin (SEQ ID NO:7).

5 Sialidase catalytic domains and other domains of a fusion protein of the present invention can optionally be joined by linkers, such as but not limited to peptide linkers. A variety of peptide linkers are known in the art. A preferred linker is a peptide linker comprising glycine, such as G-G-G-G-S (SEQ ID NO:10).

10 The present invention also comprises nucleic acid molecules that fusion proteins of the present invention that comprise a catalytic domain of a sialidase. The nucleic acid molecules can have codons optimized for expression in particular cell types, such as, for example E. coli or human cells. The nucleic acid molecules or the present invention that encode fusion proteins of the present invention can also comprise other nucleic acid sequences, including but not limited to sequences that enhance gene expression. The
15 nucleic acid molecules can be in vectors, such as but not limited to expression vectors.

IV Pharmaceutical Compositions

20 The present invention includes compounds of the present invention formulated as pharmaceutical compositions. The pharmaceutical compositions comprise a pharmaceutically acceptable carrier prepared for storage and preferably subsequent administration, which have a pharmaceutically effective amount of the compound in a pharmaceutically acceptable carrier or diluent. Acceptable carriers or diluents for therapeutic use are well known in the
25 pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, 18th Ed., Mack Publishing Co., Easton, PA (1990)). Preservatives, stabilizers, dyes and even flavoring agents can be provided in the pharmaceutical composition. For example, sodium benzoate, sorbic acid and esters of p-hydroxybenzoic acid can be added as preservatives. In addition,
30 antioxidants and suspending agents can be used.

Depending on the target cell, the compounds of the present invention can be formulated and used as tablets, capsules or elixirs for oral administration; salves or ointments for topical application; suppositories for rectal administration; sterile solutions, suspensions, and the like for use as inhalants or nasal sprays.

5 Injectables can also be prepared in conventional forms either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Suitable excipients are, for example, water, saline, dextrose, mannitol, lactose, lecithin, albumin, sodium glutamate, cysteine hydrochloride and the like. In addition, if desired, the injectable pharmaceutical
10 compositions can contain minor amounts of nontoxic auxiliary substances, such as wetting agents, pH buffering agents and the like.

The pharmaceutically effective amount of a test compound required as a dose will depend on the route of administration, the type of animal or patient being treated, and the physical characteristics of the specific animal under
15 consideration. The dose can be tailored to achieve a desired effect, but will depend on such factors as weight, diet, concurrent medication and other factors which those skilled in the medical arts will recognize. In practicing the methods of the present invention, the pharmaceutical compositions can be used alone or in combination with one another, or in combination with other therapeutic or
20 diagnostic agents. These products can be utilized *in vivo*, preferably in a mammalian patient, preferably in a human, or *in vitro*. In employing them *in vivo*, the pharmaceutical compositions can be administered to the patient in a variety of ways, including topically, parenterally, intravenously, subcutaneously, intramuscularly, colonically, rectally, nasally or intraperitoneally, employing a
25 variety of dosage forms. Such methods can also be used in testing the activity of test compounds *in vivo*.

In preferred embodiments, these pharmaceutical compositions may be in the form of orally-administrable suspensions, solutions, tablets or lozenges; nasal sprays; inhalants; injectables, topical sprays, ointments, powders, or gels.

30 When administered orally as a suspension, compositions of the present invention are prepared according to techniques well-known in the art of

pharmaceutical formulation and may contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners/flavoring agents known in the art. As immediate release tablets, these compositions may contain

5 microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants known in the art. Components in the formulation of a mouthwash or rinse include antimicrobials, surfactants, cosurfactants, oils, water and other additives such as sweeteners/flavoring agents known in the art.

10 When administered by a drinking solution, the composition comprises one or more of the compounds of the present invention, dissolved in water, with appropriate pH adjustment, and with carrier. The compound may be dissolved in distilled water, tap water, spring water, and the like. The pH can preferably be adjusted to between about 3.5 and about 8.5. Sweeteners may be added, e.g., 1%

15 (w/v) sucrose.

Lozenges can be prepared according to U.S. Patent No. 3,439,089, herein incorporated by reference for these purposes.

When administered by nasal aerosol or inhalation, the pharmaceutical compositions are prepared according to techniques well-known in the art of

20 pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art. See, for example, Ansel, H. C. et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Sixth Ed. (1995). Preferably these

25 compositions and formulations are prepared with suitable nontoxic pharmaceutically acceptable ingredients. These ingredients are known to those skilled in the preparation of nasal dosage forms and some of these can be found in Remington's *Pharmaceutical Sciences*, 18th Ed., Mack Publishing Co., Easton, PA (1990, a standard reference in the field. The choice of suitable carriers is

30 highly dependent upon the exact nature of the nasal dosage form desired, e.g., solutions, suspensions, ointments, or gels. Nasal dosage forms generally contain

large amounts of water in addition to the active ingredient. Minor amounts of other ingredients such as pH adjusters, emulsifiers or dispersing agents, preservatives, surfactants, jelling agents, or buffering and other stabilizing and solubilizing agents may also be present. Preferably, the nasal dosage form should
5 be isotonic with nasal secretions.

Nasal formulations can be administered as drops, sprays, aerosols or by any other intranasal dosage form. Optionally, the delivery system can be a unit dose delivery system. The volume of solution or suspension delivered per dose can preferably be anywhere from about 5 to about 2000 microliters, more preferably
10 from about 10 to about 1000 microliters, and yet more preferably from about 50 to about 500 microliters. Delivery systems for these various dosage forms can be dropper bottles, plastic squeeze units, atomizers, nebulizers or pharmaceutical aerosols in either unit dose or multiple dose packages.

The formulations of this invention may be varied to include; (1) other
15 acids and bases to adjust the pH; (2) other tonicity imparting agents such as sorbitol, glycerin and dextrose; (3) other antimicrobial preservatives such as parahydroxy benzoic acid esters, sorbate, benzoate, propionate, chlorbutanol, phenylethyl alcohol, benzalkonium chloride, and mercurials; (4) other viscosity imparting agents such as sodium carboxymethylcellulose, microcrystalline
20 cellulose, polyvinylpyrrolidone, polyvinyl alcohol and other gums; (5) suitable absorption enhancers; (6) stabilizing agents such as antioxidants, like bisulfite and ascorbate, metal chelating agents such as sodium edetate and drug solubility enhancers such as polyethylene glycols.

25

V. Method of preventing or treating infection by a pathogen

The present invention also includes methods of preventing or treating infection by a pathogen. In one aspect, the method includes: treating a subject that is infected with a
30 pathogen or at risk of being infected with a pathogen with a pharmaceutical composition of the present invention that comprises a compound that comprises at least one anchoring

domain that can anchor the compound at or near the surface of a target cell and at least one therapeutic domain comprising a peptide or protein that has at least one extracellular activity that can prevent the infection of a target cell by a pathogen. In some preferred embodiments, the method includes applying a therapeutically effective amount of a pharmaceutical composition of the present invention to epithelial cells of a subject. The subject to be treated can be an animal or human subject.

In another aspect, the method includes: treating a subject that is infected with a pathogen or at risk of being infected with a pathogen with a pharmaceutical composition of the present invention that comprises a protein-based compound that comprises a sialidase activity. In some preferred embodiments, the method includes applying a therapeutically effective amount of a pharmaceutical composition of the present invention to epithelial cells of a subject. The sialidase activity can be an isolated naturally occurring sialidase protein, or a recombinant protein substantially homologous to at least a portion of a naturally occurring sialidase. A preferred pharmaceutical composition comprises a sialidase with substantial homology to the *A. viscosus* sialidase (SEQ ID NO:12). The subject to be treated can be an animal or human subject.

In yet another aspect, the method includes: treating a subject that is infected with a pathogen or at risk of being infected with a pathogen with a pharmaceutical composition of the present invention that comprises a protein-based compound that comprises a sialidase catalytic domain. In some preferred embodiments, the method includes applying a therapeutically effective amount of a pharmaceutical composition of the present invention to epithelial cells of a subject. The sialidase catalytic domain is preferably can substantially homologous to the catalytic domain of a naturally occurring sialidase. A preferred pharmaceutical composition comprises a sialidase catalytic domain with substantial homology to amino acids 274-666 the *A. viscosus* sialidase (SEQ ID NO:12). The subject to be treated can be an animal or human subject.

A pathogen can be a viral, bacterial, or protozoan pathogen. In some embodiments, the pathogen is one of the following: influenza viruses, parainfluenza virus, respiratory syncytial virus (RSV), coronavirus, rotavirus, *Streptococcus pneumoniae*, *Mycoplasma pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis*, *Pseudomonas aeruginosa*, and *Helicobacter pylori*. In one preferred embodiment, the pathogen is influenza virus.

- Compounds of the present invention can be designed for human use or animal use. In some aspects of the present invention, a compound of the present invention can be used to prevent pathogen infection in a class of animals, such as mammals. In some aspects of the present invention, a composition can be used for human and animal use
- 5 (although the formulation may differ). In these aspects, the active domains of a compound can be effective against more than one pathogen species, type, subtype, or strain and can be active in more than one host species. For example, some preferred compounds of the present invention that comprise, for example, active domains such as
- 10 protease inhibitors that prevent processing of the HA protein of influenza virus, or sialidases that remove sialic acid receptors from target cells, or anchoring domains such as domains that bind heparin or heparan sulfate, can be used in birds, mammals, or humans. Such compounds that can be effective against a range of pathogens with the capacity to infect different host species can also be used in humans to combat infection by pathogens that are naturally hosted in other species.
- 15 In some preferred embodiments of the present invention, the pharmaceutical composition prevents infection by influenza, and a therapeutically effective amount of the pharmaceutical composition is applied to the respiratory epithelial cells of a subject. This can be done by the use of an inhaler, or by the use of a nasal spray. Preferably, the inhaler or nasal spray is used from one to four times a day.
- 20 Because influenza viruses primarily infect the upper respiratory tract, removing the receptor sialic acid locally in the nasal cavity, pharynx, trachea and bronchi can prevent infections or interrupt early infections. The sialidase can be delivered to the upper respiratory tract as a nasal spray or as an inhalant, and it can be used either in therapeutic mode during early stage of influenza (or other infection) or in prophylactic mode before
- 25 the infection occurs. Alternatively, it can be delivered to the lower respiratory tract as an inhalant to treat influenza and to prevent influenza complications, such as bronchopneumonia. Similarly, the sialidase can be delivered as nasal spray or inhalant to prevent or reduce infection by parainfluenza virus and coronavirus. It can also be delivered as an inhalant or nasal spray to prevent or reduce airway colonization by
- 30 pathogenic bacteria, including *Streptococcus pneumoniae*, *Mycoplasma pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis* and *Pseudomonas aeruginosa*. The

therapeutic compounds can optionally be adapted, by genetic or chemical engineering, or by pharmaceutical formulation, to improve their half-life or retention at the respiratory epithelium. Additionally, it can be delivered topically to the eyes or to surgical wounds in the form of drops, sprays or ointments to prevent and treat bacterial infection including
5 infection by *Pseudomonas aeruginosa*. It can also be administered orally to treat infection by *Helicobacter pylori*.

Dosage

As will be readily apparent to one skilled in the art, the useful *in vivo*
10 dosage to be administered and the particular mode of administration will vary depending upon the age, weight and type of patient being treated, the particular pharmaceutical composition employed, and the specific use for which the pharmaceutical composition is employed. The determination of effective dosage levels, that is the dose levels necessary to achieve the desired result, can be
15 accomplished by one skilled in the art using routine methods as discussed above. In non-human animal studies, applications of the pharmaceutical compositions are commenced at higher dose levels, with the dosage being decreased until the desired effect is no longer achieved or adverse side effects are reduced or disappear. The dosage for a compound of the present invention can range broadly
20 depending upon the desired affects, the therapeutic indication, route of administration and purity and activity of the compound. Typically, human clinical applications of products are commenced at lower dosage levels, with dosage level being increased until the desired effect is achieved. Alternatively, acceptable *in vitro* studies can be used to establish useful doses and routes of administration of
25 the test compound. Typically, dosages can be between about 1 ng/kg and about 10 mg/kg, preferably between about 10 ng/kg and about 1 mg/kg, and more preferably between about 100 ng/kg and about 100 micrograms/kg.

The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition (see, Fingle et al., in
30 The Pharmacological Basis of Therapeutics (1975)). It should be noted that the attending physician would know how to and when to terminate, interrupt or adjust

administration due to toxicity, organ dysfunction or other adverse effects.

Conversely, the attending physician would also know to adjust treatment to higher levels if the clinical response were not adequate. The magnitude of an administered dose in the management of the disorder of interest will vary with the severity of the condition to be treated and to the route of administration. The severity of the condition may, for example, be evaluated, in part, by standard prognostic evaluation methods. Further, the dose and perhaps dose frequency, will also vary according to the age, body weight and response of the individual patient, including those for veterinary applications.

Thus, in accordance with the present invention, there is further provided a method of treating and a pharmaceutical composition for treating influenza virus infection and prevention of influenza virus infection. The treatment involves administering to a patient in need of such treatment a pharmaceutical carrier and a therapeutically effective amount of any composition of the present invention, or a pharmaceutically acceptable salt thereof.

In one preferred regimen, appropriate dosages are administered to each patient by either inhaler, nasal spray, or by oral lozenge. It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific salt or other form employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

25

VI. Method of reducing, preventing, or treating allergic and inflammatory responses

The present invention also includes methods of reducing, preventing, or treating an allergic or inflammatory response of a subject.

30

In one aspect, the method includes: preventing or treating an allergic or inflammatory response of a subject with a pharmaceutical composition of the present

invention that comprises a protein-based compound that comprises a sialidase activity. In some preferred embodiments, the method includes applying a therapeutically effective amount of a pharmaceutical composition of the present invention to epithelial cells of a subject. The sialidase activity can be an isolated naturally occurring sialidase protein, or a
5 recombinant protein substantially homologous to at least a portion of a naturally occurring sialidase. A preferred pharmaceutical composition comprises a sialidase with substantial homology to the *A. viscosus* sialidase (SEQ ID NO:12). The subject to be treated can be an animal or human subject.

In yet another aspect, the method includes: preventing or treating an allergic or
10 inflammatory response of a subject with a pharmaceutical composition of the present invention that comprises a protein-based compound that comprises a sialidase catalytic domain. In some preferred embodiments, the method includes applying a therapeutically effective amount of a pharmaceutical composition of the present invention to epithelial cells of a subject. The sialidase catalytic domain is preferably can substantially
15 homologous to the catalytic domain of a naturally occurring sialidase. A preferred pharmaceutical composition comprises a sialidase catalytic domain with substantial homology to amino acids 274-666 the *A. viscosus* sialidase (SEQ ID NO:12). The subject to be treated can be an animal or human subject.

The allergic or inflammatory response can be and acute or chronic condition, and
20 can include, as nonlimiting examples, asthma, other allergic responses causing respiratory distress, allergic rhinitis, eczema, psoriasis, reactions to plant or animal toxins, or autoimmune conditions.

In some preferred embodiments, compounds of the present invention can be delivered as an inhalant or nasal spray to prevent or treat inflammation in the airway
25 including, but not limited to, asthma and allergic rhinitis. Compounds of the present invention comprising sialidase activity (including sialidase catalytic domain proteins and sialidase fusion proteins) can also be administered as eye drops, ear drops, or sprays, ointments, lotions, or gels to be applied to the skin. In another aspect, the method includes treating a patient who has inflammatory diseases with the present invention that comprises
30 a sialidase activity that is administered intravenously or as a local injection.

Dosage

As will be readily apparent to one skilled in the art, the useful *in vivo* dosage to be administered and the particular mode of administration will vary depending upon the age, weight and type of patient being treated, the particular pharmaceutical composition employed, and the specific use for which the pharmaceutical composition is employed. The determination of effective dosage levels, that is the dose levels necessary to achieve the desired result, can be accomplished by one skilled in the art using routine methods as discussed above. In non-human animal studies, applications of the pharmaceutical compositions are commenced at higher dose levels, with the dosage being decreased until the desired effect is no longer achieved or adverse side effects are reduced or disappear. The dosage for a compound of the present invention can range broadly depending upon the desired affects, the therapeutic indication, route of administration and purity and activity of the compound. Typically, human clinical applications of products are commenced at lower dosage levels, with dosage level being increased until the desired effect is achieved. Alternatively, acceptable *in vitro* studies can be used to establish useful doses and routes of administration of the test compound. Typically, dosages can be between about 1 ng/kg and about 10 mg/kg, preferably between about 10 ng/kg and about 1 mg/kg, and more preferably between about 100 ng/kg and about 100 micrograms/kg.

The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition (see, Fingle et al., in *The Pharmacological Basis of Therapeutics* (1975)). It should be noted that the attending physician would know how to and when to terminate, interrupt or adjust administration due to toxicity, organ dysfunction or other adverse effects. Conversely, the attending physician would also know to adjust treatment to higher levels if the clinical response were not adequate. The magnitude of an administered dose in the management of the disorder of interest will vary with the severity of the condition to be treated and to the route of administration. The severity of the condition may, for example, be evaluated, in part, by standard prognostic evaluation methods. Further, the dose and perhaps dose frequency,

will also vary according to the age, body weight and response of the individual patient, including those for veterinary applications.

In some preferred regimens, appropriate dosages are administered to each patient by either inhaler, nasal spray, or by topical application. It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific salt or other form employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

VI. Method of enhancing gene delivery by a recombinant viral vector

The present invention also includes methods of gene delivery by a recombinant viral vector. In one aspect, the method includes: administering an effective amount of a compound of the present invention that comprises a protein having sialidase activity to at least one cell prior to or concomitant with the administration of at least one recombinant viral vector. A composition of the present invention can be provided in the same formulation as at least one recombinant viral vector, or in a separate formulation.

In some preferred embodiments, the method includes applying a therapeutically effective amount of a composition of the present invention and a recombinant viral vector to cells of a subject. The subject to be treated can be an animal or human subject. In a particularly preferred embodiment, a recombinant viral vector is used to transduce epithelial target cells of a subject for gene therapy. For example, a recombinant viral vector can be used to transduce airway epithelial cells of a subject with cystic fibrosis. In this case, a compound of the present invention can be administered by use of an inhaler. A recombinant virus comprising a therapeutic gene can be administered concurrently or separately.

In other embodiments, cells can be treated with a compound of the present invention and a recombinant viral vector in vitro or "ex vivo" (that is, cells removed from a subject to be transplanted into a subject after transduction).

The sialidase activity can be an isolated naturally occurring sialidase protein, or a recombinant protein substantially homologous to at least a portion of a naturally occurring sialidase, including a sialidase catalytic domain. A preferred pharmaceutical composition comprises a sialidase with substantial homology to the *A. viscosus* sialidase (SEQ ID NO:12).

A compound of the present invention can be administered to target cells from one day before to two hours subsequent to the administration of the recombinant virus. Preferably a compound of the present invention is administered to target cells from four hours to ten minutes before administration of the recombinant virus. Administration can be

A recombinant virus is preferably a recombinant virus that can be used to transfer genes to mammalian cells, such as, preferably human cells. For example, a recombinant virus can be a retrovirus (including lentivirus), adeno-virus, adeno-associated virus (AAV) or herpes simplex virus type 1. The recombinant virus comprises at least one exogenous gene that is to be transferred to a target cell. The gene is preferably a therapeutic gene, but this need not be the case. For example, the gene can be a gene used to mark cells or confer drug resistance.

In a preferred embodiment, the present invention includes methods of improving efficacy of a gene therapy vector. The method includes treating a patient with a compound of the present invention that comprises a sialidase activity and, in the same or a separate formation, with a recombinant virus. The compound of the present invention having sialidase activity can be administered to the patient prior to, concomitant to, or even subsequent to the administration of a recombinant virus. In one embodiment, the sialidase is substantially homologous to the *Actinomyces viscosus* sialidase (SEQ ID NO:12) or a portion thereof. In one preferred embodiment, the sialidase comprises the catalytic domain of the *Actinomyces viscosus* sialidase. In another embodiment, the recombinant virus is AAV. In yet another embodiment, the disease is cystic fibrosis. In yet another embodiment, the recombinant virus comprises the cystic fibrosis transmembrane conductance regulator (CFTR) gene.

30

Dosage

As will be readily apparent to one skilled in the art, the useful *in vivo* dosage to be administered and the particular mode of administration will vary depending upon the age, weight and type of patient being treated, the particular pharmaceutical composition employed, and the specific use for which the pharmaceutical composition is employed. The determination of effective dosage levels, that is the dose levels necessary to achieve the desired result, can be accomplished by one skilled in the art using routine methods as discussed above. In non-human animal studies, applications of the pharmaceutical compositions are commenced at higher dose levels, with the dosage being decreased until the desired effect is no longer achieved or adverse side effects are reduced or disappear. The dosage for a compound of the present invention can range broadly depending upon the desired affects, the therapeutic indication, route of administration and purity and activity of the compound. Typically, human clinical applications of products are commenced at lower dosage levels, with dosage level being increased until the desired effect is achieved. Alternatively, acceptable *in vitro* studies can be used to establish useful doses and routes of administration of the test compound. Typically, dosages can be between about 1 ng/kg and about 10 mg/kg, preferably between about 10 ng/kg and about 1 mg/kg, and more preferably between about 100 ng/kg and about 100 micrograms/kg.

The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition (see, Fingle et al., in *The Pharmacological Basis of Therapeutics* (1975)). It should be noted that the attending physician would know how to and when to terminate, interrupt or adjust administration due to toxicity, organ dysfunction or other adverse effects. Conversely, the attending physician would also know to adjust treatment to higher levels if the clinical response were not adequate. The magnitude of an administered dose in the management of the disorder of interest will vary with the severity of the condition to be treated and to the route of administration. The severity of the condition may, for example, be evaluated, in part, by standard prognostic evaluation methods. Further, the dose and perhaps dose frequency,

will also vary according to the age, body weight and response of the individual patient, including those for veterinary applications.

In some preferred regimens, appropriate dosages are administered to each patient by either inhaler, nasal spray, or by topical application. It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific salt or other form employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

Examples

15

Example 1: Synthesizing aprotinin genes, purifying and testing aprotinin fusion proteins.

Introduction

Influenza viral protein hemagglutinin (HA) is the major influenza envelope protein. It plays an essential role in viral infection. The importance of HA is evidenced by the fact that it is the major target for protective neutralizing antibodies produced by the host immune response (Hayden, FG. (1996) In *Antiviral drug resistance* (ed. D. D. Richman), pp. 59-77. Chichester, UK: John Wiley & Sons Ltd.). It is now clear that HA has two different functions in viral infection. First, HA is responsible for the attachment of the virus to sialic acid cell receptors. Second, HA mediates viral entry into target cells by triggering fusion of the viral envelope with cellular membranes.

HA is synthesized as a precursor protein, HA0, which is transferred through the Golgi apparatus to the cell surface as a trimeric molecular complex. HA0 is further cleaved to generate the C terminus HA1 (residue 328 of HA0) and the N terminus of HA2. It is generally believed that the cleavage occurs at the cell surface or on released viruses. The cleavage of HA0 into HA1/HA2 is not required for HA binding to a sialic acid receptor; however, it is essential for viral infectivity (Klenk, HD and Rott, R. (1988)

Adv Vir Res. 34:247-281; Kido, H, Niwa, Y, Beppu, Y and Towatari, T. (1996) *Advan Enzyme Regul* 36:325-347; Skehel, JJ and Wiley, DC. (2000) *Annu Rev Biochem* 69:531-569).

Sensitivity of HA0 to host proteases is determined by the proteolytic site in the external loop of HA0 molecule. The proteolytic site may contain either a single Arg or Lys residue (monobasic cleavage site) or several Lys and/or Arg residues in R-X-K/R-R motif (multibasic cleavage site). Only the influenza A virus subtypes H5 and H7 have HA proteins carrying the multibasic cleavage site. All other influenza A, B and C viruses contain HA proteins having the monobasic cleavage site. Influenza A viruses having multibasic cleavage sites are more virulent and induce systemic infection in hosts whereas viruses with a monobasic HA site initiate infection only in the respiratory tract in mammals or in the respiratory and enteric tracts in avian species (Klenk, HD and Garten W. 1994. *Trend Micro* 2:39-43 for review). Fortunately, human infection by the highly virulent avian influenza A H5 and H7 subtypes, which carry the multibasic cleavage site, has so far only occurred in a handful of cases discovered mostly in Hong Kong. The vast majority of influenza infections are caused by viruses with HA proteins are cleaved at the monobasic cleavage site.

Influenza virus HA subtypes 5 and 7 that contain multibasic cleavage sites are activated by furin, a member of the subtilisin-like endoproteases, or the pre-protein convertase family. Furin cleaves the virus intracellularly and is ubiquitously present in many cell types, allowing the virulent, systemic infection seen with such viruses (Klenk, HD and Garten W. 1994. *Trend Micro* 2:39-43; Nakayama, K. 1997. *Biochem* 327:625-635). All other influenza viruses, which have HAs with monobasic cleavage sites, are activated by secreted, trypsin-like serine proteases. Enzymes that have been implicated in influenza virus activation include: plasmin (Lazarowitz SG, Goldberg AR and Choppin PW. 1973. *Virology* 56:172-180), mini-plasmin (Murakami M, Towatari T, Ohuchi M, Shiota M, Akao M, Okumura Y, Parry MA and Kido H. (2001) *Eur J Biochem* 268: 2847-2855), tryptase Clara (Kido H, Chen Y and Murakami M. (1999) In B.Dunn (ed.), *Proteases of infectious agents*: p.205-217, Academic Press, New York, N.Y), kallikrein, urokinase, thrombin (Scheiblaue H, Reinacher M, Tashiro M and Rott R. (1992) *J Infec Dis* 166:783-791), blood clotting factor Xa (Gotoh B, Ogasawara T, Toyoda T, Inocencio

- N, Hamaguchi M and Nagai Y. (1990) *EMBO J* 9:4189-4195), acrosin (Garten W, Bosch FX, Linder D, Rott R and Klenk HD. (1981) *Virology* 115:361-374.), proteases from human respiratory lavage (Barbey-Morel CL, Oeltmann TN, Edwards KM and Wright PF. (1987) *J Infect Dis* 155:667-672) and bacterial proteases from *Staphylococcus aureus* (Tashiro M, Ciborowski P, Reinacher M, Pulverer G, Klenk HD and Rott R. (1987) *Virology* 157:421-430) and *Pseudomonas aeruginosa* (Callan RJ, Hartmann FA, West SE and Hinshaw VS. (1997) *J Virol* 71:7579-7585). Activation of influenza viruses by host serine proteases is generally considered to occur extracellularly either at the plasma membrane or after virus release from the cell.
- 10 Aprotinin, also called Trasylol, or bovine pancreatic trypsin inhibitor (BPTI) is a polypeptide having 58 amino acids. It belongs to the family of Kunitz-type inhibitors and competitively inhibits a wide spectrum of serine proteases, including trypsin, chymotrypsin, plasmin and plasma kallikrein. Aprotinin has long been used as a human therapeutics, such as treatment of pancreatitis, various states of shock syndrome,
- 15 hyperfibrinolytic haemorrhage and myocardial infarction. It is also used in open-heart surgery, including cardiopulmonary bypass operations, to reduce blood loss (Fritz H and Wunderer G. (1983) *Arzneim-Forsch* 33:479-494).
- The safety of aprotinin in human has been well documented through years of clinical applications. In addition, aprotinin is apparently a very weak immunogen as
- 20 aprotinin-specific antibodies have not been observed in human sera so far (Fritz H and Wunderer G. (1983) *Arzneim-Forsch* 33:479-494). Another desired feature of aprotinin as a drug candidate is its superb stability. It can be kept at room temperature for at least 18 months without any loss of activity (Fritz H and Wunderer G. (1983) *Arzneim-Forsch* 33:479-494).
- 25 To achieve significant viral inhibition in animal studies that have been performed, aprotinin was administered at high doses. For example, 280 micrograms to 840 micrograms per day of aprotinin was injected intraperitoneally into each mouse for 6 days (Zhirnov OP, Ovcharenko AV and Bukrinskaya AG. (1984) *J Gen Virol* 65:191-196); a lower dosage was required for aerosol inhalation, still, each mouse was given 63 -
- 30 126 micrograms per day for 6 days (Ovcharenko AV and Zhirnov OP. (1994) *Antiviral Res* 23:107-118). A very high dose of aprotinin would be required in human based on

extrapolation from the mouse data. Therefore to achieve better efficacy in human, the potency of aprotinin molecule needs to be significantly improved.

Aprotinin functions by competitively inhibiting serine proteases that are mostly on the surface of host respiratory epithelial cells. Local concentration of aprotinin in the vicinity of host proteases is therefore the key factor determining competitive advantage of aprotinin. We use two approaches that work synergistically to boost competitive advantage of aprotinin on the surface of respiratory epithelium.

First, the avidity (functional affinity) of aprotinin is increased by making multivalent aprotinin fusion proteins consisting of two, three, or more aprotinin proteins connected via linkers. Such a molecule is able to bind to membrane proteases in a multivalent fashion, which has significant kinetic advantage over the aprotinin monomer. Monomeric aprotinin binds to bovine trypsin very tightly with dissociation constant (K_i) being 6.0×10^{-14} mol/l. However, its affinity compared to other proteases, such as chymotrypsin, plasmin and Kallikrein, which have been implicated in activation of influenza viruses, is much lower with K_i being at the level of 10^{-8} to 10^{-9} mol/l (Fritz H and Wunderer G. (1983) *Arzneim-Forsch* 33:479-494). Multimerization can increase aprotinin's affinity to these proteases exponentially.

Second, we fuse aprotinin with a respiratory epithelium-anchoring domain. The anchoring domain localizes aprotinin to the proximity of host membrane-associated proteases and maintains a high local concentration of aprotinin on epithelial surface. The anchoring domain also increases retention time of the drug on the respiratory epithelium.

Cloning

Aprotinin is a single chain polypeptide having 58 amino acid residues and 3 intra-chain disulfide bonds (SEQ ID NO:1). The amino acid sequence of aprotinin is shown in **Figure 1**. Genes encoding aprotinin and aprotinin fusion proteins are synthesized by PCR using overlapping oligonucleotides with codons optimized for *E. Coli* expression as templates. The PCR products are cloned into pCR2.1-TOPO vector (Invitrogen). After sequencing, the genes are subcloned into an expression vector pQE (Qiagen). The vector carries a purification tag, Hisx6, to allow easy purification of the recombinant proteins. The constructs are used to transform *E. Coli*. The transformed cells grown in LB-

ampicillin medium to mid-log phase are induced by IPTG according to standard protocols. Cells are pelleted and lysed in phosphate-buffered-saline (PBS) by sonication. The enzymes, which have His₆ purification tag, are purified using a nickel column (Qiagen).

5 The following aprotinin fusion proteins are made:

1. Dimeric and trimeric aprotinin. Two or three aprotinin genes are linked via a flexible linker as the following constructs:

Aprotinin—(GGGGS (SEQ ID NO:10))_n (n=3, 4 or 5)—Aprotinin;

10 and

Aprotinin—(GGGGS(SEQ ID NO:10))_n (n=3, 4 or 5)—Aprotinin—
(GGGGS(SEQ ID NO:10))_n (n=3, 4 or 5)—Aprotinin

The length of the linker sequence may determine three-dimensional flexibility of the multimeric aprotinin and thereby influence functional affinity of the molecule. Therefore
15 constructs having linkers with various lengths are made.

Fully functional recombinant monomeric aprotinin has been produced in *E. Coli* (Auerswald EA, Horlein D, Reinhardt G, Schroder W and Schnabel E. (1988). *Biol Chem Hoppe-Seyler* Vol 369, Suppl., pp27-35). We therefore expect proper folding of multivalent aprotinin proteins in *E. coli* cells. Besides expressing protein in various
20 common *E. Coli* cell strains, such as BL21, JM83, etc, the multivalent aprotinin proteins are also expressed in OrigamiTM cells (Novagen, Bad Soden, Germany). The OrigamiTM cell strain does not have thioredoxin and glutathione reductase and thus has an oxidizing cytoplasm. This cell strain has been used to successfully express a number of proteins that contain disulfide bonds (Besette PH, Aslund F, Beckwith J and Georgiou G. (1999)
25 *Pro Natl Acad Sci USA* 96:13703-13708; Venturi M, Seifert C and Hunte C. (2001) *J Mol Biol* 315:1-8.).

2. The epithelium cell-anchoring aprotinin. An epithelium cell-anchoring sequence is fused with aprotinin. The epithelium-anchoring sequence can be any peptide or polypeptide sequence that has affinity towards the surface of epithelial cells. We have

selected three human GAG-binding sequences: PF4 (aa 47-70; **SEQ ID NO: 2**), IL-8 (aa 46-72; **SEQ ID NO: 3**), and AT III (aa 118-151; **SEQ ID NO: 4**) (**Figure 2**).

These sequences bind to heparin/heparan sulfate with nanomolar-level affinities (**Table 1**). Heparin/Heparan Sulfate are ubiquitously present on the respiratory epithelium. In separate constructs, the GAG-binding sequences are fused with the aprotinin gene on the N terminus and on the C terminus via a generic linker sequence GGGGS as the following constructs:

(GAG domain—GGGGS(**SEQ ID NO:10**)—Aprotinin); and

(Aprotinin—GGGGS(**SEQ ID NO:10**)—GAG domain)

Table 1. Affinities to Heparin

Protein	Kd nM (ref)	
PF4	27	(44)
IL-8	<5	(43)
ATIII	11	(42)
ApoE	620	(45)

Photometric trypsin inhibition assay

The trypsin inhibition activity of aprotinin and aprotinin fusion proteins is measured by a photometric assay described previously in detail (Fritz H and Wunderer G. (1983) *Arzneim-Forsch* 33:479-494). Briefly, in this assay aprotinin inhibits the trypsin-catalyzed hydrolysis of Na-benzoyl-L-arginine-p-nitroanilide (BzArgpNA or L-BAPA)

(Sigma), which is followed photometrically at 405 nm. One trypsin unit (U_{BAPA}) corresponds to the hydrolysis of 1 micromole substrate per min. One inhibitor unit (IU_{BAPA}) decreases the activity of two trypsin units by 50%, which corresponds arithmetically to the inhibition of 1 U_{BAPA} of trypsin. The specific activity of aprotinin is
5 given in IU_{BAPA}/mg polypeptide.

Surface plasmon resonance assay

The affinities of dimeric and trimeric aprotinin with various linkers are compared against the monomeric aprotinin using surface plasmon resonance assay, or BIAcore
10 analysis (BIAcore, Piscataway, NJ) with human plasmin as the target. Similarly, BIAcore assay with heparin as the target is used to analyze affinity between GAG binding aprotinin fusion proteins and heparin.

When plasmin is used as the target, purified human plasmin (Sigma) is immobilized on the CM5 chip according manufacturer's instructions (BIAcore,
15 Piscataway, NJ). When heparin is the target, biotinylated albumin and albumin-heparin (Sigma) are captured on a streptavidin-coated BIAcore SA chip as described previously (Xiang Y and Moss B. (2003) *J Virol* 77:2623-2630).

Example 2: Establishing improved tissue culture models for studies on influenza virus infection. 20

Stocks of Influenza Viruses

Influenza viral strains are obtained from ATCC and the repository at St. Jude Children's Research Hospital. All experiments involving influenza viruses are conducted at Bio-safety level II.

25 Viruses are propagated by injection into the allantoic cavity of nine-day-old chicken embryos as described (Zhirmov OP, Ovcharenko AV and Bukrinskaya AG. (1985) *J Gen Virol* 66:1633-1638). Alternatively, viral stocks are grown on Madin-Darby canine kidney (MDCK) cells in minimal essential medium (MEM) supplemented with 0.3% bovine serum albumin and 0.5 micrograms of trypsin per ml. After

incubating for 48 to 72 hours, the culture medium is clarified by low speed centrifugation. Viral particles are pelleted by ultracentrifugation through a 25% sucrose cushion. Purified viruses are suspended in 50% glycerol-0.1M Tris buffer (pH 7.3) and stored at -20°C .

5

Plaque Assays

Infectivity and titer of the viral stocks are determined by two kinds of plaque assays, a conventional one and a modified one (Tobita, K, Sugiura, A, Enomoto, C and Furuyama, M. (1975) *Med Microbiol Immunol* 162:9-14; Zhirmov OP, Ovcharenko AV and Bukrinskaya AG. (1982) *Arch Virol* 71:177-183). The conventional plaque assay is routinely used as a virus titration method. It requires exogenous trypsin in agar overlay added immediately after virus infection to MDCK monolayers (Tobita, K, Sugiura, A, Enomoto, C and Furuyama, M. (1975) *Med Microbiol Immunol* 162:9-14). This method artificially increases infectivity of the viral stocks being tested by activating all the viral particles that have uncleaved HA.

Zhirmov et. al. designed a modified plaque assay consisting of a double agar overlay, with trypsin being included in the second layer which is added 24 hours after infection (Zhirmov OP, Ovcharenko AV and Bukrinskaya AG. (1982) *Arch Virol* 71:177-183). Three days after infection, cells are fixed with a 10% formaldehyde solution, agarose layers are removed, fixed cells are stained with hematoxylin-eosin solution and plaques are counted. The modified plaque assay allows accurate determination of the real infectivity of viral stocks that contain both cleaved and uncleaved HA. Combining results from both conventional and modified plaque assays, one can distinguish viruses containing cleaved or uncleaved HA and correlate infectivity of viral stocks with the status of HA cleavage.

25

Human Cell Culture Models

1. Short-term culture of primary human epithelial cells. Conventional *in vitro* influenza virus infection is mostly carried out in MDCK cells with exogenous trypsin added to the culture medium. This is far from being physiological and is inappropriate for the work

30

proposed here because trypsin is not the protease that activate influenza viruses *in vivo*. Very limited numbers of *in vitro* tissue culture models that are able to support the growth of influenza virus without an exogenous protease have been reported so far, those being primary cultures with primate cells of renal origin, cells lining the allantoic and amniotic cavities of embryonated eggs, fetal tracheal ring organ cultures and primary human adenoid epithelial cells (Endo Y, Carroll KN, Ikizler MR and Wright PF. (1996) *J Virol* 70:2055-2058). Among these, the latest work with primary human adenoid epithelial cells is the closest mimic of human conditions. In this case, Endo et. al. (Endo Y, Carroll KN, Ikizler MR and Wright PF. (1996) *J Virol* 70:2055-2058) isolated epithelial cells from surgical samples of human adenoids, and cultured the epithelial cells on a collagen matrix (Vitrogen 100, Celtrix Laboratories, Palo Alto, California) in Transwell inserts (Costar, Cambridge, Mass). Cells were maintained in 50% Ham's F12 and 50% Eagles minimal essential media with supplements of growth factors and trace elements. The cells reached confluency in 10 to 14 days, remaining largely as a monolayer but with discrete patches of ciliated cells, which maintained regular ciliary activity for 1 to 3 weeks after reaching confluency. In this system, influenza A virus grew to a titer of 10^6 PFU/ml with a multiplicity of infection of 0.001 (Endo Y, Carroll KN, Ikizler MR and Wright PF. (1996) *J Virol* 70:2055-2058). Progressive cytopathogenic effects were also present during infection. The biggest drawback of this system is that it requires fresh human adenoid tissue.

To solve this problem, primary human adenoid epithelial cells are replaced with primary human airway epithelial cells that are commercially available (Cambrex), and the cells are grown under the same conditions. Such short-term culture of primary human airway epithelial cells is relatively quick to establish and is useful as the first-line experimental model for most of the *in vitro* infection and antiviral experiments.

2. Well-differentiated human airway epithelium (WD-HAE). In order to best mimic the *in vivo* condition of human airway, the model of well-differentiated human airway epithelium (WD-HAE) is used. WD-HAE is stratified epithelium that has all the differentiated cells of the normal human airway epithelium, including functional ciliated cells and mucus secreting cells. Therefore, in this model system influenza viruses are

most likely to be activated by host proteases that are physiologically relevant. Although WD-HAE has been widely used to study respiratory viral infections, such as respiratory syncytial virus (RSV) Zhang L, Peebles ME, Boucher RC, Collins PL and Pickles RJ. (2002) *J Virol* 76:5654-5666) measles virus (Sinn PL, Williams G, Vongpunswad S, Cattaneo R and McCray PB. (2002) *J Virol* 76:2403-2409, or human rhinovirus, it has not previously been used to study influenza viruses.

A detailed protocol of WD-HAE has been described previously (Krunkosky TM, Fischer BM, Martin LD, Jones N, Akley NJ and Adler KB. (2000) *Am J Respir Cell Mol Biol* 22:685-692). Briefly, commercial primary human bronchial epithelial cells (Cambrex) are cultured on Transwell-clear culture inserts (Costar) that are thin-coated with rat-tail collagen I. Cells are cultured submerged for the first 5 to 7 days in medium containing a 1:1 mixture of bronchial epithelial cell growth medium (BEGM) (Cambrex) and DMEM with high glucose with supplement of growth factors (Krunkosky TM, Fischer BM, Martin LD, Jones N, Akley NJ and Adler KB. (2000) *Am J Respir Cell Mol Biol* 22:685-692). When cultures are 70% confluent (days 5 to 7), the air-liquid interface is created by removing the apical medium and exposing cells only to medium on their basal surface. Cells are cultured for additional 14 days in air-liquid interphase, for a total of 21 days in culture, and are then ready for experiments. The differentiated epithelium can be maintained *in vitro* for weeks.

Epithelial morphology and degree of differentiation is documented by routine histology (Endo Y, Carroll KN, Ikizler MR and Wright PF. (1996) *J Virol* 70:2055-2058). Briefly, following fixation with 10% buffered formalin, the epithelial cells are embedded in paraffin, sectioned and stained with hematoxylin and eosin, and with periodic acid-Schiff stain for mucus secreting cells.

Influenza infection is carried out in the above two model systems by adding 0.001 to 1 MOI of viruses to the differentiated cells. The titer and infectivity of viruses in the supernatant are followed over a period of 3 to 7 days. The level of influenza viral amplification and the infectivity of influenza viruses are evaluated using conventional and modified plaque assays.

Example 3: Comparing functions of the aprotinin fusion proteins in vitro***Anti-Viral Effects of Aprotinin Fusion Proteins***

1. Pre-infection treatment. Aprotinin fusion proteins are added to primary human cell cultures at various concentrations and allowed to incubate with the cells for 1 hour.

5 The cells are washed with fresh medium and immediately inoculated with influenza viruses at MOI 0.01 to 1. Cells are washed again after 1 hour and cultured for 3 to 5 days. Titer and infectivity of viruses in the supernatant are measured at various time points by two plaque assays. The cytopathic effect caused by viral infection is evaluated by staining viable cells with crystal violet and quantifying by measuring absorption at

10 570 nm at the end of the experiment. The percentage of cell protection by aprotinin fusion proteins is calculated by $100 \times \{(\text{aprotinin treated sample} - \text{untreated infected sample}) / (\text{uninfected control} - \text{untreated infected sample})\}$. The drug efficacy for cell protection is described by its Effective Concentration that achieves 50% of the cell protection (EC_{50}). Since HA activation only occurs to newly released viral particles, the

15 first round of viral infection occurs normally and viral titer rises in the first 24 hours after infection. However, starting from the second round, infectivity of viruses drops and viral titer gradually decreases as result of aprotinin treatment. Results from this experiment differentiate various types of different aprotinin fusion proteins by their efficacies in a single prophylactic treatment.

20 Alternatively, timing of initial viral inoculation is altered from immediately after aprotinin treatment to 2-24 hours post treatment. Viral titer, infectivity and cytopathic effect are measured for 3 to 5 day after infection as described above. Results from these experiments distinguish various aprotinin fusion proteins by the lengths of the effective window after a single prophylactic treatment.

25

2. Post-infection Treatment. For multi-dose treatment, cells are first infected by viral inoculations at 0.001 to 0.1 MOI for 1 hour. Various concentrations of aprotinin fusion proteins are added immediately afterwards, additional treatments are applied at 8-hour intervals during the first 48 hours post infection. Cells are cultured until day 7 post

infection. Viral titer and infectivity in the media are followed during the whole process. Cytopathic effect is evaluated at the end of the experiment.

For single dose treatment, cells are first infected by viral inoculations at 0.001 to 0.1 MOI for 1 hour. Treatments of aprotinin fusion proteins at various concentrations are applied at different time points during the first 48 hours after infection, but each cell sample only receives one treatment during the whole experiment. Cells are cultured until day 7 post infection. Viral titer and infectivity in the media are followed during the whole process. Cytopathic effect is evaluated at the end of the experiment. Results from these experiments distinguish different types of aprotinin fusion proteins for their therapeutic potency.

Inhibition of HA Cleavage by Aprotinin Fusion Proteins

To demonstrate that aprotinin fusion proteins inhibit influenza viral infection by inhibiting cleavage of influenza HA protein, a human primary epithelial cell culture is infected with influenza virus at MOI of 1. Aprotinin fusion proteins are added to the culture either right before viral inoculation or immediately after the viral infection. At 6.5 hour post infection, the culture is incubated for 1 hour in MEM lacking cold methionine and containing ^{35}S -labeled methionine (Amersham) at a concentration of 100 microCi/ml (pulse). Thereafter, the cells are washed twice with MEM containing a 10-fold concentration of cold methionine and incubated in MEM for additional 3 hours (chase). After labeling, cells are dissolved in radioimmunoprecipitation assay (RIPA) buffer, HA is precipitated by anti-serum against the particular strain of virus used for infection (anti-influenza sera can be obtained from ATCC and Center of Disease Control and Prevention), and immunocomplex is then purified by protein G-Sepharose (Amersham). Samples are fractionated by SDS-PAGE followed by autoradiography. In samples untreated by aprotinin fusion proteins, HA1 and HA2 are expected to be the predominant HA species; while in aprotinin treated samples, HA0 is expected to be the major type of HA present.

Example 4: Synthesizing genes of five sialidases, expressing and purifying the sialidase proteins.

Introduction

Influenza viruses belong to the *orthomyxoviridae* family of RNA viruses. Both
5 type A and type B viruses have 8 segmented negative-strand RNA genomes enclosed in a
lipid envelope derived from the host cell. The viral envelope is covered with spikes that
are composed of three proteins: hemagglutinin (HA), that attaches virus to host cell
receptors and mediates fusion of viral and cellular membranes; neuraminidase (NA),
which facilitates the release of the new viruses from the host cell; and a small number of
10 M2 proteins that serve as ion channels. For *Influenza A virus*, HA and NA both undergo
antigenic drift and antigenic shift, the viral subtypes are distinguished by serologic
differences between their HA and NA proteins. There are total 15 types of HA (H1-H15)
and 9 types of NA (N1-N9), but only three HA (H1-H3) and two NA (N1 and N2) have
been found in human *Influenza A virus* so far (Granoff, A. & Webster, R. G., ed.
15 *Encyclopedia of Virology*, 2nd Edition, Vol 2). In contrast to *Influenza A virus*, no distinct
antigenic subtypes are recognized for *Influenza virus B*.

While *Influenza B virus* circulates only in humans, *Influenza A virus* can be
isolated from a whole host of animals, such as pigs, horses, chickens, ducks and other
kinds of birds, which accounts for genetic reassortment of *Influenza A virus* that results in
20 antigenic shift. Wild aquatic birds are considered to be the primordial reservoir of all
influenza viruses for avian and mammalian species. There is extensive evidence for
transmission of the virus between aquatic birds and other species including pigs and
horses and indirect transmission to humans through pigs. Direct transmission from pigs
or chickens to humans has also been documented (Ito, T. (2000) *Microbiol Immunol*
25 44(6):423-430).

The host cell receptor for influenza viruses is the cell surface sialic acid. Sialic
acids are α -keto acids with 9-carbon backbones that are usually found at the outermost
positions of the oligosaccharide chains that are attached to glycoproteins and glycolipids.
One of the major types of sialic acids is N-acetylneuraminic acid (Neu5Ac), which is the
30 biosynthetic precursor for most of the other types. Two major linkages between Neu5Ac

and the penultimate galactose residues of carbohydrate side chains are found in nature, Neu5Ac α (2,3)-Gal and Neu5Ac α (2,6)-Gal. Both Neu5Ac α (2,3)-Gal and Neu5Ac α (2,6)-Gal molecules can be recognized by *Influenza A virus* as the receptor (Schauer, R. (1982) *Adv. Carbohydrate Chem & Biochem* 40:131-235), while human viruses seem to
5 prefer Neu5Ac α (2,6)-Gal, avian and equine viruses predominantly recognize Neu5Ac α (2,3)-Gal (Ito, T. (2000) *Microbiol Immunol* 44(6):423-430).

Infections by influenza type A and B viruses are typically initiated at the mucosal surface of the upper respiratory tract. Viral replication is primarily limited to the upper respiratory tract but can extend to the lower respiratory tract and causes
10 bronchopneumonia that can be fatal. The risk of death is one per 10,000 infections, but is significantly greater for high-risk groups with pre-existing cardiopulmonary conditions and for immunologically naïve individuals during a pandemic.

A therapeutic compound comprising a sialidase that can effectively degrade both receptor sialic acids, Neu5Ac α (2,6)-Gal and Neu5Ac α (2,3)-Gal, can confer protection
15 against the broadest range of influenza viruses, including animal viruses. It can also remain effective as the viral strains change yearly. Because sialidase targets the host cell rather than virus and acts at the "choking point" in a viral life cycle, generation of resistant virus is improbable. Protein-bound sialic acid turns over homogeneously on cell surface with half-life of 33 hours (Kreisel, W, Volk, BA, Buchsel, R. and Reutter, W.
20 (1980) *Proc Natl Acad Sci USA* 77:1828-1831). Therefore we estimate that once-a-day or twice-a-day administration of a sialidase would confer sufficient protection against influenza.

Sialidases are found in higher eukaryotes, as well as in some mostly pathogenic microbes, including viruses, bacteria and protozoans. Viral and bacterial sialidases have
25 been well characterized, and the three-dimensional structures of some of them have been determined (Crennell, SJ, Garman, E, Laver, G, Vimr, E and Taylor, G. (1994) *Structure* 2:535-544; Janakiraman, MN, White, CL, Laver, WG, Air, GM and Luo, M. (1994) *Biochemistry* 33:8172-8179; Pshezhetsky, A, Richard, C, Michaud, L, Igoudora, S, Wang, S, Elsliger, M, Qu, J, Leclerc, D, Gravel, R, Dallaire, L and Potier, M. (1997) *Nature*
30 *Genet* 15: 316-320). Several human sialidases have also been cloned in the recent years (Milner, CM, Smith, SV, Carrillo MB, Taylor, GL, Hollinshead, M and Campbell, RD.

- (1997) *J Bio Chem* 272:4549-4558; Monti, E, Preti, A, Nesti, C, Ballabio, A and Borsani G. 1999. *Glycobiol* 9:1313-1321; Wada, T, Yoshikawa, Y, Tokuyama, S, Kuwabara, M, Akita, H and Miyagi, T. (1999) *Biochem Biophy Res Communi* 261:21-27; Monti, E, Bassi, MT, Papini, N, Riboni, M, Manzoni, M, Veneranodo, B, Croci, G, Preti, A, Ballabio, A, Tettamanti, G and Borsani, G. (2000) *Bichem J* 349:343-351). All the sialidases characterized share a four amino acid motif in the amino terminal portion followed by the Asp box motif which is repeated three to five times depending on the protein. (Monti, E, Bassi, MT, Papini, N, Riboni, M, Manzoni, M, Veneranodo, B, Croci, G, Preti, A, Ballabio, A, Tettamanti, G and Borsani, G. (2000) *Bichem J* 349:343-351; Copley, RR, Russell, RB and Ponting, CP. (2001) *Protein Sci* 10:285-292). While the overall amino acid identity of the sialidase superfamily is relatively low at about 20-30%, the overall fold of the molecules, especially the catalytic amino acids, are remarkably similar (Wada, T, Yoshikawa, Y, Tokuyama, S, Kuwabara, M, Akita, H and Miyagi, T. (1999) *Biochem Biophy Res Communi* 261:21-27; Monti, E, Bassi, MT, Papini, N, Riboni, M, Manzoni, M, Veneranodo, B, Croci, G, Preti, A, Ballabio, A, Tettamanti, G and Borsani, G. (2000) *Bichem J* 349:343-351; Copley, RR, Russell, RB and Ponting, CP. (2001) *Protein Sci* 10:285-292).

The sialidases are generally divided into two families: "small" sialidases have molecular weight of about 42 kDa and do not require divalent metal ion for maximal activity; "large" sialidases have molecular weight above 65 kDa and may require divalent metal ion for activity (Wada, T, Yoshikawa, Y, Tokuyama, S, Kuwabara, M, Akita, H and Miyagi, T. (1999) *Biochem Biophy Res Communi* 261:21-27; Monti, E, Bassi, MT, Papini, N, Riboni, M, Manzoni, M, Veneranodo, B, Croci, G, Preti, A, Ballabio, A, Tettamanti, G and Borsani, G. (2000) *Bichem J* 349:343-351; Copley, RR, Russell, RB and Ponting, CP. (2001) *Protein Sci* 10:285-292).

Over fifteen sialidase proteins have been purified and they vary greatly from one another in substrate specificities and enzymatic kinetics. To confer a broad-spectrum protection against influenza viruses, a sialidase needs to effectively degrade sialic acid in both $\alpha(2,6)$ -Gal and $\alpha(2,3)$ -Gal linkages and in the context of glycoproteins and some glycolipids. Viral sialidases, such as those from *influenza A virus*, *fowl plague virus* and *Newcastle disease virus*, are generally specific for Neu5Ac $\alpha(2,3)$ -Gal and only degrade

Neu5Ac α (2,6)-Gal very inefficiently. Small bacterial sialidases generally react poorly to sialic acid in the context of glycoproteins and glycolipids. By contrast, large bacterial sialidases can effectively cleave sialic acid in both (α ,2-6) linkage and (α ,2-3) linkage in the context of most natural substrates (**Figure 4**; Vimr, DR. (1994) *Trends Microbiol* 2: 271-277; Drzeniek, R. (1973) *Histochem J* 5:271-290; Roggentin, P, Kleineidam, RG and Schauer, R. (1995) *Biol Chem Hoppe-Seyler* 376:569-575; Roggentin, P, Schauer, R, Hoyer, LL and Vimr, ER. (1993) *Mol Microb* 9:915-921). Because of their broad substrate specificities, large bacterial sialidases are better candidates.

Among the large bacterial sialidases with known substrate specificity shown in **Figure 4**, *Vibrio cholerae* sialidase requires Ca^{2+} for activity making it less preferred. More preferred sialidases include the 71 kDa enzyme from *Clostridium perfringens*, the 113 kDa enzyme from *Actinomyces viscosus* and sialidase of *Arthrobacter ureafaciens*. A third sialidase, the 68 kDa enzyme from *Micromonospora viridifaciens*, has been known to destroy influenza viral receptor (Air, GM and Laver, WG. (1995) *Virology* 211:278-284), and is also a candidate.

These enzymes have high specific activity (600 U/mg protein for *C. perfringens* (Corfield, AP, Veh, RW, Wember, M, Michalski, JC and Schauer, R. (1981) *Biochem J* 197:293-299) and 680 U/mg protein for *A. viscosus* (Teufel, M, Roggentin, P. and Schauer, R. (1989) *Biol Chem Hoppe Seyler* 370:435-443)), are fully active without divalent metal iron, and have been cloned and purified as recombinant proteins from *E. coli* (Roggentin, P, Kleineidam, RG and Schauer, R. (1995) *Biol Chem Hoppe-Seyler* 376:569-575, Teufel, M, Roggentin, P. and Schauer, R. (1989) *Biol Chem Hoppe Seyler* 370:435-443, Sakurada, K, Ohta, T and Hasegawa, M. (1992) *J Bacteriol* 174: 6896-6903). In addition, *C. perfringens* is stable in solution at 2-8°C for several weeks, and at 4°C in the presence of albumin for more than two years (Wang, FZ, Akula, SM, Pramod, NP, Zeng, L and Chandran, B. (2001) *J Virol* 75:7517-27). *A. viscosus* is labile towards freezing and thawing, but is stable at 4°C in 0.1 M acetate buffer, pH 5 (Teufel, M, Roggentin, P. and Schauer, R. (1989) *Biol Chem Hoppe Seyler* 370:435-443).

Although the chances of inducing immune reactions using bacterial sialidases is very low because the proteins will be used topically in the upper respiratory tract and will

not be absorbed systemically a human enzyme would be more desirable for long-term use in human subjects.

Four sialidase genes have been cloned from human so far: NEU1/G9/lysosomal sialidase (Pshezhetsky, A, Richard, C, Michaud, L, Igdoura, S, Wang, S, Elsliger, M, Qu, J, Leclerc, D, Gravel, R, Dallaire, L and Potier, M. (1997) *Nature Genet* 15: 316-320. 5 , Milner, CM, Smith, SV, Carrillo MB, Taylor, GL, Hollinshead, M and Campbell, RD. (1997). *J Bio Chem* 272:4549-4558); NEU3, a membrane-associated sialidase isolated from human brain (Wada, T, Yoshikawa, Y, Tokuyama, S, Kuwabara, M, Akita, H and Miyagi, T. (1999) *Biochem Biophys Res Communi* 261:21-27, Monti, E, Bassi, MT, 10 Papini, N, Riboni, M, Manzoni, M, Venerando, B, Croci, G, Preti, A, Ballabio, A, Tettamanti, G and Borsani, G. (2000) *Biochem J* 349:343-351), NEU2 a 42 kDa sialidase expressed in human skeletal muscle at a very low level (Monti, E, Preti, A, Nesti, C, Ballabio, A and Borsani G. (1999) *Glycobiol* 9:1313-1321), and NEU4 a 497 amino acid protein (Genbank NM080741) expressed in all human tissues examined (Monti, E, Preti, 15 A, Venerando, B and Borsani, G. (2002) *Neurochem Res* 27:646-663).

Amino acid sequence comparison reveals NEU2 (SEQ ID NO:8) and NEU4 (SEQ ID NO:9) are both cytosolic sialidases. 9 out of 12 of the amino acid residues which form the catalytic site of *S. typhimurium* sialidase are conserved in both NEU2 and NEU4 (Monti, E, Preti, A, Nesti, C, Ballabio, A and Borsani G. (1999) *Glycobiol* 9:1313- 20 1321, Figure 3). In addition, NEU4 also shows a stretch of about 80 amino acid residues (aa 294-373) that appears unique among known mammalian sialidases (Monti, E, Preti, A, Venerando, B and Borsani, G. (2002) *Neurochem Res* 27:646-663). Unlike the selected large bacterial sialidases, the substrate specificity of NEU2 and NEU4 is unknown. It will need to be tested if NEU2 and NEU4 can effectively degrade the 25 influenza virus receptors.

Sialidase assay

NEU2, NEU4 and *M. viridifaciens* enzymes will be stored in PBS and 50% glycerol at -20°C. *C. perfringens* and *A. viscosus* enzymes are stored in 10mM acetate 30 buffer (pH5) at 4°C. Protein preps are characterized by HPLC and SDS-PAGE

electrophoresis. Specific activities and stability of the enzymes will be monitored by sialidase assay.

The enzymatic activity of sialidases are determined by fluorimetric 2'-(4-methylumbelliferyl)-alpha-D-N-acetylneuraminic acid (4MU-NANA) (Sigma) as the substrate. Specifically, reactions are set up in duplicate in 0.1M Na citrate/phosphate buffer pH5.6, in the presence of 400 micrograms bovine serum albumin, with 0.2 mM 4MU-NANA in a final volume of 100 microliters, and incubated at 37°C for 5-10min. Reactions are stopped by addition of 1 ml of 0.2 M glycines/NaOH pH10.2. Fluorescence emission is measured on a fluorometer with excitation at 365 nm and emission at 445 nm, using 4-methylumbelliferone (4-MU) to obtain a calibration curve.

Example 5: Comparing functions of the sialidases in vitro and selecting one sialidase for further studies.

1. Stocks of Influenza Viruses

Influenza viral strains are obtained from the ATCC and the repository at St. Jude Children's Research Hospital. Viral stocks are grown on Madin-Darby canine kidney (MDCK) cells in minimal essential medium (MEM) supplemented with 0.3% bovine serum albumin and 0.5 micrograms of trypsin per ml. After incubating for 48 to 72 hours, the culture medium is clarified by low speed centrifugation. Viral particles are pelleted by ultracentrifugation through a 25% sucrose cushion. Purified viruses are suspended in 50% glycerol-0.1M Tris buffer (pH 7.3) and stored at -20°C. Viral titer is determined by plaque assay (Tobita, K, Sugiura, A, Enomoto, C and Furuyama, M. (1975) *Med Microbiol Immunol* 162: 9-14), or TCID₅₀, which is the dose of virus required to infect 50% of the MDCK cells.

Selected human and animal influenza A strains with specificity towards Neu5Ac alpha(2,6)-Gal or Neu5Ac alpha(2,3)-Gal and have high affinity to the receptors (measured by high hemagglutination activity) are chosen for *in vitro* tests:

1. Strains that recognize receptor Neu5Ac alpha(2,6)-Gal include human isolates A/aichi/2/68, A/Udon/307/72, A/Prot Chaimers/1/73 and A/Victoria/3/75, etc.

(Connor, RJ, Kawaoka, Y, Webster, RG and Paulson JC. (1994) *Virology* 205:17-23).

2. Strains that have Neu5Ac alpha(2,3)-Gal specificity include animal isolates A/duckUkraine/1/63, A/duckMemphis/928/74, A/duckhokk/5/77,
5 A/Eq/Miami/1/63, A/Eq/Ur/1/63, A/Eq/Tokyo/71, A/Eq/Prague/71, etc (Connor, RJ, Kawaoka, Y, Webster, RG and Paulson JC. (1994) *Virology* 205:17-23).

2. Hemagglutination Assay

10 This assay is used to rapidly determine the efficiency of each enzyme to destroy receptors Neu5Ac alpha(2,6)-Gal and Neu5Ac alpha(2,3)-Gal.

Specifically, 6 ml of Chicken red blood cells (SPAFAS Inc., Norwich, CT) are diluted in two times the volume of PBS, centrifuge for 5 min at 500 x g and re-suspended in PBS of original volume. Sialidases are added to the chicken erythrocytes at various concentrations and allowed to incubate at room temperature for 30 min. The cells are
15 then washed three times to remove sialidase proteins, and then are resuspended in PBS to 6 ml. Control cells are incubated with BSA and washed. Various strains of influenza virus, which recognize either Neu5Ac alpha(2,6)-Gal or Neu5Ac alpha(2,3)-Gal as the receptor as listed above, are prepared in microtiter plates as serial dilutions in PBS (100 microliters) of the original viral stocks. Sialidase-treated or control chicken red blood cell
20 suspensions (100 microliters of the 0.5% solution prepared above) are added to each well at 4°C. The plates are read after 2 h. The lowest concentration of virus that causes the blood cell to agglutinate is defined as one hemagglutination unit. We will be looking for enzymes that effectively abolish hemagglutination by all viral strains.

25 3. Viral Inhibition Assay

Confluent monolayers of MDCK cells are treated with various concentrations of sialidases for 1 h, washed twice with buffer, then infected with various strains of influenza virus. After incubation for 1 hr, the cells are washed again to remove unbound virus. To estimate the decrease in viral binding sites on cell surface, the cells are overlaid

with agar and incubated at 37°C. The number of plaques in the sialidase treated cells will be compared against those in control cells. Alternatively, the cells will be cultured in regular medium at 37°C, and viral titers in the culture media are measured at various time during culture as TCID₅₀.

5 To demonstrate that sialidase treatment can inhibit a pre-existing infection, MDCK monolayers are first infected with a low titer of virus. After washing off the unbound virus, the cells are then cultured in the presence of a sialidase. Fresh sialidase is added to cell culture every 24 h. Viral titer in the cultured medium is measured over a 72-hour period.

10 4. Cytotoxicity assay

 Primary human bronchial epithelial cells are purchased (Clonetics) and cultured in supplemented minimal medium following manufacture's instruction. Sialidases are added to the culture medium at various concentrations. Cell growth over a period of 7-10 days will be measured. Cells will also be observed regularly for microscopic cytopathic
15 effects.

Example 6: Constructing and testing sialidase fusion proteins.

1. Choosing a GAG-binding sequence as the anchoring domain.

 One sialidase is selected for its best overall properties, including anti-viral
20 activity, toxicity, stability, ease of production, etc. We will then genetically link it to a GAG-binding sequence, sub-clone the fusion genes into pQE vector, express and purify the fusion proteins from *E. coli*.

 We have selected six possible human GAG-binding sequences: PF4 (aa 47-70) (SEQ ID NO:2), IL-8 (aa 46-72) (SEQ ID NO:3), AT III (aa 118-151) (SEQ ID NO:4),
25 ApoE (aa 132-165) (SEQ ID NO:5), amphiregulin (aa 25-45) (SEQ ID NO:6), and human angio-associated migratory cell protein (AAMP) (aa 14-25) (SEQ ID NO:7) (Figure 2). These sequences generally bind to heparin with nanomolar-level affinities; however, their affinities may vary from one another by an order of magnitude (Table 1). Since it is not clear which anchoring domain will enable the most effective functioning of

the sialidase, all four GAG-binding sequences are fused with the sialidase gene either on the N terminus or the C terminus via a generic linker sequence GGGGS as the following constructs:

(GAG binding domain— GGGGS(SEQ ID NO:10) —Sialidase); or

5

(Sialidase—GGGGS(SEQ ID NO:10)—GAG binding domain)

Different fusion proteins are compared by a modified viral inhibition assay. Specifically, confluent monolayers of MDCK cells are treated with same amount of each fusion protein for a limited duration, such as 30 min. The cells are then washed twice with buffer to remove unbound sialidase fusion proteins, and incubated in culture medium for an additional 1 hour. Afterwards, strains of influenza virus are added to the cells for 1 hr and then cells are washed again to remove unbound virus. Viral titers in the culture media are measured during 72-h cultures as TCID₅₀. The un-fused sialidase protein will be used to compare against the fusion proteins in this assay. If the results are too close to rank all fusion proteins, we will make the assay more stringent by shortening treatment window for the fusion proteins, lowering protein concentrations and increasing the level of viral challenge.

2. Optimizing the fusion protein construct

After selecting the best fusion protein from the earlier experiments, the construct is further optimized by testing different linker length. In this regard, the following constructs are made:

(Sialidase— (GGGGS(SEQ ID NO:10))_n (n=0, 1, 2, 3, or 4) —GAG binding domain)

The proteins are expressed and purified and compared in the modified viral protection assay as described above.

In addition, if earlier data indicate that higher affinity of the fusion protein towards heparan sulfate brings better potency, we also plan to test if the potency can be further improved by increasing the GAG-binding affinity. This can be achieved by creating a multivalent GAG binding mechanism in the fusion protein in constructs like these:

(Sialidase—(GGGGS(SEQ ID NO:10))_n—HS binding domain—GAG binding domain);

or:

(GAG binding domain—(GGGGS(SEQ ID NO:10))_n—Sialidase—

(GGGGS(SEQ ID NO:10))_n—GAG binding domain)

- 5 The purified fusion proteins are ranked based on their activities in the modified viral protection assay as described above.

3. Cytotoxicity assay

- 10 The effects of the fusion proteins on normal cell growth and morphology are monitored by culturing primary human bronchial epithelial cells with various concentrations of the fusion proteins and following growth curve of the cells and observing any microscopic cytopathic effects.

Example 7: Fusion Proteins against Other Infectious Microbes

- 15 Fusion proteins composed of a functional domain and an anchorage domain are designed for many more different applications. For example, a sialidase fusion protein as proposed here can also be used as a therapeutic/prophylactic agent against infections by other viruses and bacteria besides influenza viruses, because many other infectious microbes, such as paramyxoviruses (Wassilewa, L. (1977) *Arch Virol* 54:299-305),
 20 coronaviruses (Vlasak, R., Luytjes, W., Spaan, W. and Palese, P. (1988) *Proc Natl Acad Sci USA* 85:4526-4529), rotaviruses (Fukudome, K., Yoshie, O. and Konno, T. (1989) *Virology* 172:196-205) and *Pseudomonas aeruginosa* (Ramphal, R. and Pyle, M. (1983) *Infect Immun* 41:339-44) etc, are also known to use sialic acid as cellular receptors. For
 25 example, aprotinin fused with a heparin-binding domain can make a fusion protein that be used to prevent/treat infection of other viruses besides influenza that require host serine proteases for activation, such as parainfluenza virus.

Example 8: Cloning Sialidase Catalytic Domain Fusion Proteins

According to the published literature on the large bacterial sialidases, the 51 kDa *Arthrobacter ureafaciens* sialidase, the 71 kDa sialidase from *Clostridium perfringens* and the 113 kDa sialidase from *Actinomyces viscosus* seem to have similar specific activities and broad substrate specificity toward various sialic acid conjugates (Biology of the Sialic Acids (1995) , 270-273; Corfield et al., Biochem. J., (1981) 197(2), 293-299; Roggentin et al., Biol. Chem. Hoppe Seyler, (1995) 376(9), 569-575; Teufel et al., Biol. Chem. Hoppe Seyler, (1989) 370(5), 435-443). A third sialidase, the 68 kDa enzyme from *Micromonospora viridifaciens*, was also known to destroy the influenza viral receptor (Air and Laver, Virology, (1995) 211(1), 278-284; (1995) , 270-273).

A. viscosus is part of the normal flora of human oral cavity and gastrointestinal tract (Sutter, Rev. Infect. Dis., (1984) 6 Suppl 1, S62-S66). Since the sialidase from *A. viscosus* is normally secreted by the bacterium hosted on human mucosal surface, it should be tolerated by the human mucosal immune system. Therefore, it is unlikely that *A. viscosus* sialidase will be immunogenic when delivered topically to the human airway surface. We think that this feature makes *A. viscosus* sialidase a good candidate for a therapeutic agent.

We determined that a fragment of the *A. viscosus* sialidase, extending from amino acid 274 to amino acid 667, should contain the catalytic domain (referred to as AvCD) of the sialidase and should be fully active on it own. We later cloned the AvCD fragment and demonstrated that this AvCD fragment and other *A. viscosus* sialidase fragments comprising at least amino acids 290-666 of the *A. viscosus* sialidase protein sequence (SEQ ID NO:12), such as the fragment extending from amino acid 274 to amino acid 681, the fragment extending from amino acid 274 to amino acid 666, the fragment extending from amino acid 290 to amino acid 666, and the fragment extending from amino acid 290 to amino acid 681, have sialidase activity.

The complete sequence of *A. viscosus* sialidase protein and gene were obtained from GenBank (A49227 and L06898). Based on homology with sialidases with known 3D structures (*M. viridifaciens* and *S. typhimurium*), we assigned the catalytic domain (CD) sequence to be located between amino acids 274-667 (SEQ ID NO:16). To clone the catalytic domain of *A. viscosus* sialidase (AvCD), this region of the *A. viscosus*

sialidase gene was engineered with codons optimized for expression in *E. coli* (SEQ ID NO:15). The codon-optimized AvCD nucleotide sequence encoding amino acids 274-667 of the *A. viscosus* sialidase (SEQ ID NO:15) was produced by chemical synthesis of overlapping oligonucleotides which were annealed, amplified by PCR and cloned into the expression vector pTrc99a (Amersham, New Jersey, USA).

Sialidase fusion constructs were made using standard molecular cloning methods. The His₆-AR construct was made by fusing six histidines (His₆) to the N-terminal residue of the AvCD sequence. The His₆-AvCD construct has the nucleotide sequence of SEQ ID NO:17 and translated amino acid sequence of SEQ ID NO:18. These sequences are depicted in Figure 7.

To make the AR-AvCD construct, an anchoring domain was directly fused with the N-terminal residue of the AvCD sequence. The anchoring domain, referred to as AR, was derived from the GAG binding sequence of human amphiregulin precursor (GenBank # AAH09799). Nucleotide sequences encoding amino acids 125 to 145 (Figure 2, SEQ ID NO:7) of the human amphiregulin precursor were synthesized chemically as two overlapping oligonucleotides. The AR-AvCD construct has the nucleotide sequence of SEQ ID NO:19 and translated amino acid sequence of SEQ ID NO:20.

Another construct, AR-G4S-AvCD, was made by fusing the same AR-encoding sequence used in the AR-AvCD construct with a sequence encoding a five-amino-acid linker (GGGGS; SEQ ID NO:10) which then was fused with the AvCD sequence such that in a translation product, the linker was fused to N-terminus of the catalytic domain of the *A. viscosus* sialidase. The nucleotide sequence (SEQ ID NO:34) and translated amino acid sequence (SEQ ID NO:35) of this construct are depicted in Figure 9. All constructs were cloned into the pTrc99a expression vector.

In addition, four constructs were made in which the catalytic domain of the *A. viscosus* sialidase was fused to the N-terminus of the AR (GAG-binding domain of human amphiregulin; SEQ ID NO:7). In Construct #4 (SEQ ID NO:27), the catalytic domain of the *A. viscosus* sialidase consisted of amino acids 274-666 of SEQ ID NO:12 fused to the GAG-binding domain of amphiregulin (SEQ ID NO:7). In Construct #5 (SEQ ID NO:29), the catalytic domain of the *A. viscosus* sialidase consisted of amino acids 274-

681 of SEQ ID NO:12 fused to the GAG-binding domain of amphiregulin (SEQ ID NO:7). In Construct #6 (SEQ ID NO:31), the catalytic domain of the *A. viscosus* sialidase consisted of amino acids 290-666 of SEQ ID NO:12 fused to the GAG-binding domain of amphiregulin (SEQ ID NO:7). In Construct #7 (SEQ ID NO:33), the catalytic domain of the *A. viscosus* sialidase consisted of amino acids 290-681 of SEQ ID NO:12 fused to the GAG-binding domain of amphiregulin (SEQ ID NO:7). All of these constructs displayed comparable sialidase activity in assays.

Example 9: Production of Sialidase Catalytic Domain Fusion Proteins

To produce the sialidase fusion proteins, the expression constructs were transformed into E.coli BL21. A single colony was inoculated into 2.5 ml of LB broth and grown overnight at 37°C with shaking. In the morning 2 ml of overnight culture was inoculated into 500 ml of TB medium in a 2 liter shake flask and the culture was allowed to grow to OD₆₀₀=4.0 (2-4 hours) at 37°C with shaking. Protein expression was induced by addition of IPTG to a final concentration of 1 mM and continued for 3 hr with shaking. Cells were harvested by centrifugation at 5,000xg for 10 min. Cell were washed once (resuspended in PBS and recentrifuged) and resuspended in 15 ml of Lysis buffer.

Compositions of media and buffers used in protein expression and purification.

TB medium for protein expression

Solution 1

Bacto-tryptone - 12 g

Yeast extract - 24 g

H₂O to 800 ml

Solution 2

KH₂PO₄ (anhydrous) - 2.3 g

K₂HPO₄ (anhydrous) - 12.5 g

H₂O to 100 ml

Autoclave solutions 1 and 2 separately, cool, mix and add the following:

60 ml of 20% glycerol (filter sterilized)

20 ml of 20% glucose (filter sterilized)

Lysis buffer

50 mM phosphate, pH 8.0

5 10% glycerol

300 mM NaCl

Bacterial cells suspended in lysis buffer were lysed by sonication and cell debris was removed by centrifugation. Clarified lysate was passed through an SP-Sepharose column (bed volume 15 ml, flow rate 120 cm/hour). The column was reconditioned to lower pH and salt with one volume of PBS to ensure good retention of Fludase during endotoxin removal. Endotoxin was removed by washing the column with 5 volumes of PBS containing 1% Triton X-100, 0.5% Sodium Deoxycholate and 0.1% SDS. The detergents were washed away with 3 volumes of PBS and 3 volumes of lysis buffer.

15 Proteins were eluted from the column with lysis buffer that contained 0.8 M NaCl. The fraction eluted from SP-Sepharose was adjusted to 1.9 M $(\text{NH}_4)_2\text{SO}_4$ (most contaminating proteins are salted out at this step) and clarified by centrifugation. The supernatant was loaded onto Butyl-Sepharose column (flow rate 120 cm/hour). The column was washed with 2 volumes of 1.3 M $(\text{NH}_4)_2\text{SO}_4$ and the fusion was eluted with 0.65 M $(\text{NH}_4)_2\text{SO}_4$.

20 For the final step, size exclusion chromatography was performed on Sephacryl S-200 equilibrated with PBS buffer at a flow rate of 25 cm/hour. Sialidase activity was determined against 4-MU-NANA as described in the following paragraph. Protein concentration was determined using Bio-Rad's Bradford kit. Protein purity was assessed by SDS-PAGE and estimated to be >98%. Specific activity of the enzyme was about 937

25 U/mg. Endotoxin in final preparations was measured using LAL test (Cambrex) and estimated to be <0.5 EU/ml.

For purification of His6 containing fusion protein, cation exchange on SP-Sepharose was replaced with Metal Chelate Affinity Chromatography on Ni-NTA. All buffers remained the same with the exception that elution from Ni-NTA was performed

30 by 0.25 M imidazole in lysis buffer.

Example 10: Sialidase Assay to Measure Activity of Sialidase Catalytic Domain Fusion Proteins

The sialidase activity of the AR-AvCD protein encoded by Construct #2 was assayed and compared with that of native sialidases purified from *C. perfringens* (Sigma, St. Louis, MO) and *A. ureafaciens* (Prozyme, San Leandro, CA). In addition, a fusion protein produced from a construct in which the amphiregulin GAG sequence (SEQ ID NO: 7) was fused to the Neu 2 human sialidase (SEQ ID NO:8) was also assayed for sialidase activity.

The sialidase activity expressed as units per mg sialidase was measured by the sialidase assay using the artificial fluorogenic substrate 4-MU-NANA (Sigma). One unit of sialidase is defined as the amount of enzyme that releases 10 nmol of MU from 4-MU-NANA in 10 min at 37°C (50 mM CH₃COOH - NaOH buffer, pH 5.5) in reaction that contains 20 nmol of 4-MU-NANA in a 0.2 ml volume. Reactions are stopped by addition of 1 ml of 0.2 M glycine/NaOH pH 10.2. Fluorescence emission is measured on a fluorometer with excitation at 365 nm and emission at 445 nm, using 4-methylumbelliferone (4-MU) to obtain a calibration curve (Potier et al., Anal. Biochem., (1979) 94(2), 287-296).

Table 2. Specific activity of sialidases (units per mg).

Sialidase	Specific activity
AR-NEU2	8
AR-AvCD	937
<i>C. perfringens</i>	333
<i>A. ureafaciens</i>	82

Our results show that the AvCD fusion protein (AR-AvCD) has the highest specific activity among all the tested sialidases (Table 2). The specific activity of AR-AvCD is over 100 times higher than that of a human sialidase fusion (AR-NEU2), and over two times higher than that of *C. perfringens* sialidase. Experimental results comparing the stability of the sialidases indicate very high stability of AR-AvCD: No

loss of activity for AR-AvCD was detected after 20 weeks at 25°C or 4°C in solution. By comparison, AR-NEU2 solution exhibited a half-life of 5 and 2 weeks when stored at 25°C and 37°C, respectively.

5 **Example 11: Optimization of the N-terminus of Sialidase Catalytic Domain Fusion Proteins**

The N-terminus of the AR-AvCD fusion protein was partially cleaved under certain conditions that resulted in small degrees of protein heterogeneity in the purified AR-AvCD prep. To solve this problem, we designed an approach to optimize the N-terminus of the sialidase fusion construct. A library containing AR-AvCD with random amino acids at the N-terminus was constructed as follows. AR-AvCD was amplified by PCR using a primer pair in which the primer annealing on 5'-end of the gene contained a randomized sequence in positions corresponding to amino acids 2 and 3. The nucleotide sequence of the primer and the encoded amino acid sequence are shown below.

15

ttttcgtctcccatgvnnvnnaagcgcaaaaaaaaaaggcggca (SEQ ID NO:21)

MetXxxXxxLysArgLysLysLysGlyGly (SEQ ID NO:22)

In SEQ ID NO:21, "n" stands for any nucleotide (a, c, g, or t) and "v" stands for nucleotides a, g or c. By designing the sequence in such a way (disallowing the nucleotide t in the first position of codons) we avoided introduction of stop codons as well as aromatic amino acids (Phe, Tyr, Trp) and Cys. The *Esp3I* restriction endonuclease site (shown in bold) was introduced to allow generation of *NcoI* compatible overhang. The primer annealing to 3'-end of the gene carried *HindIII* site following the stop codon.

20 The PCR product was digested with *Esp3I* - *HindIII* was ligated into pTrc99a expression vector digested with *NcoI* - *HindIII*. The ligation mix was transformed into *E.coli* and the cells were grown overnight in liquid culture containing Ampicillin.

The next day the culture was diluted with fresh medium, grown to OD₆₀₀=0.8 and induced with IPTG for 2 hours. Cells were harvested, homogenized and the fusions were subjected to two-step purification by liquid chromatography. Clarified lysate was loaded onto SP-Sepharose equilibrated with lysis buffer (50 mM HEPES, pH 8.0, 0.3 M NaCl,

30

10% glycerol). The column was washed with 0.45 M NaCl and the fusions were eluted with 0.9 M NaCl. The eluate was diluted with 10% glycerol to bring the concentration of NaCl to 0.2 M and loaded onto Heparin-Sepharose column. The column was developed with a linear gradient of NaCl. The fractions that contained sialidase activity were
5 resolved on SDS-PAGE, electroblotted onto PVDF membrane and the 43 kDa band was subjected to amino-terminal sequencing.

The predominant N-terminal residues of the isolated sialidase fusion protein were either Val or Gly followed by the N-terminal residues of the AR tag. We then synthesized new sialidase fusion constructs, Constructs #2 and #3, by introducing a Val
10 in front of the AR sequence such that the first six amino acids encoded by Constructs #2 and #3 were (Met-Val-Lys-Arg-Lys-Lys (SEQ ID NO:23)). N-terminal sequencing of proteins made from these new fusion constructs showed 100% homogeneity with the initiation Met being completely removed (which is desirable for therapeutic proteins) and Val being the first N-terminal residue followed by the AR tag sequence. These data are
15 consistent with earlier publications that reported the common rules of N-terminal processing and protein stability as function of protein's N-terminal amino acid residue (Hirel et al., Proc. Natl. Acad. Sci. U. S. A, (1989) 86(21), 8247-8251; Varshavsky, Proc. Natl. Acad. Sci. U. S. A, (1996) 93(22), 12142-12149).

The nucleotide sequences of new fusion Construct #2 (AR-AvCD with optimized
20 N-terminus) (SEQ ID NO:24) and its amino acid sequence translation (SEQ ID NO:25) is depicted in Figure 10. The nucleotide sequences of new fusion Construct #3 (AR-G4S-AvCD with optimized N-terminus) (SEQ ID NO:36) and its amino acid sequence translation (SEQ ID NO:37) is depicted in Figure 11. The amino acid sequence of processed proteins isolated from E. coli infected with Construct #2 is provided herein as
25 SEQ ID NO:38 and the amino acid sequence of processed proteins isolated from E. coli infected with Construct #3 is provided herein as SEQ ID NO:39.

Example 12: Comparing Activities of Sialidase Constructs with or without an Anchoring Domain

30 To evaluate if the AR sequence indeed improves the cell-surface activity of a sialidase fusion protein, we incubated proteins purified from E. coli that were

transformed with Construct #2; SEQ ID NO:24, depicted in Figure 7) or Construct #1 (His₆-AvCD; SEQ ID NO:17, depicted in Figure 5) with primary human bronchial epithelial cells and measured cell-bound sialidase activity after extensive washing. For cells incubated with Construct #2 protein (SEQ ID NO:25), up to 10% of the sialidase was found to be cell-bound, and the cell-bound sialidase activity increased in a dose-dependent manner with the input concentration of Construct #2 protein. However, Construct #1 protein (SEQ ID NO:18) incubated cells only exhibited background level of sialidase activity. Furthermore, we treated MDCK cells with either Construct #2 protein or Construct #1 protein and measured the level of residual α (2,6)-linked sialic acid on the surface of the cells (Figure 8). At equal levels of enzymatic activity below 100 mU per well, Construct #2 protein demonstrated significantly higher potency than Construct #1 protein. These results indicate that the AR domain indeed enhances the function of sialidase.

15

Example 13: In vitro Activities of Sialidase Fusion Proteins

Stocks of Influenza Viruses

Influenza viral strains are obtained from ATCC and the repository at St. Jude Children's Research Hospital. All experiments involving influenza viruses are conducted at Bio-safety level II.

Viruses are propagated on Madin-Darby canine kidney (MDCK) cells in minimal essential medium (MEM) supplemented with 0.3% bovine serum albumin and 0.5 micrograms of trypsin per ml. After incubating for 48 to 72 hours, the culture medium is clarified by low speed centrifugation. Viral particles are pelleted by ultracentrifugation through a 25% sucrose cushion. Purified viruses are suspended in 50% glycerol-0.1M Tris buffer (pH 7.3) and stored at -20°C.

25

Cell protection assay

To evaluate the ability of the Construct #2 AR-AvCD protein to protect cells against influenza viruses, we first treated MDCK cells with AR-AvCD made from Construct #2 or a broad-spectrum bacterial sialidase isolated from *A. ureafaciens*, and challenged the cells with a broad selection of human influenza viruses (IFV), including

30

human IFV A of H1, H2 and H3 subtypes, human IFV B as well as an avian IFV strain. As shown in Figure 9, the fusion protein made from Construct #2 demonstrated 80 to 100% of cell protection that was comparable to the effect of *A. ureafaciens* sialidase.

To perform the assay, MDCK cells were treated with 10 mU of AR-AvCD protein (made using Construct #2) or the isolated sialidase of *A. ureafaciens* at 37°C for 2 hrs. The cells were subsequently challenged with influenza viruses at MOI 0.1 for 1 hr. The cells were washed and incubated in fresh DMDM:F12 supplemented with 0.2% ITS (GIBCO) and 0.6 µg/ml acetylated trypsin (Sigma). The cells were stained with 0.5% crystal violet and 20% methanol for 5 min and rinsed with tap water. The level of viable cells in each well was quantitated by extracting crystal violet by 70% ethanol and reading at 570 nM. Cell protection was calculated by $100 \times \{(\text{sialidase treated sample} - \text{virus only}) / (\text{uninfected sample} - \text{virus only})\}$.

IFV inhibition assay

We evaluated inhibition of IFV amplification by AR-AvCD protein (made using Construct #2) and AR-G₄S-AvCD protein (made using Construct #3) using a cell-based ELISA method (Belshe et al., J Virol., (1988) 62(5), 1508-1512).

To perform the assay, MDCK monolayers in 96 well plates were treated with 16 mU of the sialidases AR-AvCD made from Construct #2 or AR-G₄S-AvCD made from Construct #3 in EDB/BSA buffer (10 mM Sodium Acetate, 150 mM NaCl, 10 mM CaCl₂, 0.5 mM MgCl₂, and 0.5% BSA) for 2 hrs at 37 °C. Both the sialidase treated and the untreated control cells (treated with only EDB/BSA buffer) were infected with 0.1 MOI of virus. After 1 hour, the cells were washed two times with PBS and incubated in DMEM:F12 supplemented with 0.2% ITS (Gibco) and 0.6 µg/ml acetylated trypsin (Sigma). Forty to 48 hours post-infection, the levels of cell-bound virus were determined by using a cell-based ELISA assay. Specifically, cells were fixed in 0.05% glutaraldehyde in PBS and were incubated with 50 µl of 10³ dilution of either anti-influenza A NP antiserum or anti-influenza B (Fitzgerald Inc.) in 0.5% BSA and PBS at 37°C for 1 hr. After washing, each well was incubated with HRP-protein G in 0.5% BSA and PBS for 1 hr. After final washes, 50µl of 25 mM sodium citrate (pH 4.5) containing 0.02% 3,3',5,5'-tetramethylbenzidine dihydrochloride (Sigma) and 0.01% hydrogen

peroxide was allowed to react with the cells at room temperature for 5 min. The reactions were stopped by adding 50 µl of 1M H₂SO₄, and quantitated by measuring optical densities at 450 nM. Percentage viral replication inhibition is calculated by 100% x {(virus only samples – sialidase treated samples)/(virus only samples – uninfected samples)}.

Data on inhibition of viral replication and cell protection EC₅₀'s and selective indexes for recombinant sialidase fusion proteins AR-AvCD made from Construct #2 and AR-G₄S-AvCD made from Construct #3 for a variety of human influenza A and influenza B viruses, as well as equine viruses are shown in Figure 12.

As shown in Figure 10, sialidase fusion proteins strongly inhibited amplification of a broad selection of influenza viruses. Notably, 80-100% viral inhibition (Figure 10) as well as cell protection (Figure 9) was achieved although a maximum of 70-80% of cell surface sialic acid was removed by the sialidase treatment (Figure 8). This finding demonstrates that it is unnecessary to completely eliminate cell surface sialic acid in order to achieve the desired therapeutic effect of treating with the sialidase fusion proteins of the present invention. The residual 20-30% of the surface sialic acid, while being inaccessible for the sialidase fusion proteins, is probably inaccessible for influenza viruses as well.

Cytotoxicity of sialidase fusion proteins

To evaluate the cytotoxicity of AR-AvCD or AR-G₄S-AvCD proteins (made from Constructs #2 and #3), MDCK cells were seeded at low density in 96-well plates and cultured for 5 days in DMEM containing 10% FBS and up to 20 U of AR-AvCD protein or AR-G₄S-AvCD protein per well (both sialidases remained fully active during the entire experiment). Cell density in AR-AvCD or AR-G₄S-AvCD treated or control wells were determined every day by staining the cells with crystal violet and measuring absorption at 570 nM. No inhibition of cell growth was observed even at the highest concentration of AR-AvCD or AR-G₄S-AvCD (100 U/ ml) in the culture. Therefore IC₅₀, which is the drug concentration that inhibits cell growth by 50%, for AR-AvCD or AR-G₄S-AvCD is above 100 U/ml.

Example 14: In vivo Activities of Sialidase Catalytic Domain Fusion Protein

Ferrets can be infected with human unadapted influenza viruses and produce signs of disease comparable to those of humans, which can be treated by antiviral compounds, such as zanamivir (Relenza). (Mendel et al., *Antimicrob Agents Chemother*, (1998) 42(3), 640-646; Smith and Sweet, *Rev. Infect. Dis.*, (1988) 10(1), 56-75; Reuman et al., *J. Virol. Methods*, (1989) 24(1-2), 27-34). To evaluate in vivo efficacy of our compounds, we tested AR-AvCD protein (made from Construct #2) in the ferret model. Specifically, 24 young female ferrets (0.5-0.8 kg) (Marshall Farms, North Rose, NY) that tested negative for the presence of anti-hemagglutinin antibodies in sera were included in the study. Two animals were placed in each cage and allowed to acclimate for 3 days before the experiment. The animals were randomly divided into three groups: 8 animals were treated with drug dilution buffer and viral challenge, 12 animals were treated with AR-AvCD and viral challenge, and 4 animals were treated with AR-AvCD only. A preparation of AR-AvCD dissolved in phosphate buffered saline (PBS) that contains 500 U/ml in sialidase activity and 0.7 mg/ml in protein concentration was used in the study. Animals in the drug treatment groups received 1 ml of AR-AvCD at each dose, which amounts to about 1 mg/kg in dosage level.

Ferrets were anesthetized and inoculated intranasally (0.5 ml into each nostril) with AR-AvCD or PBS twice (8 am and 6pm) and daily for a total of 7 days (2 days prior to the viral challenge and 5 days post virus inoculation). The ferrets were observed following the drug application for signs of intolerance. Viral inoculation was carried out on day 3 between 10-11 am. The viral challenge was done with human A/Bayern/7/95 (H1N1)-like virus at dose 10^5 TCID₅₀ ($\geq 10^4$ ferret ID₅₀). The nasal washes were collected from all animals starting day 2 post AR-AvCD treatment and continued until day 7. To collect nasal washes, 1 ml of sterile PBS was administered intranasally, the sneezed liquid was harvested and its volume was recorded. The nasal washes were centrifuged. The pelleted cells were re-suspended and counted in a hemacytometer under a microscope. The supernatants were collected, aliquoted and stored at -80°C. The protein concentration in cell-free nasal washes was determined by using the Bio-Rad protein reagent according to manufacturer's protocol (Bio-Rad, Hercules, CA). For virus titration of the nasal washes, inoculated MDCK cells were incubated for 3 days at 36°C

in a CO₂ incubator. The monolayers were inspected visually for cytopathic effect (CPE) and aliquots of the cell culture supernatants from each well were tested for the virus presence by a standard hemagglutination assay with guinea pig red blood cells. Viral titer was determined by the Spearman Karber method (1996).

5 In uninfected animals given intranasal AR-AvCD (n=4), no apparent effect on the inflammatory cell counts and protein concentrations in the nasal washes was observed (Figure 15 A and B). Nasal washes from these animals were followed for 7 days and were all negative for viral shedding. No signs of drug-related toxicity were detected in these animals at the drug dose used in this study. In the vehicle-treated group, virus
10 replicated in the nasal epithelium of all 8 ferrets. Viral shedding reached peak values of 4.4 to 5.9 log₁₀TCID₅₀ (mean peak titer of 4.9) on day 1 or 2 post challenge, diminished over time and became negative by day 5 (Figure 13). By contrast, only 3 of 12 AR-AvCD-treated ferrets were positive for viral shedding on day 1 post challenge (Figure 13), and their nasal viral titers were about 100 times lower than those in the vehicle-
15 treated animals (mean 2.4±0.3 vs. 4.4±0.4 log₁₀TCID₅₀) (Figure 13). After day 1, the response to the AR-AvCD treatment varied substantially. Three animals were completely protected against infection, signs of illness, and inflammatory response (Figure 13), ferret tag # 803, 805, 806). The protection was also confirmed by a lack of seroconversion on day-14 post challenge. One ferret (tag #780) did not shed virus during
20 the first three days post challenge, but it died on day 4 post infection from an unrelated injury. The shedding in the remaining 8 ferrets varied during the course of infection, ranging from ferret #812 that shed virus for a day only, to the ferret #791 that shed virus for 5 days.

Infection in the ferrets that shed virus for at least one day was confirmed by more
25 than a 16-fold rise in the post-challenge anti-HA antibody titer (seroconversion). There was no apparent effect of AR-AvCD treatment on the anti-HA titers in post-challenge sera (320-1280, vs. 160-1280, vehicle- and drug-treated group, respectively).

In ferrets that shed the virus despite the AR-AvCD treatment (n=8), the inflammatory response was reduced and animals appeared to be more alert and active
30 compared to the untreated ferrets that were invariably lethargic and feverish. For this group of 8 infected, AR-AvCD-treated animals, the mean AUC (area under the curve)

value calculated for the nasal protein concentrations was reduced by approximately 40% (2.68 vs. 4.48, arbitrary units) compared to the vehicle-treated infected animals (Figure 11B). In vehicle-treated infected animals, the number of inflammatory cells in nasal washes was increased to approximately 100-fold above those in uninfected animals on day 2 post challenge. These levels were sustained for 4 additional days. The AR-AvCD-treated animals exhibited a significant reduction in the number of inflammatory cells in the nasal washes. Specifically, the AUC value for cell counts was reduced by approximately 3-fold in the AR-AvCD-treated animals compared to the vehicle-treated infected animals (1965 vs. 674, arbitrary units, Figure 11A). The observed reduction in the inflammatory response indicates the importance of inhibiting viral replication at the early stage of infection.

Example 15. Inhibition of Bacterial Cell Adhesion by Sialidase Fusion Proteins

Bacteria

S. pneumoniae: 10 encapsulated strains of different serotypes are selected from the clinical isolates deposited at ATCC. Bacteria are maintained as frozen stocks and passaged on tryptic soy agar plates containing 3% sheep blood (Difco & Micropure Medical Inc.) for 18 hr at 37°C in 5% CO₂. To label pneumococci with radioisotope, an inoculum is taken from a 1- to 2-day plate culture, added to lysine-deficient tryptic soy broth containing 70 µCi of [³H] lysine per ml and incubated at 37°C in 5% CO₂. The growth of each culture is monitored by light absorbance at 595 nm. At late logarithmic phase, the bacteria are harvested, washed twice by centrifugation (13,000rpm x 3min), and resuspended in L-15 medium (without phenol red) plus 0.1% BSA (L-15-BSA) (Cundell and Tuomanen, Microb. Pathog., (1994) 17(6), 361-374) (Barthelson et al., Infect. Immun., (1998) 66(4), 1439-1444).

H. influenzae: 5 strains of type b (Hib) and 10 nontypable strains (NTHi) are obtained from the clinical isolates deposited at ATCC. All strains are stocked in brain heart infusion (BHI, Difco) containing hemin (ICN) and NAD (Sigma) and kept frozen until use; then they are cultured on BHI agar supplemented with hemin and NAD and grown

for 14 hr at 37°C with 5% CO₂. (Kawakami et al., Microbiol. Immunol., (1998) 42(10), 697-702). To label the bacteria with [³H], *H. influenzae* cells are inoculated in BHI broth containing hemin, NAD and [³H]leucine at 250 µCi/ml and allowed to grow until late logarithmic phase and then harvested, washed and resuspended in L-15-BSA (Barthelson et al., Infect. Immun., (1998) 66(4), 1439-1444).

Cell Adhesion Assay

All [³H]-labeled bacteria are suspended in L-15-BSA after washing, the bacterial concentration is determined by visual counting with a Petroff-Hausser chamber, radioactivity is determined by scintillation counting, and the specific activity of the [³H]-labeled cells is calculated. Preparations of bacteria with 7cpm/1000 cells or greater are used. The bacteria are diluted to 5 x 10⁸ cells/ml. BEAS-2B cell monolayers are incubated with [³H]-labeled bacterial suspension containing 5 x 10⁷ bacteria at 37°C in 5% CO₂. After 30 min, unbound bacteria are removed by washing with L-15-BSA for 5 times. Bacteria attached to the WD-HAE tissue samples are quantitated by scintillation counting.

Desialylation of BEAS-2B cells by sialidase fusion proteins and effects on cell adhesion by *H. influenzae* and *S. pneumoniae*.

BEAS-2B cells are incubated with 1-50 mU of AR-AvCD for 2 hours. Cell adhesion assay will be carried out using *H. influenzae* and *S. pneumoniae* strains as described above. Mock treated cells are used as positive control. Efficacy of AR-AvCD is quantitated as the EC₅₀, which is the amount of enzyme to achieve 50% inhibition on bacterial adhesion.

Example 16. Improving Transduction Efficiency of AAV Vector using Sialidase Fusion Proteins

In vitro Experiments

An experiment demonstrating effect of AR-AvCD is performed in a way similar to the procedure published (Bals et al., J Virol., (1999) 73(7), 6085-6088). A monolayer

of Well-Differentiated Airway Epithelium (WDAE) cells is maintained in transwells (Karp et al., *Methods Mol. Biol.*, (2002) 188, 115-137; Wang et al., *J Virol.*, (1998) 72(12), 9818-9826). In order to eliminate sialic acid from the cell surface the culture medium is replaced with serum free medium in which 0.5-10 units of AR-AvCD are dissolved. The cells are treated for 30 min to 6 hours. The cell monolayers are washed, transduced with AAV, and transduction efficiency is estimated using standard procedures. Several transwells are treated with medium only (without AR-AvCD) to serve the purpose of control (basal transduction efficiency). Additional controls may include the transwells treated with AR-AvCD only to assess cytotoxic effect of desialylation. A reporter virus is used for facile detection of transduced cells. Examples of reporter AAV and their use have been described in literature and include AAV-CMV-eGFP, AAV2LacZ (Bals et al., *J Virol.*, (1999) 73(7), 6085-6088; Wang et al., *Hum. Gene Ther.*, (2004) 15(4), 405-413) and alkaline phosphatase (Halbert et al., *Nat. Biotechnol.*, (2002) 20(7), 697-701). The efficiency is estimated by light microscopy of the cells that were fixed and treated with appropriate substrate (if lacZ or AP containing virus is used) or fluorescent microscopy of live cells (if GFP is used). According to the experiments conducted at NexBio with NHBE primary epithelial cells (Cambrex, Walkersville, MD) the maximum amount of removal of sialic acid is achieved in less than one hour when 10 units of AR-AvCD per transwell are used. Other cell lines used (e.g. MDCK) become desialylated with much less AR-AvCD administered (0.1 U for 1 hour). It is therefore our estimate that a treatment of WDAE with 10 U of AR-AvCD for 2 hours will be sufficient to remove accessible sialic acid and provide significant enhancement of transduction of WDAE cells with AAV.

25 Testing Effect of AR-AvCD Treatment on AAV Transduction in an Animal Model.

To demonstrate effect of AR-AvCD treatment in animal model an experiment similar to previously described is conducted (Flotte et al., *Proc. Natl. Acad. Sci. U. S. A.*, (1993) 90(22), 10613-10617; Halbert et al., *Nat. Biotechnol.*, (2002) 20(7), 697-701). Several hours (1-6) prior to administration of AAV AR-AvCD is delivered to mice lungs by nasal aspiration of aerosole or lyophilized AR-AvCD powder according to previously published protocol (Flotte et al., *Proc. Natl. Acad. Sci. U. S. A.*, (1993) 90(22), 10613-

10617). AAV carrying reporter gene (alkaline phosphatase) is delivered by nasal aspiration, mice are euthanized 4 weeks later and transduced cells are detected in fixed lungs as previously described (Halbert et al., J Virol., (1998) 72(12), 9795-9805).

5 ***Example 17. Sialidase Treatment Inhibits Mast Cell Functions and Smooth Muscle Contraction in the Trachea.***

10 Using experimental methods described previously (Cocchiara et al., J Neuroimmunol., (1997) 75(1-2), 9-18), it will be demonstrated that treatment by compounds of the present invention prevents substance P (SP) induced histamine release by mast cells. Using another set of experiments (Stenton et al., J Pharmacol. Exp. Ther., (2002) 302(2), 466-474), treatment by compounds of the present invention will inhibit β -hexosaminidase release by mast cells stimulated by two PAR-activating peptides (PAR stands for proteinase-activated receptors).

15 Compounds of the present invention will be administered intratracheally in guinea pigs and the airway reactivity will be assessed in the animals as described previously (Jarreau et al., Am. Rev. Respir. Dis., (1992) 145(4 Pt 1), 906-910; Stenton et al., J Pharmacol. Exp. Ther., (2002) 302(2), 466-474). Sialidase treatment should not induce nonspecific airway hyperreactivity judged by the reaction to multiple inducers. In
20 addition, sialidase treatment should reduce substance P-induced bronchoconstriction. Similarly, compounds of the present invention will be used to treated the isolated guinea pig and rat trachea and lung (Kai et al., Eur. J. Pharmacol., (1992) 220(2-3), 181-185; Stenton et al., J Pharmacol. Exp. Ther., (2002) 302(2), 466-474). Again recombinant sialidase treatment will have no effect on smooth muscle contractions induced by
25 acetylcholine, histamine and 5-hydroxytryptamine. In addition, it will inhibit tracheal contraction induced by antigen (ovalbumin) or compound 48/80.

30

Bibliography

- Achyuthan, KE and Achyuthan AM. 2001. Comparative enzymology, biochemistry and pathophysiology of human exo- α -sialidases (neuraminidases). *Comparative Biochem & Physiol part B* 129:29-64.
- 5
- Air, GM and Laver, WG. 1995. Red cells bound to influenza virus N9 neuraminidase are not released by the N9 neuraminidase activity. *Virology* 211:278-284.
- 10 Auerswald EA, Horlein D, Reinhardt G, Schroder W and Schnabel E. 1988. Expression, isolation and characterization of recombinant [Arg¹⁵, Glu⁵²] Aprotinin. *Biol Chem Hoppe-Seyler* Vol 369, Suppl., pp27-35.
- Barbey-Morel CL, Oeltmann TN, Edwards KM and Wright PF. 1987. Role of
- 15 respiratory tract proteases in infectivity of influenza A virus. *J Infect Dis* 155:667-672.
- Bessette PH, Aslund F, Beckwith J and Georgiou G. 1999. Efficient folding of proteins with multiple disulfide bonds in the *Escherichia coli* cytoplasm. *Proc Natl Acad Sci USA* 96:13703-13708.
- 20
- Callan RJ, Hartmann FA, West SE and Hinshaw VS. 1997. Cleavage of influenza A virus H1 hemagglutinin by swine respiratory bacterial proteases. *J Virol* 71:7579-7585.
- Connor, RJ, Kawaoka, Y, Webster, RG and Paulson JC. 1994. Receptor specificity in
- 25 human, avian, and equine H2 and H3 influenza virus isolates. *Virology* 205:17-23.
- Copley, RR, Russell, RB and Ponting, CP. 2001. Sialidase-like Asp-boxes: sequence-similar structures within different protein folds. *Prot Sci* 10:285-292.

- Corfield, AP, Veh, RW, Wember, M, Michalski, JC and Schauer, R. 1981. The release of N-acetyl- and N-glycolloyl-neuraminic acid from soluble complex carbohydrates and erythrocytes by bacterial, viral and mammalian sialidases. *Biochem J* 197:293-299.
- 5 Crennell, SJ, Garman, E, Laver, G, Vimr, E and Taylor, G. 1994. Crystal structure of *Vibrio Cholerae* neuraminidase reveals dual lectin-like domains in addition to the catalytic domain. *Structure* 2:535-544.
- Drzeniek, R. Substrate specificity of neuraminidases. 1973. *Histochem J* 5:271-290.
- 10 Endo Y, Carroll KN, Ikizler MR and Wright PF. 1996. Growth of influenza virus in primary, differentiated epithelial cells derived from adenoids. *J Virol* 70:2055-2058.
- Fritz H and Wunderer G. 1983. Biochemistry and applications of aprotinin, the kallikrein inhibitor from bovine organs. *Arzneim-Forsch* 33:479-494.
- 15 Fukudome, K., Yoshie, O. and Konno, T. 1989. Comparison of human, simian, and bovine rotaviruses for requirement of sialic acid in hemagglutination and cell adsorption. *Virology* 172:196-205.
- 20 Garten W, Bosch FX, Linder D, Rott R and Klenk HD. 1981. Proteolytic activation of the influenza virus hemagglutinin: the structure of the cleavage site and the enzymes involved in cleavage. *Virology* 115:361-374.
- 25 Goger, B, Halden, Y, Rek, A, Mosl, R, Pye, D, Gallagher, J and Kungl, AJ. 2002. Different affinities of glycosaminoglycan oligosaccharides for monomeric and dimeric interleukin-8: a model for chemokine regulation at inflammatory sites. *Biochem* 41:1640-1646.

- Gotoh B, Ogasawara T, Toyoda T, Inocencio N, Hamaguchi M and Nagai Y. 1990. An endoprotease homologous to the blood clotting factor X as a determinant of viral tropism in chick embryo. *EMBO J* 9:4189-4195.
- 5 Granoff, A. & Webster, R. G., ed. *Encyclopedia of Virology*, 2nd Edition, Vol 2.
- Gust, ID, Hampson, AW. and Lavanchy, D. 2001. Planning for the next pandemic. *Rev Med Virol* 11:59-70.
- 10 Hayden, FG. 1996. Amantadine and rimantadine-mechanisms. In *Antiviral drug resistance* (ed. D. D. Richman), pp. 59-77. Chichester, UK: John Wiley & Sons Ltd.
- Hosoya M, Matsuyama S, Baba M, Susuki H and Shigeta S. 1992. Effects of protease inhibitors on replication of various myxoviruses. *Antimicrobial Agents and*
- 15 *Chemotherapy* 36:1432-1436.
- Ito, T. 2000. Interspecies transmission and receptor recognition of influenza A virus. *Microbiol Immunol* 44(6):423-430.
- 20 Janakiraman, MN, White, CL, Laver, WG, Air, GM and Luo, M. 1994. Structure of influenza virus neuraminidase B/lee/40 complexed with sialic acid and a dehydro analog at 1.8-A resolution: implications for the catalytic mechanism. *Biochemistry* 33:8172-8179.
- 25 Kido, H, Niwa, Y, Beppu, Y and Towatari, T. 1996. Cellular proteases involved in the pathogenicity of enveloped animal viruses, human immunodeficiency virus, influenza virus A and sendai virus. *Advan Enzyme Regul* 36:325-347.
- Kido H, Chen Y and Murakami M. 1999. Cellular proteinases and viral infection:
- 30 influenza virus, sendai virus and HIV-1, p.205-217. In B. Dunn (ed.), *Proteases of infectious agents*. Academic Press, New York, N.Y.

- Klenk, HD and Rott, R. 1988. The molecular biology of influenza virus pathogenicity. *Adv Vir Res* 34:247-281.
- Klenk, HD and Garten W. 1994. Host cell proteases controlling virus pathogenicity. *Trend Micro* 2:39-43.
- Kreisel, W, Volk, BA, Buchsel, R. and Reutter, W. 1980. Different half-lives of the carbohydrate and protein moieties of a 110,000-dalton glycoproteins isolated from plasma membranes of rat liver. *Proc Natl Acad Sci USA* 77:1828-1831.
- Krunkosky TM, Fischer BM, Martin LD, Jones N, Akley NJ and Adler KB. 2000. Effects of TNF- β on expression of ICAM-1 in human airway epithelial cells in vitro. *Am J Respir Cell Mol Biol* 22:685-692.
- Lazarowitz SG, Goldberg AR and Choppin PW. 1973. Proteolytic cleavage by plasmin of the HA polypeptide of influenza virus: host cell activation of serum plasminogen. *Virology* 56:172-180.
- Lee, MK and Lander, AD. 1991. Analysis of affinity and structural selectivity in the binding of proteins to glycosaminoglycans: development of a sensitive electrophoretic approach. *Pro Natl Acad Sci USA* 88:2768-2772.
- Meltzer, MI, Cox, NJ and Fukuda, K. 1999. The economic impact of pandemic influenza in the United States: priorities for intervention. *Emerg Infect Dis* 5:659-671.
- Meyer, FA, King, M and Gelman, RA., 1975. On the role of sialic acid in the rheological properties of mucus. *Biochimica et Biophysica Acta* 392: 223-232.
- Milner, CM, Smith, SV, Carrillo MB, Taylor, GL, Hollinshead, M and Campbell, RD. 1997. Identification of a sialidase encoded in the human major histocompatibility complex. *J Bio Chem* 272:4549-4558.

- Monti, E, Preti, A, Venerando, B and Borsani, G. 2002. Recent development in mammalian sialidase molecular biology. *Neurochem Res* 27:646-663.
- Monti, E, Preti, A, Nesti, C, Ballabio, A and Borsani G. 1999. Expression of a novel
5 human sialidase encoded by the NEU2 gene. *Glycobiol* 9:1313-1321.
- Monti, E, Bassi, MT, Papini, N, Riboni, M, Manzoni, M, Veneranodo, B, Croci, G, Preti, A, Ballabio, A, Tettamanti, G and Borsani, G. 2000. Identification and expression of NEU3, a novel human sialidase associated to the plasma membrane. *Biochem J* 349:343-
10 351.
- Murakami M, Towatari T, Ohuchi M, Shiota M, Akao M, Okumura Y, Parry MA and Kido H. 2001. Mini-plasmin found in the epithelial cells of bronchioles triggers infection by broad-spectrum influenza A viruses and Sendai virus. *Eur J Biochem* 268:
15 2847-2855.
- Nakayama, K. 1997. Furin: a mammalian subtilisin/kex2p-like endoprotease involved in process of a wide variety of precursor proteins. *Biochem* 327:625-635.
- 20 Ovcharenko AV and Zhirnov OP. 1994. Aprotinin aerosol treatment of influenza and paramyxovirus bronchopneumonia of mice. *Antiviral Res* 23:107-118.
- Pshezhetsky, A, Richard, C, Michaud, L, Igdoura, S, Wang, S, Elsliger, M, Qu, J, Leclerc, D, Gravel, R, Dallaire, L and Potier, M. 1997. Cloning, expression and
25 chromosomal mapping of human lysosomal sialidase and characterization of mutations in sialidosis. *Nature Genet* 15: 316-320.
- Ramphal, R. and Pyle, M. 1983. Evidence for mucins and sialic acid as receptors for *Pseudomonas aeruginosa* in the lower respiratory tract. *Infect Immun* 41:339-44.
30

Roggentin, P, Kleineidam, RG and Schauer, R. 1995. Diversity in the properties of two sialidase isoenzymes produced by *Clostridium perfringens* spp.. *Biol Chem Hoppe-Seyler* 376:569-575.

- 5 Roggentin, P, Schauer, R, Hoyer, LL and Vimr, ER. 1993. The sialidase superfamily and its spread by horizontal gene transfer. *Mol Microb* 9:915-921.

Rosenberg A. ed. *Biology of the Sialic Acids*. 1995. pp270-273.

- 10 Sakurada, K, Ohta, T and Hasegawa, M. 1992. Cloning, expression and characterization of the *Micromonospora viridifaciens* neuraminidase gene in *Streptomyces lividans*. *J Bacteriol* 174: 6896-6903.

- Schauer, S. ed., pp233. *Sialic Acids Chemistry, Metabolism and Function*. Springer-
15 Verlag, 1982.

Schauer, R. 1982. Chemistry, metabolism, and biological functions of sialic acids. *Adv. Carbohydrate Chem & Biochem* 40:131-235.

- 20 Scheiblaue H, Reinacher M, Tashiro M and Rott R. 1992. Interactions between bacteria and influenza A virus in the development of influenza pneumonia. *J Infec Dis* 166:783-791.

- Sinn PL, Williams G, Vongpunsawad S, Cattaneo R and McCray PB. 2002. Measles
25 virus preferentially transduces the basolateral surface of well-differentiated human airway epithelia. *J Virol* 76:2403-2409.

Skehel, JJ and Wiley, DC. 2000. Receptor binding and membrane fusion in virus entry: the influenza hemagglutinin. *Annu Rev Biochem* 69:531-569.

30

- Tashiro M, Klenk HD and Rott R. 1987. Inhibitory effect of a protease inhibitor, leupeptin, on the development of influenza pneumonia, mediated by concomitant bacteria. *J Gen Virol* 68:2039-2043.
- 5 Tashiro M, Ciborowski P, Reinacher M, Pulverer G, Klenk HD and Rott R. 1987. Synergistic role of staphylococcal proteases in the induction of influenza virus pathogenicity. *Virology* 157:421-430.
- 10 Teufel, M, Roggentin, P. and Schauer, R. 1989. Properties of sialidase isolated from *Actinomyces viscosus* DSM43798. *Biol Chem Hoppe Seyler* 370:435-443.
- Tobita, K, Sugiura, A, Enomoto, C and Furuyama, M. 1975. Plaque assay and primary isolation of influenza A viruses in an established line of canine kidney cells (MDCK) in the presence of trypsin. *Med Microbiol Immunol* 162:9-14.
- 15 Venturi M, Seifert C and Hunte C. 2001. High level production of functional antibody Fab fragments in an oxidizing bacterial cytoplasm. *J Mol Biol* 315:1-8.
- 20 Vimr, DR. 1994. Microbial sialidases: does bigger always mean better? *Trends Microbiol* 2: 271-277.
- Vlasak, R., Luytjes, W., Spaan, W. and Palese, P. 1988. Human and bovine coronaviruses recognize sialic acid-containing receptors similar to those of influenza C viruses. *Proc Natl Acad Sci USA* 85:4526-4529.
- 25 Wada, T, Yoshikawa, Y, Tokuyama, S, Kuwabara, M, Akita, H and Miyagi, T. 1999. Cloning, expression, and chromosomal mapping of a human ganglioside sialidase. *Biochem Biophys Res Commun* 261:21-27.

- Wang, FZ, Akula, SM, Pramod, NP, Zeng, L and Chandran, B. 2001. Human herpesvirus 8 envelope glycoproteins K8.1A interaction with the target cells involves heparan sulfate. *J Virol* 75:7517-27
- 5 Wassilewa, L. 1977. Cell receptor for paramyxoviruses. *Arch Virol* 54:299-305.
- Weisgraber, KH, Rall, SC, Mahley, RW, Milne, RW and Marcel, Y. 1986. Human apolipoprotein E, determination
- 10 Witt, DP and Lander AD. 1994. Differential binding of chemokines to glycosaminoglycan subpopulations. *Curr Bio* 4:394-400.
- Wood, J. 2001. Developing vaccines against pandemic influenza. *Phil Trans R Soc Lond B* 356:1953-1960.
- 15 Xiang Y and Moss B. 2003. Molluscum contagiosum virus interleukin-18 (IL-18) binding protein is secreted as a full-length form that bind cell surface glycosaminoglycans through the C-terminal tail and a furin-cleaved form with only the IL-18 binding domain. *J Virol* 77:2623-2630.
- 20 Zambon, M. 2001. The pathogenesis of influenza in humans. *Rev Med Virol* 11:227-241.
- Zhang L, Peebles ME, Boucher RC, Collins PL and Pickles RJ. 2002. Respiratory syncytial virus infection of human airway epithelial cells is polarized, specific to ciliated cells, and without obvious cytopathology. *J Virol* 76:5654-5666.
- 25 Zhirnov OP, Ovchartenko AV and Bukrinskaya AG. 1982. Protective effect of protease inhibitors in influenza virus infected animals. *Arch Virol* 73:263-272
- 30

- Zhirnov OP, Ovcharenko AV and Bukrinskaya AG. 1982. A modified plaque assay method for accurate analysis of infectivity of influenza viruses with uncleaved hemagglutinin. *Arch Virol* 71:177-183.
- 5 Zhirnov OP. 1983. Proteolytic activation of myxoviruses and a new strategy in the treatment of viral diseases. *Problems Virol.* 4:9-12. (In Russian).
- Zhirnov OP, Ovcharenko AV and Bukrinskaya AG. 1984. Suppression of influenza virus replication in infected mice by protease inhibitors. *J Gen Virol* 65:191-196.
- 10 Zhirnov OP, Ovcharenko AV and Bukrinskaya AG. 1985. Myxovirus replication in chicken embryos can be suppressed by aprotinin due to the blockage of viral glycoprotein cleavage. *J Gen Virol* 66:1633-1638.
- 15 Zhirnov OP. 1987. High protection of animals lethally infected with influenza virus by aprotinin-rimantadine combination. *J Med Virol* 21:161-167.
- Zhirnov OP, Ikizler MR and Wright PF. 2002. Cleavage of influenza A virus hemagglutinin in human respiratory epithelium is cell associated and sensitive to
20 exogenous antiproteases. *J Virol* 76:8682-8689.
- Bartlett J.G., Breiman R.F., Mandell L.A., & File T.M., Jr. (1998) Community-acquired pneumonia in adults: guidelines for management. The Infectious Diseases Society of America. *Clin.Infect.Dis.* 26, 811-838.
- 25 Andrews J., Nadjm B., Gant V., & Shetty N. (2003) Community-acquired pneumonia. *Curr.Opin.Pulm.Med.* 9, 175-180.
- File T.M. (2000) The epidemiology of respiratory tract infections. *Semin.Respir.Infect.* 15, 184-194.

- Macfarlane J. (1994) An overview of community acquired pneumonia with lessons learned from the British Thoracic Society Study. *Semin.Respir.Infect.* **9**, 153-165.
- Matsushima T., Miyashita N., & File T.M., Jr. (2002) Etiology and management of community-acquired pneumonia in Asia. *Curr.Opin.Infect.Dis.* **15**, 157-162.
- 5 Ball P. (1995) Epidemiology and treatment of chronic bronchitis and its exacerbations. *Chest* **108**, 43S-52S.
- Faden H. (2001) The microbiologic and immunologic basis for recurrent otitis media in children. *Eur.J Pediatr.* **160**, 407-413.
- Garcia-Rodriguez, JA and Martinez, MJF. Dynamics of nasopharyngeal colonization by
10 potential respiratory pathogens. *J Antimicrob Chemother* 50[Suppl S2], 59-73. 2002.
- Soriano F. & Rodriguez-Cerrato V. (2002) Pharmacodynamic and kinetic basis for the selection of pneumococcal resistance in the upper respiratory tract. *J Antimicrob Chemother* **50 Suppl S2**, 51-58.
- 15 Mbaki N., Rikitomi N., Nagatake T., & Matsumoto K. (1987) Correlation between *Branhamella catarrhalis* adherence to oropharyngeal cells and seasonal incidence of lower respiratory tract infections. *Tohoku J Exp.Med.* **153**, 111-121.
- Zopf D. & Roth S. (1996) Oligosaccharide anti-infective agents. *Lancet* **347**, 1017-1021.
- Cundell D.R., Weiser J.N., Shen J., Young A., & Tuomanen E.I. (1995) Relationship
20 between colonial morphology and adherence of *Streptococcus pneumoniae*. *Infect.Immun.* **63**, 757-761.
- Karlsson K.A. (1998) Meaning and therapeutic potential of microbial recognition of host glycoconjugates. *Mol.Microbiol.* **29**, 1-11.

- Andersson B., Porras O., Hanson L.A., Lagergard T., & Svanborg-Eden C. (1986) Inhibition of attachment of *Streptococcus pneumoniae* and *Haemophilus influenzae* by human milk and receptor oligosaccharides. *J Infect.Dis.* **153**, 232-237.
- Bals R., Xiao W., Sang N., Weiner D.J., Meegalla R.L., & Wilson J.M. (1999)
- 5 Transduction of well-differentiated airway epithelium by recombinant adeno-associated virus is limited by vector entry. *J Virol.* **73**, 6085-6088.
- Barthelson R., Mobasser A., Zopf D., & Simon P. (1998) Adherence of *Streptococcus pneumoniae* to respiratory epithelial cells is inhibited by sialylated oligosaccharides. *Infect.Immun.* **66**, 1439-1444.
- 10 Cundell D.R. & Tuomanen E.I. (1994) Receptor specificity of adherence of *Streptococcus pneumoniae* to human type-II pneumocytes and vascular endothelial cells in vitro. *Microb.Pathog.* **17**, 361-374.
- Fakih M.G., Murphy T.F., Pattoli M.A., & Berenson C.S. (1997) Specific binding of *Haemophilus influenzae* to minor gangliosides of human respiratory epithelial cells.
- 15 *Infect.Immun.* **65**, 1695-1700.
- Kawakami K., Ahmed K., Utsunomiya Y., Rikitomi N., Hori A., Oishi K., & Nagatake T. (1998) Attachment of nontypable *Haemophilus influenzae* to human pharyngeal epithelial cells mediated by a ganglioside receptor. *Microbiol.Immunol.* **42**, 697-702.
- Solzbacher D., Hanisch F.G., van Alphen L., Gilsdorf J.R., & Schroten H. (2003) Mucin
- 20 in middle ear effusions inhibits attachment of *Haemophilus influenzae* to mucosal epithelial cells. *Eur.Arch.Otorhinolaryngol.* **260**, 141-147.
- van Alphen L., Geelen-van den Broek L., Blaas L., van Ham M., & Dankert J. (1991) Blocking of fimbria-mediated adherence of *Haemophilus influenzae* by sialyl gangliosides. *Infect.Immun.* **59**, 4473-4477.
- 25 Ahmed K., Matsumoto K., Rikitomi N., & Nagatake T. (1996) Attachment of *Moraxella catarrhalis* to pharyngeal epithelial cells is mediated by a glycosphingolipid receptor. *FEMS Microbiol.Lett.* **135**, 305-309.

Hazlett L.D., Moon M., & Berk R.S. (1986) In vivo identification of sialic acid as the ocular receptor for *Pseudomonas aeruginosa*. *Infect.Immun.* **51**, 687-689.

Baker N., Hansson G.C., Leffler H., Riise G., & Svanborg-Eden C. (1990) Glycosphingolipid receptors for *Pseudomonas aeruginosa*. *Infect.Immun.* **58**, 2361-2366.

- 5 Schultze B., Gross H.J., Brossmer R., & Herrler G. (1991) The S protein of bovine coronavirus is a hemagglutinin recognizing 9-O-acetylated sialic acid as a receptor determinant. *J Virol.* **65**, 6232-6237.

- Wuppermann F.N., Hegemann J.H., & Jantos C.A. (2001) Heparan sulfate-like glycosaminoglycan is a cellular receptor for *Chlamydia pneumoniae*. *J Infect.Dis.* **184**,
10 181-187.

Beswick E.J., Travelstead A., & Cooper M.D. (2003) Comparative studies of glycosaminoglycan involvement in *Chlamydia pneumoniae* and *C. trachomatis* invasion of host cells. *J Infect.Dis.* **187**, 1291-1300.

- Martinez I. & Melero J.A. (2000) Binding of human respiratory syncytial virus to cells: implication of sulfated cell surface proteoglycans. *J Gen.Virol.* **81**, 2715-2722.
15

Thomas R.J. & Brooks T.J. (2004) Oligosaccharide receptor mimics inhibit *Legionella pneumophila* attachment to human respiratory epithelial cells. *Microb.Pathog.* **36**, 83-92.

Hirno S., Kelm S., Schauer R., Nilsson B., & Wadstrom T. (1996) Adhesion of *Helicobacter pylori* strains to alpha-2,3-linked sialic acids. *Glycoconj.J* **13**, 1005-1011.

- 20 Simon P.M., Goode P.L., Mobasser A., & Zopf D. (1997) Inhibition of *Helicobacter pylori* binding to gastrointestinal epithelial cells by sialic acid-containing oligosaccharides. *Infect.Immun.* **65**, 750-757.

- Miller-Podraza H., Bergstrom J., Milh M.A., & Karlsson K.A. (1997) Recognition of glycoconjugates by *Helicobacter pylori*. Comparison of two sialic acid-dependent specificities based on haemagglutination and binding to human erythrocyte
25 glycoconjugates. *Glycoconj.J* **14**, 467-471.

- Crocker P.R. & Varki A. (2001) Siglecs, sialic acids and innate immunity. *Trends Immunol.* **22**, 337-342.
- Angata T. & Brinkman-Van der Linden E. (2002) I-type lectins. *Biochim.Biophys.Acta* **1572**, 294-316.
- 5 Lyczak J.B., Cannon C.L., & Pier G.B. (2002) Lung infections associated with cystic fibrosis. *Clin.Microbiol.Rev.* **15**, 194-222.
- Flotte T.R. & Carter B.J. (1998) Adeno-associated virus vectors for gene therapy of cystic fibrosis. *Methods Enzymol.* **292**, 717-732.
- Wagner J.A., Reynolds T., Moran M.L., Moss R.B., Wine J.J., Flotte T.R., & Gardner P.
10 (1998) Efficient and persistent gene transfer of AAV-CFTR in maxillary sinus. *Lancet* **351**, 1702-1703.
- Martinez I. & Melero J.A. (2000) Binding of human respiratory syncytial virus to cells: implication of sulfated cell surface proteoglycans. *J Gen. Virol.* **81**, 2715-2722.
- Park P.W., Pier G.B., Hinkes M.T., & Bernfield M. (2001) Exploitation of syndecan-1
15 shedding by *Pseudomonas aeruginosa* enhances virulence. *Nature* **411**, 98-102.
- Monti E., Preti A., Venerando B., & Borsani G. (2002) Recent development in mammalian sialidase molecular biology. *Neurochem.Res.* **27**, 649-663.
- (1995) *Biology of the Sialic Acids*, 270-273.
- Roggentin P., Kleineidam R.G., & Schauer R. (1995) Diversity in the properties of two
20 sialidase isoenzymes produced by *Clostridium perfringens* spp. *Biol.Chem.Hoppe Seyler* **376**, 569-575.
- Sutter V.L. (1984) Anaerobes as normal oral flora. *Rev.Infect.Dis.* **6 Suppl 1**, S62-S66.
- Gaskell A., Crennell S., & Taylor G. (1995) The three domains of a bacterial sialidase: a beta-propeller, an immunoglobulin module and a galactose-binding jelly-roll. *Structure.*
25 **3**, 1197-1205.

Alvarez P., Buscaglia C.A., & Campetella O. (2004) Improving protein pharmacokinetics by genetic fusion to simple amino acid sequences. *J.Biol.Chem.* **279**, 3375-3381.

Potier M., Mameli L., Belisle M., Dallaire L., & Melancon S.B. (1979) Fluorometric assay of neuraminidase with a sodium (4-methylumbelliferyl-alpha-D-N-acetylneuraminate) substrate. *Anal.Biochem.* **94**, 287-296.

Hirel P.H., Schmitter M.J., Dessen P., Fayat G., & Blanquet S. (1989) Extent of N-terminal methionine excision from Escherichia coli proteins is governed by the side-chain length of the penultimate amino acid. *Proc.Natl.Acad.Sci.U.S.A* **86**, 8247-8251.

Varshavsky A. (1996) The N-end rule: functions, mysteries, uses. *Proc.Natl.Acad.Sci.U.S.A* **93**, 12142-12149.

Belshe R.B., Smith M.H., Hall C.B., Betts R., & Hay A.J. (1988) Genetic basis of resistance to rimantadine emerging during treatment of influenza virus infection. *J Virol.* **62**, 1508-1512.

Mendel D.B., Tai C.Y., Escarpe P.A., Li W., Sidwell R.W., Huffman J.H., Sweet C., Jakeman K.J., Merson J., Lacy S.A., Lew W., Williams M.A., Zhang L., Chen M.S., Bischofberger N., & Kim C.U. (1998) Oral administration of a prodrug of the influenza virus neuraminidase inhibitor GS 4071 protects mice and ferrets against influenza infection. *Antimicrob Agents Chemother* **42**, 640-646.

Smith H. & Sweet C. (1988) Lessons for human influenza from pathogenicity studies with ferrets. *Rev.Infect.Dis.* **10**, 56-75.

Reuman P.D., Keely S., & Schiff G.M. (1989) Assessment of signs of influenza illness in the ferret model. *J.Virol.Methods* **24**, 27-34.

Virology Methods Manual. 1996. London, San Diego, New York, boston, Sydney, Todyo, Toronto: Academic Press, Harcourt Brace & Company.

- Karp P.H., Moninger T.O., Weber S.P., Nesselhauf T.S., Launsbach J.L., Zabner J., & Welsh M.J. (2002) An in vitro model of differentiated human airway epithelia. Methods for establishing primary cultures. *Methods Mol.Biol.* **188**, 115-137.
- Wang G., Davidson B.L., Melchert P., Slepushkin V.A., van Es H.H., Bodner M., Jolly
5 D.J., & McCray P.B., Jr. (1998) Influence of cell polarity on retrovirus-mediated gene transfer to differentiated human airway epithelia. *J Virol.* **72**, 9818-9826.
- Wang A.Y., Peng P.D., Ehrhardt A., Storm T.A., & Kay M.A. (2004) Comparison of adenoviral and adeno-associated viral vectors for pancreatic gene delivery in vivo. *Hum.Gene Ther.* **15**, 405-413.
- 10 Halbert C.L., Allen J.M., & Miller A.D. (2002) Efficient mouse airway transduction following recombination between AAV vectors carrying parts of a larger gene. *Nat.Biotechnol.* **20**, 697-701.
- Flotte T.R., Afione S.A., Conrad C., McGrath S.A., Solow R., Oka H., Zeitlin P.L., Guggino W.B., & Carter B.J. (1993) Stable in vivo expression of the cystic fibrosis
15 transmembrane conductance regulator with an adeno-associated virus vector. *Proc.Natl.Acad.Sci.U.S.A* **90**, 10613-10617.
- Halbert C.L., Standaert T.A., Wilson C.B., & Miller A.D. (1998) Successful readministration of adeno-associated virus vectors to the mouse lung requires transient immunosuppression during the initial exposure. *J Virol.* **72**, 9795-9805.
- 20 Cocchiara R., Bongiovanni A., Albeggiani G., Azzolina A., Lampiasi N., Di Blasi F., & Geraci D. (1997) Inhibitory effect of neuraminidase on SP-induced histamine release and TNF-alpha mRNA in rat mast cells: evidence of a receptor-independent mechanism. *J Neuroimmunol.* **75**, 9-18.
- Stenton G.R., Nohara O., Dery R.E., Vliagoftis H., Gilchrist M., Johri A., Wallace J.L.,
25 Hollenberg M.D., Moqbel R., & Befus A.D. (2002) Proteinase-activated receptor (PAR)-1 and -2 agonists induce mediator release from mast cells by pathways distinct from PAR-1 and PAR-2. *J Pharmacol.Exp.Ther.* **302**, 466-474.

Jarreau P.H., Harf A., Levame M., Lambre C.R., Lorino H., & Macquin-Mavier I. (1992) Effects of neuraminidase on airway reactivity in the guinea pig. *Am.Rev.Respir.Dis.* **145**, 906-910.

Kai H., Makise K., Matsumoto S., Ishii T., Takahama K., Isohama Y., & Miyata T.
5 (1992) The influence of neuraminidase treatment on tracheal smooth muscle contraction.
Eur.J.Pharmacol. **220**, 181-185.

10 All publications, including patent documents, Genbank sequence database entries
including nucleotide and amino acid sequences and accompanying information, and
scientific articles, referred to in this application and the bibliography and attachments are
incorporated by reference in their entirety for all purposes to the same extent as if each
individual publication were individually incorporated by reference.

15 All headings are for the convenience of the reader and should not be used to limit
the meaning of the text that follows the heading, unless so specified.

CLAIMS

We Claim:

5

1. A sialidase catalytic domain protein, wherein said sialidase catalytic domain protein comprises an amino acid sequence that begins at any of the amino acids from amino acid 270 to amino acid 290 of said *Actinomyces viscosus* sialidase protein sequence (SEQ ID NO:12) and ends at any of the amino acids from
10 amino acid 665 to amino acid 901 of said *Actinomyces viscosus* sialidase protein sequence (SEQ ID NO:12); wherein said sialidase catalytic domain protein lacks the *Actinomyces viscosus* sialidase protein sequence that consists of the sequence extending from amino acid 1 to amino acid 269, further wherein said sialidase catalytic domain protein has sialidase activity.

15

2. A nucleic acid molecule comprising a nucleotide sequence encoding the sialidase catalytic domain protein of claim 1.

3. The nucleic acid molecule of claim 2 in an expression vector.

20

4. The sialidase catalytic domain protein of claim 1 comprising SEQ ID NO:14.

5. A nucleic acid molecule comprising a nucleotide sequence encoding the sialidase catalytic domain protein of claim 4.

25

6. The nucleic acid molecule of claim 5 in an expression vector.

7. The sialidase catalytic domain protein of claim 1, wherein said sialidase catalytic domain protein comprises an amino acid sequence that begins at any of the amino acids from amino acid amino acids 270 to 290 of said *Actinomyces viscosus* sialidase protein sequence (SEQ ID NO:12) and ends at any of amino acid residues 665 to 681 of said *Actinomyces viscosus* sialidase protein sequence (SEQ ID NO:12).
8. A nucleic acid molecule comprising a nucleotide sequence encoding the sialidase catalytic domain protein of claim 7.
9. The nucleic acid molecule of claim 8 in an expression vector.
10. The sialidase catalytic domain protein of claim 7, wherein said sialidase catalytic domain protein comprises SEQ ID NO:16.
11. A nucleic acid molecule comprising a nucleotide sequence encoding the sialidase catalytic domain protein of claim 10.
12. The nucleic acid molecule of claim 11 in an expression vector.
13. The sialidase catalytic domain protein of claim 7, wherein said sialidase catalytic domain protein comprises an amino acid sequence that begins amino acid 274 of said *Actinomyces viscosus* sialidase protein sequence (SEQ ID NO:12) and ends at amino acid residues 681 of said *Actinomyces viscosus* sialidase protein sequence (SEQ ID NO:12).
14. A nucleic acid molecule comprising a nucleotide sequence encoding the sialidase catalytic domain protein of claim 13.
15. The nucleic acid molecule of claim 14 in an expression vector.

16. The sialidase catalytic domain protein of claim 7, wherein said sialidase catalytic domain protein comprises an amino acid sequence that begins amino acid 290 of said *Actinomyces viscosus* sialidase protein sequence (SEQ ID NO:12) and ends at amino acid residues 666 of said *Actinomyces viscosus* sialidase protein sequence (SEQ ID NO:12).
17. A nucleic acid molecule comprising a nucleotide sequence encoding the sialidase catalytic domain protein of claim 16.
18. The nucleic acid molecule of claim 17 in an expression vector.
19. The sialidase catalytic domain protein of claim 7, wherein said sialidase catalytic domain protein comprises an amino acid sequence that begins amino acid 290 of said *Actinomyces viscosus* sialidase protein sequence (SEQ ID NO:12) and ends at amino acid residues 681 of said *Actinomyces viscosus* sialidase protein sequence (SEQ ID NO:12).
20. A nucleic acid molecule comprising a nucleotide sequence encoding the sialidase catalytic domain protein of claim 19.
21. The nucleic acid molecule of claim 20 in an expression vector.
22. A fusion protein comprising:
at least one catalytic domain of a sialidase; and
a purification domain, a protein tag, a protein stability domain, a solubility domain, a protein size-increasing domain, a protein folding domain, a protein localization domain, an anchoring domain, an N-terminal domain, a C-terminal domain, a catalytic activity domain, a binding domain, or a catalytic activity-enhancing domain

23. The fusion protein of claim 22, further comprising at least one peptide linker.
24. The fusion protein of claim 22, wherein said catalytic domain is substantially homologous to the catalytic domain of the *Clostridium perfringens* sialidase, substantially homologous to the *Actinomyces viscosus* sialidase, substantially homologous to the *Arthrobacter ureafaciens* sialidase, substantially homologous to the *Micromonospora viridifaciens* sialidase, substantially homologous to the human Neu2 sialidase, or substantially homologous to the human Neu4 sialidase.
25. The fusion protein of claim 24, wherein said catalytic domain is substantially homologous to the catalytic domain of the *Actinomyces viscosus* sialidase.
26. The fusion protein of claim 25 wherein said catalytic domain comprises **SEQ ID NO: 16**.
27. The fusion protein of claim 26, comprising at least one protein purification domain.
28. The fusion protein of claim 27, wherein said at least one protein purification domain is a His tag, a calmodulin binding domain, a maltose binding protein domain, a streptavidin domain, a streptavidin binding domain; an intein domain, or a chitin binding domain.
29. The fusion protein of claim 28, wherein said protein purification domain is a His tag.
-
30. The fusion protein of claim 29, wherein said fusion protein comprises **SEQ ID NO:18**.
31. A nucleic acid molecule encoding the fusion protein of claim 30.

32. The nucleic acid molecule of claim 31 in an expression vector.
33. The nucleic acid molecule of claim 31, comprising **SEQ ID NO:17**.
- 5 34. The nucleic acid molecule of claim 33 in an expression vector.
35. The fusion protein of claim 25, comprising at least one anchoring domain.
- 10 36. The fusion protein of claim 35, wherein said anchoring domain is a GAG-binding domain.
37. The fusion protein of claim 36, wherein said anchoring domain is substantially homologous to the GAG-binding domain of human platelet factor 4 (**SEQ ID NO:2**), substantially homologous to the GAG-binding domain of human interleukin 8 (**SEQ ID NO:3**), substantially homologous to the GAG-binding domain of human antithrombin III (**SEQ ID NO:4**), substantially homologous to the GAG-binding domain of human apoprotein E (**SEQ ID NO:5**), substantially homologous to the GAG-binding domain of human angio-associated migratory protein (**SEQ ID NO:6**), or substantially homologous to the GAG-binding domain of human amphiregulin (**SEQ ID NO:7**).
- 15 20 38. The fusion protein of claim 37, wherein said anchoring domain is substantially homologous to the human amphiregulin GAG-binding domain (**SEQ ID NO:7**).
- 25 39. The fusion protein of claim 38, wherein said anchoring domain comprises the human amphiregulin GAG-binding domain (**SEQ ID NO:7**).
40. The fusion protein of claim 39, wherein said catalytic domain of a sialidase comprises **SEQ ID NO:16**.
- 30

41. The fusion protein of claim 40, comprising **SEQ ID NO:20**.
42. A nucleic acid molecule encoding the fusion protein of claim 41.
- 5 43. The nucleic acid molecule of claim 42 in an expression vector.
44. The nucleic acid molecule of claim 42, comprising **SEQ ID NO:19**.
45. The nucleic acid molecule of claim 44 in an expression vector.
- 10 46. The fusion protein of claim 40, comprising **SEQ ID NO:25**.
47. A nucleic acid molecule encoding the fusion protein of claim 46.
- 15 48. The nucleic acid molecule of claim 47 in an expression vector.
49. The nucleic acid molecule of claim 47, comprising **SEQ ID NO:24**.
50. The nucleic acid molecule of claim 49 in an expression vector.
- 20 51. The fusion protein of claim 40, comprising **SEQ ID NO:38**.
52. The fusion protein of claim 40, further comprising a peptide linker connecting
said human amphiregulin GAG-binding domain to said catalytic domain of a
25 sialidase.
53. The fusion protein of claim 52, comprising **SEQ ID NO:35**.
54. A nucleic acid molecule molecule encoding the fusion protein of claim 53.
- 30 55. The nucleic acid molecule of claim 54, comprising **SEQ ID NO:34**.

56. An expression vector comprising the nucleic acid molecule of claim 55.
57. The fusion protein of claim 52, comprising **SEQ ID NO:37**.
58. A nucleic acid molecule encoding the fusion protein of claim 57.
59. The nucleic acid molecule of claim 58, comprising **SEQ ID NO:36**.
60. An expression vector comprising the nucleic acid molecule of claim 59.
61. The fusion protein of claim 52, comprising **SEQ ID NO:39**.
62. The fusion protein of claim 39, comprising **SEQ ID NO:27**.
63. A nucleic acid molecule encoding the fusion protein of claim 62.
-
64. An expression vector comprising the nucleic acid molecule of claim 63.
65. The nucleic acid molecule of claim 63, comprising **SEQ ID NO:26**.
66. An expression vector comprising the nucleic acid molecule of claim 65.
67. The fusion protein of claim 39, comprising **SEQ ID NO:29**.
68. A nucleic acid molecule encoding the fusion protein of claim 67.
69. An expression vector comprising the nucleic acid molecule of claim 68.
70. The nucleic acid molecule of claim 68, comprising **SEQ ID NO:28**.

71. An expression vector comprising the nucleic acid molecule of claim 70.
72. The fusion protein of claim 39, comprising **SEQ ID NO:31**.
- 5 73. A nucleic acid molecule encoding the fusion protein of claim 72.
74. An expression vector comprising the nucleic acid molecule of claim 73.
- 10 75. The nucleic acid molecule of claim 73, comprising **SEQ ID NO:30**.
77. An expression vector comprising the nucleic acid molecule of claim 75.
78. The fusion protein of claim 39, comprising **SEQ ID NO:33**.
- 15 79. A nucleic acid molecule encoding the fusion protein of claim 78.
80. An expression vector comprising the nucleic acid molecule of claim 79.
- 20 81. The nucleic acid molecule of claim 79, comprising **SEQ ID NO:32**.
82. An expression vector comprising the nucleic acid molecule of claim 81.
83. A pharmaceutical formulation comprising a sialidase.

25

30

84. The pharmaceutical formulation of claim 83, wherein said sialidase is substantially homologous to at least a portion of *Clostridium perfringens* sialidase, substantially homologous to at least a portion of the *Actinomyces viscosus* sialidase, substantially homologous to at least a portion of the *Arthrobacter ureafaciens* sialidase, substantially homologous to at least a portion of the *Micromonospora viridifaciens* sialidase, substantially homologous to at least a portion of the human Neu2 sialidase, or substantially homologous to at least a portion of the human Neu4 sialidase.
85. The pharmaceutical formulation of claim 84, wherein said sialidase is substantially homologous to at least a portion of the *Actinomyces viscosus* sialidase.
86. The pharmaceutical formulation of claim 85, wherein said sialidase comprises **SEQ ID NO:12**.
87. A pharmaceutical formulation comprising the composition of claim 1.
88. A pharmaceutical formulation comprising the composition of claim 7
89. A pharmaceutical formulation comprising the composition of claim 22.
90. A pharmaceutical formulation comprising the composition of claim 26.
91. A pharmaceutical formulation comprising the composition of claim 46.
92. A pharmaceutical formulation comprising the composition of claim 62.
93. The pharmaceutical formulation of claim 83, 85, 87, 88, 89, 90, 91, or 92 formulated as a spray.

94. The pharmaceutical formulation of claim 83, 85, 87, 88, 89, 90, 91, or 92, formulated as an inhalant.
- 5 95. The pharmaceutical formulation of claim 83, 85, 87, 88, 89, 90, 91, or 92, formulated as a solution for injection.
96. The pharmaceutical formulation of claim 83, 85, 87, 88, 89, 90, 91, or 92, formulated as a solution for eye drops.
- 10 97. The pharmaceutical formulation of claim 83, 85, 87, 88, 89, 90, 91, or 92 formulated as a cream, salve, gel, or ointment.
98. The pharmaceutical formulation of claim 83, 85, 87, 88, 89, 90, 91, or 92, formulated as a pill, tablet, lozenge, suspension, or solution that can be administered orally.
- 15 99. A method of treating or preventing viral infection by influenza, parainfluenza, or respiratory syncytial virus, comprising:
applying a therapeutically effective amount of the composition of any of claims
20 83, 85, 87, 88, 89, 90, 91, or 92 to epithelial cells of a subject.
100. The method of claim 99, wherein said applying is by use of a nasal spray.
101. The method of claim 99, wherein said applying is by use of an inhaler.
- 25 102. The method of claim 99, wherein said applying is performed from once to four times a day.
103. A method of treating or preventing infection by a bacterial pathogen, comprising:
30 administering a therapeutically effective amount of the composition of claim 83, 85, 87, 88, 89, 90, 91, or 92 to a subject.

104. The method of claim 103, wherein said pathogen is *Streptococcus pneumoniae*,
Mycoplasma pneumoniae, *Haemophilus influenzae*, *Moraxella catarrhalis*,
Pseudomonas aeruginosa, or *Helicobacter pylori*.
- 5
105. The method of claim 103, wherein said administering is by use of a nasal spray.
106. The method of claim 103, wherein said administering is by use of an inhaler.
- 10 107. The method of claim 103, wherein said administering is by topical application.
108. The method of claim 103, wherein said administering is by oral administration.
109. The method of claim 103, wherein said administering is performed from once to
15 four times a day.
110. A method of treating or preventing allergy or inflammation, comprising:
administering a therapeutically effective amount of the composition of claim 83,
85, 87, 88, 89, 90, 91, or 92 to a subject.
- 20
111. The method of claim 110, wherein said inflammation is associated with asthma,
allergic rhinitis, eczema, psoriasis, exposure to plant or animal toxins, or
autoimmune conditions.
- 25 112. The method of claim 110, wherein said administering is by use of a nasal spray.
113. The method of claim 110, wherein said administering is by use of an inhaler.
114. The method of claim 110, wherein said administering is by use of eye drops.
- 30 115. The method of claim 110, wherein said administering is by topical application.

116. The method of claim 110, wherein said administering is by local or intravenous injection.
- 5 117. The method of claim 110, wherein said administering is performed from once to four times a day.
118. A method of enhancing gene delivery by a recombinant viral vector, comprising:
administering an effective amount of the composition of claim 83, 85, 87, 88, 89,
10 90, 91, or 92 to epithelial cells of a subject prior to or concomitant with the
administration of said recombinant viral vector.
119. The method of claim 118, wherein said recombinant viral vector is a retroviral
vector, a Herpes viral vector, an adenoviral vector, or an adeno-associated viral
15 vector.
120. The method of claim 119, wherein said recombinant viral vector is a recombinant
adeno-associated viral vector.
- 20 121. The method of claim 120, wherein said recombinant adeno-associated viral vector
comprises a gene encoding the cystic fibrosis transmembrane conductance
regulator (CFTR).
122. The method of claim 119, wherein said administering is by use of an inhaler.
25
123. The method of claim 119, wherein said administering is performed from once to
four times a day.
124. A pharmaceutical formulation comprising the composition of claim 62.
30
125. The pharmaceutical formulation of claim 124, formulated as a spray.

126. The pharmaceutical formulation of claim 124, formulated as an inhalant.
127. A method of treating or preventing influenza infection, comprising:
5 applying a therapeutically effective amount of the composition of claim 124
 to epithelial cells of a subject.
128. The method of claim 127, wherein said applying is by use of a nasal spray.
- 10 129. The method of claim 127, wherein said applying is by use of an inhaler.
130. The method of claim 127, wherein said applying is performed from once to four
 times a day.

15




FIG. 1

2 / 13

PF4 (SEQ ID NO:2): ⁴⁷NGRRICLDLQAPLYKKIIKKLLES⁷⁰
IL-8 (SEQ ID NO:3): ⁴⁶GRELCLDPKENWVQRVVEKFLKRAENS⁷²
ATIII (SEQ ID NO:4): ¹¹⁸QIHFFFAKLNCRLYRKANKSSKLVSANRLFGDKS¹⁵¹
ApoE (SEQ ID NO:5): ¹³²ELRVRLASHLRKLRKRLLRDADDLQKRLAVYQAG¹⁶⁵
AAMP (SEQ ID NO:6): ¹⁴RRLRRMESESES²⁵
Ampiregulin (SEQ ID NO:7): ²⁵KRKKKGGKNGKNTTNTKKKNP⁴⁵

FIG. 2

3 / 13

NEU2 (SEQ ID NO: 8): 1 MASLPVLQKE SVFQSGAHA- -YRIPALLYL PGQQSLLAFA EQRASKKDEH
 YR+P+LL + P +LLAF EQR S D H
 NEU4 (SEQ ID NO: 9): 1 MGVPRTPSRT VLFERERTGL TYRVPSLLPV PPGPTLLAFV EQRISPDDSH

NEU2: 49 AELIVLRRGD YDAPTHQVQW QAQEVVAQAR LDGHRSMNPC PLYDAQTGTL FLFFIAIPGQ
 A +VLRRG +W A ++ A HRSMNPC P++DA TGT+ FLFFIA+ G
 NEU4: 51 AHRLVLRRGT LAGGSV--RW GALHVLGTAA LAEHRSMNPC PVHDAGTGTV FLFFIAVLGH

NEU2: 110 VTEQQQLQTR ANVTRLCQVT STDHGRTWSS PRDLTDAAIG PAYREWSTFA VGPGHCLQLN
 E Q+ T N RLC V S D G +W S RDLT+ AIG A ++W+TFA VGPGH +QL
 NEU4: 109 TPEAVQIATG RNAARLCCVA SRDAGLSWGS ARDLTEEAIG GAVQDWATFA VGPGHGVQLP

NEU2: 170 DRARSLVPPA YAYRKLHP-- ---IQRPIPS AFCFLSHDHG RTWARGHFVA QD-TLECQVA
 R L+VPA Y YR I R P +F F S DHG RTW G V + ECQ+A
 NEU4: 169 S-GR-LLVPA YTYRVDRLEC FGKICRTSPH SFAFYSDDHG RTWRCGGLVP NLRSGECQLA

NEU2: 224 EVETGEQRVV TL-NARSHLR ARVQAQSTND GLDFQESQLV KKLVEPPPQG CQGSVISFPS
 V+ G+ NARS L +RVQA ST++ G F ++ V L E G CQGS++ FP
 NEU4: 227 AVDGGQAGSF LYCNARSPLG SRVQALSTDE GTSFLPAERV ASLPETAW-G CQGSIVGFPA

NEU2: 283 P-----

NEU4: 286 PAPNRPRDDS WSVGPRSPLO PPLLGPGVHE PPEEAADVPR GGQVPGGPFS RLQPRGDGP

NEU2: 284 ----- ---RSGPGSP QWLLYTHPTH SWQRADLGAY LNPRPPAPEA
 WLLY+HP R +G L+ P P +
 NEU4: 346 RQPGPRPGVSG DVGSWTALP MPFAAPPQSP TWLLYSHFVG RRARLHMGR LSQSPLDPRS

NEU2: 321 WSEPVLAKG SCAYSDLQSM GTGPDGSPLF GCLYEANDY- --EEIVFLMF TLKQAFPAEY
 W+EP ++ + YSDL S+ G P+G +F +CLYE +L++
 NEU4: 406 WTEPWVIYEG PSGYSDLASI GPAPEGGLVF ACLYESGART SYDEISFCTF SLREVLENVP

NEU2: 378 LPQ

NEU4: 466 ASPKPPNLGD KPRGCCWPS

FIG. 3

Substrate Specificity of Bacteria and Fungal Sialidases

Substrates	Sialidase activity*					Actinomyces viscosus
	Vibrio Cholerae perfringens (71Kd)	Clostridium perfringens (43Kd)	Arthrobacter ureafaciens	Salmonella typhimurium		
Oligo- and polysaccharides						
II ³ Neu5AcLac	100	100	100	100	100	100
II ⁶ Neu5AcLac	53	44	157	0.4	0.4	462
Colominic acid (α2-8)	30	33	63	0.1	0.1	300
Glycoproteins						
Fetuin (α2-3>α2-6)	340	272	59	17	17	—
α1-Acid glycoprotein (α2-6>α2-3)	1000	555	—	—	—	761
Submandibular gland mucin (α2-6)	400	139	—	—	—	123
Submaxillary gland mucin (α2-6)	—	—	56	—	—	—
Gangliosides						
Gangliosides mixtures	(360)	(350)	78	34	34	285
Synthetic 4MU-Neu5Ac	1580	605	—	1050	1050	—

* Each value represents a relative sialidase activity when the activity directed toward II³Neu5AcLac is regarded as 100.

FIG. 4

5 / 13

ccattggggcatcaccatcaccatcatctagagggagatcatccacaagctacaccagcacct
 M G H H H H H H L E G D H P Q A T P A P
 gcaccagatgctagcactgagctgccagcaagcatgtctcaggctcagcatcttgagca
 A P D A S T E L P A S M S Q A Q H L A A
 aatacggctactgataattatcgcatccagcgattacaaccgctccgaatgggtgattta
 N T A T D N Y R I P A I T T A P N G D L
 ctgattagctatgatgaacggccgaaggacaatggaaatgggtgggtccgatgcccctaac
 L I S Y D E R P K D N G N G G S D A P N
 ccgaatcatattgttcagcgctcgctccacagatggcggtaaaaacttggagcgcgccaacc
 P N H I V Q R R S T D G G K T W S A P T
 tatattcatcagggtacggagactggcaagaatgggatattccgacccctcttatgtg
 Y I H Q G T E T G K K V G Y S D P S Y V
 gtggatcatcaaaccggtacaatcttcaattttcatgtgaaatcatacgatcagggctgg
 V D H Q T G T I F N F H V K S Y D Q G W
 ggaggttagccgtgggggaacagaccggaaaaaccgcggtattattcaggcagaggtgtct
 G G S R G G T D P E N R G I I Q A E V S
 acgagcacggataatggatggacgtggacacatcgccacatcacgcggatattacgaaa
 T S T D N G W T W T I T A D I T K
 gataaaccgtggaccgcgcttttgcggcgctccggccaaggcattcagatccagctggg
 D K P W T A R F A A S G Q G I Q I Q H G
 ccgcatgccggccgtctgggtgcaacagtataaccattcgtagcgccgggtggagcggtgcag
 P H A G R L V Q Q Y T I R T A G G A V Q
 gctgtatcggtttattccgatgatcatgggaaaacgtggcaggctggcaccoccgattggg
 A V S V Y S D D H G K T W Q A G T P I G
 acgggtatggatgaaaacaaagtgttagagctgtctgacggctctctgatgctgaacagt
 T G M D E N K V V E L S D G S L M L N S
 cgtgcgtcggacgggagcggttttcgtaagggttgcgcatagcactgatgggtgggcagacc
 R A S D G S G F R K V A H S T D G G Q T
 tgggtccgaaccgggttttcggacaaaaatttgcgggattcgggttgataatgccagataatt
 W S E P V S D K N L P D S V D N A Q I I
 cgtgcgttttcctaattgctgcccccgatgacccgcgcgcgaaagtacttctctgagtc
 R A F P N A A P D D P R A K V L L L S H
 tccccaaatccacgtccgtgggtcccggtacgtggtacgataagcatgtcatgtgatgac
 S P N P R P W S R D R G T I S M S C D D
 ggggcctcatggaccacttccaaagtttttcacgaaccggtttgtgggctacacgactatt
 G A S W T T S K V F H E P F V G Y T T I
 gcagttcagagtgatggaagcatcggtctgctgtcggaggacgcgcacaatggcgctgat
 A V Q S D G S I G L L S E D A H N G A D
 tatgggtggcatctggtatcgtaattttacgatgaactggctgggagaacaatgtggacaa
 Y G G I W Y R N F T M N W L G E Q C G Q
 aaaccgcggaataagctt
 K P A E

FIG. 5

6 / 13

ccatgggttaagcgcaaaaaaaaaaggcggaacacggtaaaaatcgtcgtaaccgtaagaaa
M V K R K K K G G K N G K N R R N R K K
aaaaatcctggagatcatccacaagctacaccagcacctgcaccagatgctagcactgag
K N P G D H P Q A T P A P A P D A S T E
ctgccagcaagcatgtctcaggctcagcatcttgacgaaatacggctactgataattat
L P A S M S Q A Q H L A A N T A T D N Y
cgcatccagcgattacaaccgctccgaatgggtgatttactgattagctatgatgaacgg
R I P A I T T A P N G D L L I S Y D E R
ccgaaggacaatggaaatgggtggttcgatgcccctaaccgaatcatattgttcagcgt
P K D N G N G G S D A P N P N H I V Q R
cgctccacagatggcggtaaaacttgagcgcgccaacctatattcatcagggtacggag
R S T D G G K T W S A P T Y I H Q G T E
actggcaagaaagtgggatattccgacccctcttatgtggtggatcatcaaacgggtaca
T G K K V G Y S D P S Y V V D H Q T G T
atcttcaattttcatgtgaaatcatagcagggctggggaggttagccgtgggggaaca
I F N F H V K S Y D Q G W G G S R G G T
gacccggaaaaccgcggttattcaggcagaggtgtctacgagcacggataatggatgg
D P E N R G I I Q A E V S T S T D N G W
acgtggacacatcgccatccacggcggtatattacgaaagataaacgggtggaccgcgct
T W T H R T I T A D I T K D K P W T A R
tttgcggcggtccggccaaggcattcagatccagcatgggcccgcgcatgccggccgtctggtg
F A A S G G I Q I Q H G P H A G R L V
caacagtataccattcgtacggccggtggagcgggtgcaggctgtatcggtttattccgat
Q Q Y T I R T A G G A V Q A V S V Y S D
gatcatgggaaaacgtggcaggctggcaccggatgggacgggtatggatgaaaacaaa
D H G K T W Q A G T P I G T G M D E N K
gttgtagagctgtctgacggctctctgatgtgaacagtcgtgcgtcggacggggagcggc
V V E L S D G S L M L N S R A S D G S G
tttcgtaagggtgcatagcactgatgggtgggcagacctggtccgaaccggtttcggac
F R K V A H S T D G G Q T W S E P V S D
aaaaatttgccggattcgggtgataatgccagataattcgtgcgtttcctaattgctgcc
K N L P D S V D N A Q I I R A F P N A A
cccgatgacccgcgcgcgaaagtacttcttctgagtcattccccaaatccacgtccgtgg
P D D P R A K V L L L S H S P N P R P W
tcccgggatcgtggtacgataagcatgtcatgtgatgacggggcctcatggaccattcc
S R D R G T I S M S C D D G A S W T T S
aaagtttttcacgaaccgtttgtgggctacacgactattgcagttcagagtgatggaagc
K V F H E P F V G Y T T I A V Q S D G S
atcggtctgctgtcggaggacgcgcacaatggcgctgattatgggtggcatctggtatcgt
I G L L S E D A H N G A D Y G G I W Y R
aattttacgatgaactggctgggagaacaatgtggacaaaaaccgcggaataagott
N F T M N W L G E Q C G Q K P A E - A

FIG. 6

7 / 13

ccatgggttaagcgcaaaaaaaaaaggcggaacggtaaaaatcgtcgtaaccgtaagaaa
M V K R K K K G G K N G K N R R N R K K
aaaaatcctgggtgggtgggttctggagatcatccacaagctacaccagcacctgcacca
K N P G G G G S G D H P Q A T P A P A P
gatgctagcactgagctgccagcaagcatgtctcaggctcagcatcttgagcaaatagc
D A S T E L P A S M S Q A Q H L A A N T
gctactgataattatcgcatccagcgattacaaccgctccgaatgggtgatttactgatt
A T D N Y R I P A I T T A P N G D L L I
agctatgatgaacggcggaaggacaatggaaatgggtgggtccgatgcccctaaccggaat
S Y D E R P K D N G N G G S D A P N P N
catattgttcagcgctcgctccacagatggcggtaaaacttggagcgcgccaacctatatt
H I V Q R R S T D G G K T W S A P T Y I
catcagggtacggagactggcaagaaagtgggatattccgacccctcttatgtgggtggat
H Q G G T E T G K K V G Y S D P S Y V V D
catcaaaccggtacaatcttcaattttcatgtgaaatcatacgatcagggctggggaggt
H Q T G T I F N F H V K S Y D Q G W G G
agccgtgggggaacagacccggaaaaccgcggtattattcaggcagaggtgtctacgagc
S R G G T D P E N R G I I Q A E V S T S
acggataatggatggacgtggacacatcgacccatcaccgcggtattacgaaagataaa
T D N G W T W T H R T I T A D I T K D K
ccgtggaccgcgcttttgcggcgctccggccaaggcattcagatccagcatgggcccgc
P W T A R F A A S G Q G I Q I Q H G P H
gccggccgtctgggtgcaacagtataccattcgtacggccggtggagcggtgcaggctgta
A G R L V Q Q Y T I R T A G G A V Q A V
tcggttttattccgatgatcatgggaaaacgtggcaggctggcaccgattgggacgggt
S V Y S D D H G K T W Q A G T P I G T G
atggatgaaaacaaagtgttagagctgtctgacggctctctgatgtgaacagtcgtgcg
M D E N K V V E L S D G S L M L N S R A
tcggacgggagcggctttcgtaagggttcgcgcatagcactgatgggtgggcagacctgggtcc
S D G S G F R K V A H S T D G G Q T W S
gaaccgggtttcggaacaaaatttgcgggattcgggttgataatgccagataattcgtgcg
E P V S D K N L P D S V D N A Q I I R A
tttcctaattgctgcccccgatgacccgcgcgcgaaagtacttcttctgagtcattcccca
F P N A A P D D P R A K V L L L S H S P
aatccacgtccgtgggtcccggtacgtgggtacgataagcatgtcatgtgatgacggggcc
N P R P W S R D R G T I S M S C D D G A
tcatggaccacttccaaagtttttcacgaaccggtttgtgggctacacgactattgcagtt
S W T T S K V F H E P F V G Y T T I A V
cagagtgatggaagcatcggtctgctgtcggaggacgcgcacaatggcgctgattatgggt
Q S D G S I G L L S E D A H N G A D Y G
ggcatctggtatcgtaattttacgatgaactgggtgggagacaatgtggacaaaaacc
G I W Y R N F T M N W L G E Q C G Q K P
gcggaataagctt
A E - A

FIG. 7

8 / 13




FIG. 8

9 / 13




FIG. 9

10 / 13




FIG. 10

11 / 13




FIG. 11

12 / 13

Virus	AR-AvCD				AR-G ₄ S-AvCD			
	Inhibition of Viral Replication EC50	Selective Index	Cell Protection EC50	Selective Index	Inhibition of Viral Replication EC50	Selective Index	Cell Protection EC50	Selective Index
A/PR/8/34 (H1N1)	12.3 ± 7.4	>8163	50.3 ± 13.1	>1990	13.5 ± 1.4	>7407	43.8 ± 22.3	>2286
A/WS/33 (H1N1)	6.5 ± 3.9	>15444	17.7 ± 13.9	>5666	12.2 ± 11.0	>8197	19.1 ± 17.6	>5249
A/NWS/33 (H1N1)	2.5	>20000	10.8 ± 12.4	>9302	2.5	>40000	15.9 ± 19.3	>6249
A/Japan/305/57 (H2N2)	5.1 ± 3.4	>19512	11.3 ± 0.4	>8889	3.8 ± 1.8	>26667	14.9 ± 0.2	>6273
A/Victoria/504/2000 (H3N2)	2.5 ± 0.0	>40000	18.3 ± 1.0	>5457	2.6 ± 0.2	>38095	55.3 ± 0.4	>1810
A/HongKong/8/68 (H3N2)	2.5 ± 0.0	>40000	31.5 ± 38.2	>3175	2.5 ± 0.0	>40000	30.6 ± 30.2	>3265
B/Lee/40	4.1 ± 1.3	>24540	11.4 ± 3.7	>8791	3.5 ± 1.3	>28986	8.4 ± 2.7	>11940
B/Maryland/1/59	4.3 ± 2.5	>23392	2.5 ± 0.0	>40000	5.5 ± 4.2	>18265	5.6 ± 3.0	>17778
Turkey/Wis/66 (H9N2)	3.9 ± 0.5	>25478	16.6 ± 16.2	>6033	7.1 ± 0.9	>14035	20.4 ± 18.6	>4914
Equine/Prague/2/62 (H7N7)	n/a	n/a	15.8 ± 5.3	>6349	n/a	n/a	25.5 ± 5.7	>3922

FIG. 12

13 / 13

No.	Ferret tag no.	Virus titer log ₁₀ TCID ₅₀ /ml on day p.i. ^a					Post-challenge HI titers
		1	2	3	4	5	
Vehicle-treated group							
1	228	5.7	4.2	4.2	1.7	-	640
2	784	3.9	4.9	1.9	1.9	-	640
3	793	4.4	4.2	2.4	3.9	-	640
4	794	4.9	5.9	1.4	-	-	160
5	789	4.4	4.2	3.4	3.4	-	640
6	799	3.7	4.4	3.4	-	-	320
7	811	4.4	4.4	-	-	-	1280
8	841	4.2	4.7	2.7	1.9	-	320
	mean ^b	4.4	4.7	2.7	3.7	-	
	SD	0.4	0.7	1.0	0.4	-	
	Shed/total	8/8	8/8	7/8	5/8	0/8	
Fludase-treated group							
1	780	-	-	-	NA	NA	NA
2	791	2.2	5.2	4.9	4.2	1.7	640
3	804	-	4.7	3.7	1.7	-	1280
4	803	-	-	-	-	-	≤10
5	805	-	-	-	-	-	≤10
6	806	-	-	-	-	-	≤10
7	810	2.2	4.7	3.2	2.9	-	160
8	812	-	-	4.4	-	-	640
9	813	-	3.2	4.4	4.7	-	160
10	819	2.7	5.2	-	-	-	320
11	828	-	4.9	1.9	1.7	-	320
12	843	-	4.4	4.9	4.9	3.4	320
	mean ^b	2.4	4.6	3.9	3.4	2.6	
	SD	0.3	0.7	1.1	1.5	1.2	
	Shed/total	3/12	7/12	7/12	6/11	2/11	

^a - all nasal washes collected after day 5 post challenge were negative for virus presence.^b - mean value was calculated for the ferrets that shed virus.

Nasal washes recovered from the uninfected treated ferrets were negative for virus (not shown).

NA - not applicable, the ferret died on day 4 post infection due to an accident.

FIG. 13

SEQUENCE LISTING

<110> FANG, Fang
MALAKHOV, Michael

<120> A NOVEL CLASS OF THERAPEUTIC PROTEIN BASED MOLECULES

<130> NB-101.P11PC

<150> US 60/428,535
<151> 2002-11-22

<150> US 10/939,262
<151> 2004-09-10

<150> US 10/718,986
<151> 2003-11-21

<150> US 60/580,084
<151> 2004-06-16

<150> US 60/561,749
<151> 2004-04-13

<150> US 60/464,217
<151> 2003-04-19

<160> 39

<170> PatentIn version 3.3

<210> 1
<211> 58
<212> PRT
<213> Bos taurus

<400> 1

Arg Pro Asp Phe Cys Leu Glu Pro Pro Tyr Thr Gly Pro Cys Lys Ala
1 5 10 15

Arg Ile Ile Arg Tyr Phe Tyr Asn Ala Lys Ala Gly Leu Cys Gln Thr
20 25 30

Phe Val Tyr Gly Gly Cys Arg Ala Lys Arg Asn Asn Phe Lys Ser Ala
35 40 45

Glu Asp Cys Met Arg Thr Cys Gly Gly Ala
50 55

<210> 2
<211> 24
<212> PRT

<213> Homo sapiens

<400> 2

Asn Gly Arg Arg Ile Cys Leu Asp Leu Gln Ala Pro Leu Tyr Lys Lys
1 5 10 15

Ile Ile Lys Lys Leu Leu Glu Ser
20

<210> 3

<211> 27

<212> PRT

<213> Homo sapiens

<400> 3

Gly Arg Glu Leu Cys Leu Asp Pro Lys Glu Asn Trp Val Gln Arg Val
1 5 10 15

Val Glu Lys Phe Leu Lys Arg Ala Glu Asn Ser
20 25

<210> 4

<211> 34

<212> PRT

<213> Homo sapiens

<400> 4

Gln Ile His Phe Phe Phe Ala Lys Leu Asn Cys Arg Leu Tyr Arg Lys
1 5 10 15

Ala Asn Lys Ser Ser Lys Leu Val Ser Ala Asn Arg Leu Phe Gly Asp
20 25 30

Lys Ser

<210> 5

<211> 34

<212> PRT

<213> Homo sapiens

<400> 5

Glu Leu Arg Val Arg Leu Ala Ser His Leu Arg Lys Leu Arg Lys Arg
1 5 10 15

Leu Leu Arg Asp Ala Asp Asp Leu Gln Lys Arg Leu Ala Val Tyr Gln
20 25 30

Ala Gly

<210> 6
<211> 12
<212> PRT
<213> Homo sapiens

<400> 6

Arg Arg Leu Arg Arg Met Glu Ser Glu Ser Glu Ser
1 5 10

<210> 7
<211> 21
<212> PRT
<213> Homo sapiens

<400> 7

Lys Arg Lys Lys Lys Gly Gly Lys Asn Gly Lys Asn Thr Thr Asn Thr
1 5 10 15

Lys Lys Lys Asn Pro
20

<210> 8
<211> 379
<212> PRT
<213> Homo sapiens

<400> 8

Met Ala Ser Leu Pro Val Leu Gln Lys Glu Ser Val Phe Gln Ser Gly
1 5 10 15

Ala His Ala Tyr Arg Ile Pro Ala Leu Leu Tyr Leu Pro Gly Gln Gln
20 25 30

Ser Leu Leu Ala Phe Ala Glu Gln Arg Ala Ser Lys Lys Asp Glu His
35 40 45

Ala Glu Leu Ile Val Leu Arg Arg Gly Asp Tyr Asp Ala Pro Thr His
50 55 60

Gln Val Gln Trp Gln Ala Gln Glu Val Val Ala Gln Ala Arg Leu Asp
 65 70 75 80

Gly His Arg Ser Met Asn Pro Cys Pro Leu Tyr Asp Ala Gln Thr Gly
 85 90 95

Thr Leu Phe Leu Phe Phe Ile Ala Ile Pro Gly Gln Val Thr Glu Gln
 100 105 110

Gln Gln Leu Gln Thr Arg Ala Asn Val Thr Arg Leu Cys Gln Val Thr
 115 120 125

Ser Thr Asp His Gly Arg Thr Trp Ser Ser Pro Arg Asp Leu Thr Asp
 130 135 140

Ala Ala Ile Gly Pro Ala Tyr Arg Glu Trp Ser Thr Phe Ala Val Gly
 145 150 155 160

Pro Gly His Cys Leu Gln Leu Asn Asp Arg Ala Arg Ser Leu Val Val
 165 170 175

Pro Ala Tyr Ala Tyr Arg Lys Leu His Pro Ile Gln Arg Pro Ile Pro
 180 185 190

Ser Ala Phe Cys Phe Leu Ser His Asp His Gly Arg Thr Trp Ala Arg
 195 200 205

Gly His Phe Val Ala Gln Asp Thr Leu Glu Cys Gln Val Ala Glu Val
 210 215 220

Glu Thr Gly Glu Gln Arg Val Val Thr Leu Asn Ala Arg Ser His Leu
 225 230 235 240

Arg Ala Arg Val Gln Ala Gln Ser Thr Asn Asp Gly Leu Asp Phe Gln
 245 250 255

Glu Ser Gln Leu Val Lys Lys Leu Val Glu Pro Pro Pro Gln Gly Cys
 260 265 270

Gln Gly Ser Val Ile Ser Phe Pro Ser Pro Arg Ser Gly Pro Gly Ser
 275 280 285

Pro Gln Trp Leu Leu Tyr Thr His Pro Thr His Ser Trp Gln Arg Ala
290 295 300

Asp Leu Gly Ala Tyr Leu Asn Pro Arg Pro Pro Ala Pro Glu Ala Trp
305 310 315 320

Ser Glu Pro Val Leu Leu Ala Lys Gly Ser Cys Ala Tyr Ser Asp Leu
325 330 335

Gln Ser Met Gly Thr Gly Pro Asp Gly Ser Pro Leu Phe Gly Cys Leu
340 345 350

Tyr Glu Ala Asn Asp Tyr Glu Glu Ile Val Phe Leu Met Phe Thr Leu
355 360 365

Lys Gln Ala Phe Pro Ala Glu Tyr Leu Pro Gln
370 375

<210> 9
<211> 424
<212> PRT
<213> Homo sapiens

<400> 9

Leu Ala Gly Gly Ser Val Arg Trp Gly Ala Leu His Val Leu Gly Thr
1 5 10 15

Ala Ala Leu Ala Glu His Arg Ser Met Asn Pro Cys Pro Val His Asp
20 25 30

Ala Gly Thr Gly Thr Val Phe Leu Phe Phe Ile Ala Val Leu Gly His
35 40 45

Thr Pro Glu Ala Val Gln Ile Ala Thr Gly Arg Asn Ala Ala Arg Leu
50 55 60

Cys Cys Val Ala Ser Arg Asp Ala Gly Leu Ser Trp Gly Ser Ala Arg
65 70 75 80

Asp Leu Thr Glu Glu Ala Ile Gly Gly Ala Val Gln Asp Trp Ala Thr
85 90 95

Phe Ala Val Gly Pro Gly His Gly Val Gln Leu Pro Ser Gly Arg Leu
100 105 110

Leu Val Pro Ala Tyr Thr Tyr Arg Val Asp Arg Leu Glu Cys Phe Gly
 115 120 125
 Lys Ile Cys Arg Thr Ser Pro His Ser Phe Ala Phe Tyr Ser Asp Asp
 130 135 140
 His Gly Arg Thr Trp Arg Cys Gly Gly Leu Val Pro Asn Leu Arg Ser
 145 150 155 160
 Gly Glu Cys Gln Leu Ala Ala Val Asp Gly Gly Gln Ala Gly Ser Phe
 165 170 175
 Leu Tyr Cys Asn Ala Arg Ser Pro Leu Gly Ser Arg Val Gln Ala Leu
 180 185 190
 Ser Thr Asp Glu Gly Thr Ser Phe Leu Pro Ala Glu Arg Val Ala Ser
 195 200 205
 Leu Pro Glu Thr Ala Trp Gly Cys Gln Gly Ser Ile Val Gly Phe Pro
 210 215 220
 Ala Pro Ala Pro Asn Arg Pro Arg Asp Asp Ser Trp Ser Val Gly Pro
 225 230 235 240
 Arg Ser Pro Leu Gln Pro Pro Leu Leu Gly Pro Gly Val His Glu Pro
 245 250 255
 Pro Glu Glu Ala Ala Val Asp Pro Arg Gly Gly Gln Val Pro Gly Gly
 260 265 270
 Pro Phe Ser Arg Leu Gln Pro Arg Gly Asp Gly Pro Arg Gln Pro Gly
 275 280 285
 Pro Arg Pro Gly Val Ser Gly Asp Val Gly Ser Trp Thr Leu Ala Leu
 290 295 300
 Pro Met Pro Phe Ala Ala Pro Pro Gln Ser Pro Thr Trp Leu Leu Tyr
 305 310 315 320
 Ser His Pro Val Gly Arg Arg Ala Arg Leu His Met Gly Ile Arg Leu
 325 330 335

Ser Gln Ser Pro Leu Asp Pro Arg Ser Trp Thr Glu Pro Trp Val Ile
340 345 350

Tyr Glu Gly Pro Ser Gly Tyr Ser Asp Leu Ala Ser Ile Gly Pro Ala
355 360 365

Pro Glu Gly Gly Leu Val Phe Ala Cys Leu Tyr Glu Ser Gly Ala Arg
370 375 380

Thr Ser Tyr Asp Glu Ile Ser Phe Cys Thr Phe Ser Leu Arg Glu Val
385 390 395 400

Leu Glu Asn Val Pro Ala Ser Pro Lys Pro Pro Asn Leu Gly Asp Lys
405 410 415

Pro Arg Gly Cys Cys Trp Pro Ser
420

<210> 10
<211> 5
<212> PRT
<213> Artificial sequence

<220>
<223> Synthetic Construct

<400> 10

Gly Gly Gly Gly Ser
1 5

<210> 11
<211> 2706
<212> DNA
<213> Actinomyces viscosus

<400> 11
atgacatcgc atagtccttt ctcccggagg cgctgccgg cctcctggg ctccctgcca 60
ctggccgcca ccggcctgat cgccgccgca ccccggcgc acgcgctcc cagctctgac 120
ggcctggccg acgtcaccat cagcaggtg aacgcgccc cgacggcct ctactccgtc 180
ggcgatgtca tgaccttcaa catcaccctg accaacacca gggcgaggc ccactcctac 240
gccccggcct cgacgaacct gtccgggaac gtctccaagt gccggtggcg caacgtcccg 300
gccgggacga ccaagaccga ctgcaccggc ctggccacgc acacggtgac cgccgaggac 360

ctcaaggccg	gtggcttcac	cccgagatc	gcctacgagg	tcaaggccgt	ggagtacgcc	420
gggaaggccc	tgagcacccc	ggagacgac	aaggcgcgga	cgagcccagt	caaggccaac	480
tcgctgcggg	tcgagtcgat	cacgccgtcg	tcgagccagg	agaactacaa	gctgggcgac	540
accgtcagct	acacggtgcg	cgtgcgctcg	gtgtcggaca	agacgatcaa	cgtcgccgcc	600
accgaatcct	ccttcgacga	cctgggcccgc	cagtgccact	ggggcggcct	caagccgggc	660
aaggcgccg	tctacaactg	caagccgctc	acccacacga	tcacgcaagc	cgacgtcgac	720
gccggccgct	ggacgccatc	gatcaccctg	acggccaccg	gaaccgacgg	cgccaccctc	780
cagacgctca	ccgccaccgg	caacccgac	aacgtcgtcg	gcgaccaccc	gcaggccacg	840
cccgacccgg	cgcccagcgc	gagcacggag	ctgccggcct	caatgagcca	ggcccagcac	900
ctggccgcca	acacggccac	cgacaactac	cgcaccccgg	cgatcaccac	cgcccccaat	960
ggggacctgc	tcatctccta	cgacgagcgc	ccgaaggaca	acggcaacgg	cggcagcgac	1020
gcccccaacc	cgaaccacat	cgtccagcgc	cgctccaccg	acggcgga	gacctggtcg	1080
gcgcccacct	acatccacca	gggcacggag	accggcaaga	aggtcggcta	ctccgacccg	1140
agctacgtcg	tcgatcacca	gacgggcaag	atcttcaact	tccacgtcaa	gtcctacgac	1200
cagggtcggg	gcggctcgcg	cgccggcacc	gacccggaga	accggggcat	catccaggcc	1260
gaggtgtcga	cctccacgga	caacggctgg	acctggacgc	accgcacgat	caccgcggac	1320
atcacgaagg	acaagccgtg	gaccgcgcgt	ttcgcgccct	cgggccaggg	catccagatt	1380
cagcacgggc	cccacgccgg	gcgcctggtg	cagcagtaca	cgatcaggac	cgccggcggc	1440
gcggtgcagg	ccgtctcggt	ctactccgac	gaccacggga	agacgtggca	ggccggcacg	1500
ccgatcggga	ccggcatgga	tgagaacaag	gtcgttgagc	tctccgacgg	ctccctcatg	1560
ctcaactcgc	gcgcctcgga	tggtccgggc	ttccgcaagg	tggcccactc	caccgacggt	1620
gggcagacct	ggagcgagcc	ggtgtccgac	aagaacctgc	ccgactcggg	ggacaacgcc	1680
cagatcatcc	gagccttccc	gaacgcgcgc	ccggacgacc	cgcgcgccaa	ggtgctgctg	1740
ctgagccact	caccgaaccc	gcggccgtgg	tcgcgtgacc	gcggcaccat	ctcgatgtcc	1800
tgcgacgacg	gcgcctcctg	gacgaccagc	aaggtcttcc	acgagccctt	cgtcggatac	1860
acgacgatcg	cggtgcagtc	cgacggcagc	atcgggctgc	tcagcgagga	cggccacaac	1920
ggcgccgact	acggcgccat	ctggtaccgc	aacttcacga	tgaactggct	cggcgagcag	1980
tgcgcccaga	agccggcgga	gccgagcccg	gcgcgcgtcg	cgacggcggc	accctcagcg	2040
gcaccgacgg	agaagccggc	cccgtcggcc	gcgccgagcg	ctgagcccac	gcaggcaccg	2100

gcaccatcct ccgcgcccga gccgagcgct gcgcccgagc cgagcagcgc cccggcgccg 2160
gagcccaaga ccgctccgag cacggagccc acaccggctc ctgcgcccag ctccgcacct 2220
gagcagaccg atggggccgac cgctgcgccc gcaccggaga cgtcctctgc accggccgcc 2280
gaaccgacgc aggccccgac ggtggcgccct tctgttgagc ccacgcaggc tccgggtgcg 2340
cagccgagct cagcacccaa gccggggggcg acgggtcggg ccccgctcgtt ggtgaaccgc 2400
aaggcgaccg gggcgggcgac ggagcctggg acgccgtcat cgagcgcgag cccggcaccg 2460
agccggaacg cggcgccgac gccgaagccg ggcatggagc ccgatgagat tgatcggccg 2520
tctgacggca ccatggcgca gccgaccggt ggcgccagcg cgccgagtgc cgcgccgacg 2580
caggcggcga aggcggcgac caggctgtct cgcacgggga ccaacgcgct gctgatcctg 2640
ggccttgccg gtgtcgcggt tgtcgccggg tacctgctgc tgcgggctcg ccgttcgaag 2700
aactga 2706

<210> 12
<211> 901
<212> PRT
<213> Actinomyces viscosus

<400> 12

Met Thr Ser His Ser Pro Phe Ser Arg Arg Arg Leu Pro Ala Leu Leu
1 5 10 15

Gly Ser Leu Pro Leu Ala Ala Thr Gly Leu Ile Ala Ala Ala Pro Pro
20 25 30

Ala His Ala Val Pro Thr Ser Asp Gly Leu Ala Asp Val Thr Ile Thr
35 40 45

Gln Val Asn Ala Pro Ala Asp Gly Leu Tyr Ser Val Gly Asp Val Met
50 55 60

Thr Phe Asn Ile Thr Leu Thr Asn Thr Ser Gly Glu Ala His Ser Tyr
65 70 75 80

Ala Pro Ala Ser Thr Asn Leu Ser Gly Asn Val Ser Lys Cys Arg Trp
85 90 95

Arg Asn Val Pro Ala Gly Thr Thr Lys Thr Asp Cys Thr Gly Leu Ala
100 105 110

Thr His Thr Val Thr Ala Glu Asp Leu Lys Ala Gly Gly Phe Thr Pro
115 120 125

Gln Ile Ala Tyr Glu Val Lys Ala Val Glu Tyr Ala Gly Lys Ala Leu
130 135 140

Ser Thr Pro Glu Thr Ile Lys Gly Ala Thr Ser Pro Val Lys Ala Asn
145 150 155 160

Ser Leu Arg Val Glu Ser Ile Thr Pro Ser Ser Ser Gln Glu Asn Tyr
165 170 175

Lys Leu Gly Asp Thr Val Ser Tyr Thr Val Arg Val Arg Ser Val Ser
180 185 190

Asp Lys Thr Ile Asn Val Ala Ala Thr Glu Ser Ser Phe Asp Asp Leu
195 200 205

Gly Arg Gln Cys His Trp Gly Gly Leu Lys Pro Gly Lys Gly Ala Val
210 215 220

Tyr Asn Cys Lys Pro Leu Thr His Thr Ile Thr Gln Ala Asp Val Asp
225 230 235 240

Ala Gly Arg Trp Thr Pro Ser Ile Thr Leu Thr Ala Thr Gly Thr Asp
245 250 255

Gly Ala Thr Leu Gln Thr Leu Thr Ala Thr Gly Asn Pro Ile Asn Val
260 265 270

Val Gly Asp His Pro Gln Ala Thr Pro Ala Pro Ala Pro Asp Ala Ser
275 280 285

Thr Glu Leu Pro Ala Ser Met Ser Gln Ala Gln His Leu Ala Ala Asn
290 295 300

Thr Ala Thr Asp Asn Tyr Arg Ile Pro Ala Ile Thr Thr Ala Pro Asn
305 310 315 320

Gly Asp Leu Leu Ile Ser Tyr Asp Glu Arg Pro Lys Asp Asn Gly Asn
325 330 335

Gly Gly Ser Asp Ala Pro Asn Pro Asn His Ile Val Gln Arg Arg Ser
 340 345 350

Thr Asp Gly Gly Lys Thr Trp Ser Ala Pro Thr Tyr Ile His Gln Gly
 355 360 365

Thr Glu Thr Gly Lys Lys Val Gly Tyr Ser Asp Pro Ser Tyr Val Val
 370 375 380

Asp His Gln Thr Gly Thr Ile Phe Asn Phe His Val Lys Ser Tyr Asp
 385 390 395 400

Gln Gly Trp Gly Gly Ser Arg Gly Gly Thr Asp Pro Glu Asn Arg Gly
 405 410 415

Ile Ile Gln Ala Glu Val Ser Thr Ser Thr Asp Asn Gly Trp Thr Trp
 420 425 430

Thr His Arg Thr Ile Thr Ala Asp Ile Thr Lys Asp Lys Pro Trp Thr
 435 440 445

Ala Arg Phe Ala Ala Ser Gly Gln Gly Ile Gln Ile Gln His Gly Pro
 450 455 460

His Ala Gly Arg Leu Val Gln Gln Tyr Thr Ile Arg Thr Ala Gly Gly
 465 470 475 480

Ala Val Gln Ala Val Ser Val Tyr Ser Asp Asp His Gly Lys Thr Trp
 485 490 495

Gln Ala Gly Thr Pro Ile Gly Thr Gly Met Asp Glu Asn Lys Val Val
 500 505 510

Glu Leu Ser Asp Gly Ser Leu Met Leu Asn Ser Arg Ala Ser Asp Gly
 515 520 525

Ser Gly Phe Arg Lys Val Ala His Ser Thr Asp Gly Gly Gln Thr Trp
 530 535 540

Ser Glu Pro Val Ser Asp Lys Asn Leu Pro Asp Ser Val Asp Asn Ala
 545 550 555 560

Gln Ile Ile Arg Ala Phe Pro Asn Ala Ala Pro Asp Asp Pro Arg Ala
 565 570 575

Lys Val Leu Leu Leu Ser His Ser Pro Asn Pro Arg Pro Trp Ser Arg
 580 585 590

Asp Arg Gly Thr Ile Ser Met Ser Cys Asp Asp Gly Ala Ser Trp Thr
 595 600 605

Thr Ser Lys Val Phe His Glu Pro Phe Val Gly Tyr Thr Thr Ile Ala
 610 615 620

Val Gln Ser Asp Gly Ser Ile Gly Leu Leu Ser Glu Asp Ala His Asn
 625 630 635 640

Gly Ala Asp Tyr Gly Gly Ile Trp Tyr Arg Asn Phe Thr Met Asn Trp
 645 650 655

Leu Gly Glu Gln Cys Gly Gln Lys Pro Ala Glu Pro Ser Pro Ala Pro
 660 665 670

Ser Pro Thr Ala Ala Pro Ser Ala Ala Pro Thr Glu Lys Pro Ala Pro
 675 680 685

Ser Ala Ala Pro Ser Ala Glu Pro Thr Gln Ala Pro Ala Pro Ser Ser
 690 695 700

Ala Pro Glu Pro Ser Ala Ala Pro Glu Pro Ser Ser Ala Pro Ala Pro
 705 710 715 720

Glu Pro Thr Thr Ala Pro Ser Thr Glu Pro Thr Pro Ala Pro Ala Pro
 725 730 735

Ser Ser Ala Pro Glu Gln Thr Asp Gly Pro Thr Ala Ala Pro Ala Pro
 740 745 750

Glu Thr Ser Ser Ala Pro Ala Ala Glu Pro Thr Gln Ala Pro Thr Val
 755 760 765

Ala Pro Ser Val Glu Pro Thr Gln Ala Pro Gly Ala Gln Pro Ser Ser
 770 775 780

Ala Pro Lys Pro Gly Ala Thr Gly Arg Ala Pro Ser Val Val Asn Pro
785 790 795 800

Lys Ala Thr Gly Ala Ala Thr Glu Pro Gly Thr Pro Ser Ser Ser Ala
805 810 815

Ser Pro Ala Pro Ser Arg Asn Ala Ala Pro Thr Pro Lys Pro Gly Met
820 825 830

Glu Pro Asp Glu Ile Asp Arg Pro Ser Asp Gly Thr Met Ala Gln Pro
835 840 845

Thr Gly Gly Ala Ser Ala Pro Ser Ala Ala Pro Thr Gln Ala Ala Lys
850 855 860

Ala Gly Ser Arg Leu Ser Arg Thr Gly Thr Asn Ala Leu Leu Ile Leu
865 870 875 880

Gly Leu Ala Gly Val Ala Val Val Gly Gly Tyr Leu Leu Leu Arg Ala
885 890 895

Arg Arg Ser Lys Asn
900

<210> 13
<211> 1887
<212> DNA
<213> Actinomyces viscosus

<400> 13
ggcgaccacc cgcaggccac gcccgccacc gcgcccgcac cgagcacgga gctgccggcc 60
tcaatgagcc aggcccagca cctggccgcc aacacggcca ccgacaacta ccgcatcccg 120
gcgatcacca ccgcccccaa tggggacctg ctcattctct acgacgagcg ccggaaggac 180
aacggcaacg gcggcagcga cgcgcccaac ccgaaccaca tcgtccagcg ccgctccacc 240
gacggcggca agacctggtc ggcgcccacc tacatccacc agggcacgga gaccggcaag 300
aaggctggct actccgaccc gagctacgtc gtcgatcacc agacgggcac gatcttcaac 360
ttccacgtca agtcctacga ccagggctgg ggcggctcgc gcggcggcac cgacccggag 420
aaccggggca tcatccaggc cgaggtgtcg acctccacgg acaacggctg gacctggacg 480
caccgcacga tcaccgcgga catcacgaag gacaagccgt ggaccgcgcy tttcgcggcc 540
tcgggccagg gcatccagat tcagcacggg cccacgcgcg ggcgcctggt gcagcagtac 600

```

acgatcagga ccgccggcgg cgcggtgcag gccgtctcgg tctactccga cgaccacggg 660
aagacgtggc aggccggcac gccgatcggg accggcatgg atgagaacaa ggtcgttgag 720
ctctccgacg gctccctcat gctcaactcg cgcgcctcgg atgggtccgg cttccgcaag 780
gtggcccact ccaccgacgg tgggcagacc tggagcgagc cgggtgtccga caagaacctg 840
cccgactcgg tggacaacgc ccagatcatc cgagccttcc cgaacgccgc gccggacgac 900
ccgcgcgcca aggtgctgct gctgagccac tcaccgaacc cgcggccgtg gtcgcgtgac 960
cgcggcacca tctcgatgtc ctgcgacgac ggcgcctcct ggacgaccag caaggctctc 1020
cacgagccct tcgtcggata cacgacgacg cgcggtgcagt ccgacggcag catcgggctg 1080
ctcagcgagg acgcccacaa cggcgccgac tacggcgga tctggtaccg caacttcacg 1140
atgaactggc tcggcgagca gtgcggccag aagccggcgg agccgagccc ggcgccgtcg 1200
ccgacggcgg caccctcagc ggcaccgacg gagaagccgg ccccgtcggc cgcgccgagc 1260
gctgagccca cgcaggcacc ggcaccatcc tccgcgcccg agccgagcgc tgcgcccag 1320
ccgagcagcg ccccggcgcc ggagcccacg accgctccga gcacggagcc cacaccggct 1380
cctgcgcca gctccgcacc tgagcagacc gatgggccga ccgctgcgcc cgcaccggag 1440
acgtcctctg caccggccgc cgaaccgacg caggccccga cgggtggcgc ttctgttgag 1500
cccacgcagg ctccgggtgc gcagccgagc tcagaccca agccgggggc gacgggtcgg 1560
gccccgtcgg tggatgaacc gaaggcgacc gggcgggcga cggagcctgg gacgccgtca 1620
tcgagcgca gcccggcacc gagccggaac gcggcgccga cgcgaagcc gggcatggag 1680
cccgatgaga ttgatcggcc gtctgacggc accatggcgc agccgaccgg tggcgccagc 1740
gcgccgagtg ccgcgccgac gcaggcgggc aaggccggca gcaggctgtc tcgcacgggg 1800
accaacgcgc tgctgatcct gggccttgcg ggtgtcgcgg ttgtcggcgg gtacctgctg 1860
ctgcgggctc gccgttcgaa gaactga 1887

```

<210> 14

<211> 628

<212> PRT

<213> *Actinomyces viscosus*

<400> 14

```

Gly Asp His Pro Gln Ala Thr Pro Ala Pro Ala Pro Asp Ala Ser Thr
1           5           10          15

```

Glu Leu Pro Ala Ser Met Ser Gln Ala Gln His Leu Ala Ala Asn Thr
20 25 30

Ala Thr Asp Asn Tyr Arg Ile Pro Ala Ile Thr Thr Ala Pro Asn Gly
35 40 45

Asp Leu Leu Ile Ser Tyr Asp Glu Arg Pro Lys Asp Asn Gly Asn Gly
50 55 60

Gly Ser Asp Ala Pro Asn Pro Asn His Ile Val Gln Arg Arg Ser Thr
65 70 75 80

Asp Gly Gly Lys Thr Trp Ser Ala Pro Thr Tyr Ile His Gln Gly Thr
85 90 95

Glu Thr Gly Lys Lys Val Gly Tyr Ser Asp Pro Ser Tyr Val Val Asp
100 105 110

His Gln Thr Gly Thr Ile Phe Asn Phe His Val Lys Ser Tyr Asp Gln
115 120 125

Gly Trp Gly Gly Ser Arg Gly Gly Thr Asp Pro Glu Asn Arg Gly Ile
130 135 140

Ile Gln Ala Glu Val Ser Thr Ser Thr Asp Asn Gly Trp Thr Trp Thr
145 150 155 160

His Arg Thr Ile Thr Ala Asp Ile Thr Lys Asp Lys Pro Trp Thr Ala
165 170 175

Arg Phe Ala Ala Ser Gly Gln Gly Ile Gln Ile Gln His Gly Pro His
180 185 190

Ala Gly Arg Leu Val Gln Gln Tyr Thr Ile Arg Thr Ala Gly Gly Ala
195 200 205

Val Gln Ala Val Ser Val Tyr Ser Asp Asp His Gly Lys Thr Trp Gln
210 215 220

Ala Gly Thr Pro Ile Gly Thr Gly Met Asp Glu Asn Lys Val Val Glu
225 230 235 240

Leu Ser Asp Gly Ser Leu Met Leu Asn Ser Arg Ala Ser Asp Gly Ser
245 250 255

Gly Phe Arg Lys Val Ala His Ser Thr Asp Gly Gly Gln Thr Trp Ser
260 265 270

Glu Pro Val Ser Asp Lys Asn Leu Pro Asp Ser Val Asp Asn Ala Gln
275 280 285

Ile Ile Arg Ala Phe Pro Asn Ala Ala Pro Asp Asp Pro Arg Ala Lys
290 295 300

Val Leu Leu Leu Ser His Ser Pro Asn Pro Arg Pro Trp Ser Arg Asp
305 310 315 320

Arg Gly Thr Ile Ser Met Ser Cys Asp Asp Gly Ala Ser Trp Thr Thr
325 330 335

Ser Lys Val Phe His Glu Pro Phe Val Gly Tyr Thr Thr Ile Ala Val
340 345 350

Gln Ser Asp Gly Ser Ile Gly Leu Leu Ser Glu Asp Ala His Asn Gly
355 360 365

Ala Asp Tyr Gly Gly Ile Trp Tyr Arg Asn Phe Thr Met Asn Trp Leu
370 375 380

Gly Glu Gln Cys Gly Gln Lys Pro Ala Glu Pro Ser Pro Ala Pro Ser
385 390 395 400

Pro Thr Ala Ala Pro Ser Ala Ala Pro Thr Glu Lys Pro Ala Pro Ser
405 410 415

Ala Ala Pro Ser Ala Glu Pro Thr Gln Ala Pro Ala Pro Ser Ser Ala
420 425 430

Pro Glu Pro Ser Ala Ala Pro Glu Pro Ser Ser Ala Pro Ala Pro Glu
435 440 445

Pro Thr Thr Ala Pro Ser Thr Glu Pro Thr Pro Ala Pro Ala Pro Ser
450 455 460

Ser Ala Pro Glu Gln Thr Asp Gly Pro Thr Ala Ala Pro Ala Pro Glu
465 470 475 480

Thr Ser Ser Ala Pro Ala Ala Glu Pro Thr Gln Ala Pro Thr Val Ala
485 490 495

Pro Ser Val Glu Pro Thr Gln Ala Pro Gly Ala Gln Pro Ser Ser Ala
500 505 510

Pro Lys Pro Gly Ala Thr Gly Arg Ala Pro Ser Val Val Asn Pro Lys
515 520 525

Ala Thr Gly Ala Ala Thr Glu Pro Gly Thr Pro Ser Ser Ser Ala Ser
530 535 540

Pro Ala Pro Ser Arg Asn Ala Ala Pro Thr Pro Lys Pro Gly Met Glu
545 550 555 560

Pro Asp Glu Ile Asp Arg Pro Ser Asp Gly Thr Met Ala Gln Pro Thr
565 570 575

Gly Gly Ala Ser Ala Pro Ser Ala Ala Pro Thr Gln Ala Ala Lys Ala
580 585 590

Gly Ser Arg Leu Ser Arg Thr Gly Thr Asn Ala Leu Leu Ile Leu Gly
595 600 605

Leu Ala Gly Val Ala Val Val Gly Gly Tyr Leu Leu Leu Arg Ala Arg
610 615 620

Arg Ser Lys Asn
625

<210> 15
<211> 1182
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<400> 15
ggagatcatc cacaagctac accagcacct gcaccagatg ctagcactga gctgccagca 60
agcatgtctc aggctcagca tcttgacgca aatacggcta ctgataatta tcgcattcca 120

gcgattacaa ccgctccgaa tgggtgattta ctgattagct atgatgaacg gccgaaggac 180
 aatggaaatg gtggttccga tgcccctaac ccgaatcata ttgttcagcg tcgctccaca 240
 gatggcggtg aaacttggag cgcgccaacc tatattcatc agggtagcga gactggcaag 300
 aaagtgggat attccgaccc ctcttatgtg gtggatcatc aaaccggtac aatcttcaat 360
 tttcatgtga aatcatacga tcagggtctgg ggaggtagcc gtgggggaac agaccggaa 420
 aaccgcggga ttattcaggc agaggtgtct acgagcacgg ataatggatg gacgtggaca 480
 catcgcacca tcaccgcgga tattacgaaa gataaaccgt ggaccgcgcg ttttgcgcg 540
 tccggccaag gcattcagat ccagcatggg ccgcatgccg gccgtctggt gcaacagtat 600
 accattcgta cggccggtgg agcgggtgcag gctgtatcgg tttattccga tgatcatggg 660
 aaaacgtggc aggctggcac cccgattggg acgggtatgg atgaaaacaa agttgtagag 720
 ctgtctgacg gctctctgat gctgaacagt cgtgcgtcgg acgggagcgg ctttcgtaag 780
 gttgcgcata gcactgatgg tgggcagacc tgggtccgaac cggtttcgga caaaaatttg 840
 ccggattcgg ttgataatgc ccagataatt cgtgcgttcc ctaatgctgc ccccgatgac 900
 ccgcgcgcga aagtacttct tctgagtcac tccccaaatc cacgtccgtg gtcccgggat 960
 cgtggtacga taagcatgtc atgtgatgac ggggcctcat ggaccacttc caaagttttt 1020
 cacgaaccgt ttgtgggcta cacgactatt gcagttcaga gtgatggaag catcgtctcg 1080
 ctgtcggagg acgcgcacaa tggcgctgat tatggtggca tctggtatcg taattttacg 1140
 atgaactggc tgggagaaca atgtggacaa aaaccgcgga aa 1182

<210> 16
 <211> 394
 <212> PRT
 <213> Actinomyces viscosus

<400> 16

Gly Asp His Pro Gln Ala Thr Pro Ala Pro Ala Pro Asp Ala Ser Thr
 1 5 10 15

Glu Leu Pro Ala Ser Met Ser Gln Ala Gln His Leu Ala Ala Asn Thr
 20 25 30

Ala Thr Asp Asn Tyr Arg Ile Pro Ala Ile Thr Thr Ala Pro Asn Gly
 35 40 45

Asp Leu Leu Ile Ser Tyr Asp Glu Arg Pro Lys Asp Asn Gly Asn Gly
 50 55 60

Gly Ser Asp Ala Pro Asn Pro Asn His Ile Val Gln Arg Arg Ser Thr
 65 70 75 80

Asp Gly Gly Lys Thr Trp Ser Ala Pro Thr Tyr Ile His Gln Gly Thr
 85 90 95

Glu Thr Gly Lys Lys Val Gly Tyr Ser Asp Pro Ser Tyr Val Val Asp
 100 105 110

His Gln Thr Gly Thr Ile Phe Asn Phe His Val Lys Ser Tyr Asp Gln
 115 120 125

Gly Trp Gly Gly Ser Arg Gly Gly Thr Asp Pro Glu Asn Arg Gly Ile
 130 135 140

Ile Gln Ala Glu Val Ser Thr Ser Thr Asp Asn Gly Trp Thr Trp Thr
 145 150 155 160

His Arg Thr Ile Thr Ala Asp Ile Thr Lys Asp Lys Pro Trp Thr Ala
 165 170 175

Arg Phe Ala Ala Ser Gly Gln Gly Ile Gln Ile Gln His Gly Pro His
 180 185 190

Ala Gly Arg Leu Val Gln Gln Tyr Thr Ile Arg Thr Ala Gly Gly Ala
 195 200 205

Val Gln Ala Val Ser Val Tyr Ser Asp Asp His Gly Lys Thr Trp Gln
 210 215 220

Ala Gly Thr Pro Ile Gly Thr Gly Met Asp Glu Asn Lys Val Val Glu
 225 230 235 240

Leu Ser Asp Gly Ser Leu Met Leu Asn Ser Arg Ala Ser Asp Gly Ser
 245 250 255

Gly Phe Arg Lys Val Ala His Ser Thr Asp Gly Gly Gln Thr Trp Ser
 260 265 270

Glu Pro Val Ser Asp Lys Asn Leu Pro Asp Ser Val Asp Asn Ala Gln
 275 280 285

Ile Ile Arg Ala Phe Pro Asn Ala Ala Pro Asp Asp Pro Arg Ala Lys
 290 295 300

Val Leu Leu Leu Ser His Ser Pro Asn Pro Arg Pro Trp Ser Arg Asp
 305 310 315 320

Arg Gly Thr Ile Ser Met Ser Cys Asp Asp Gly Ala Ser Trp Thr Thr
 325 330 335

Ser Lys Val Phe His Glu Pro Phe Val Gly Tyr Thr Thr Ile Ala Val
 340 345 350

Gln Ser Asp Gly Ser Ile Gly Leu Leu Ser Glu Asp Ala His Asn Gly
 355 360 365

Ala Asp Tyr Gly Gly Ile Trp Tyr Arg Asn Phe Thr Met Asn Trp Leu
 370 375 380

Gly Glu Gln Cys Gly Gln Lys Pro Ala Glu
 385 390

<210> 17
 <211> 6
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 17

Met Val Lys Arg Lys Lys
 1 5

<210> 18
 <211> 1281
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 18
 ccattggttaa ggcgaacaaaa aaaggcggca aaaacggtaa aaatcgtcgt aaccgtaaga 60

```

aaaaaaatcc tggagatcat ccacaagcta caccagcacc tgcaccagat gctagcactg      120
agctgccagc aagcatgtct caggctcagc atcttgacgc aaatacggct actgataatt      180
atcgcatcc agcgattaca accgctccga atgggtgattt actgattagc tatgatgaac      240
ggccgaagga caatggaaat ggtgggtccg atgccctaa cccgaatcat attgttcagc      300
gtcgctccac agatggcggg aaaacttgga gcgcgccaac ctatattcat cagggtacgg      360
agactggcaa gaaagtggga tattccgacc cctcttatgt ggtggatcat caaaccggta      420
caatcttcaa ttttcatgtg aaatcatacg atcagggtcg gggaggtagc cgtgggggaa      480
cagacccgga aaaccgcggg attattcagg cagaggtgtc tacgagcacg gataatggat      540
ggacgtggac acatcgcacc atcaccgcgg atattacgaa agataaaccg tggaccgcgc      600
gttttgccgc gtccggccaa ggcattcaga tccagcatgg gccgcgtgcc ggccgtctgg      660
tgcaacagta taccattcgt acggccgggtg gagcgggtgca ggctgtatcg gtttattccg      720
atgatcatgg gaaaacgtgg caggctggca ccccgattgg gacgggtatg gatgaaaaca      780
aagttgtaga gctgtctgac ggctctctga tgctgaacag tcgtgcgtcg gacgggagcg      840
gctttcgtaa ggttgcgcat agcactgatg gtgggcagac ctggtccgaa ccggtttcgg      900
acaaaaatth gccggattcg gttgataatg cccagataat tcgtgcgttt cctaattgctg      960
ccccgatga ccgcgcgcgc aaagtacttc ttctgagtca ttccccaat ccacgtccgt     1020
ggtcccggga tcgtggtacg ataagcatgt catgtgatga cggggcctca tggaccactt     1080
ccaaagttht tcacgaaccg tttgtgggct acacgactat tgcagttcag agtgatggaa     1140
gcatcggtct gctgtcggag gacgcgcaca atggcgctga ttatggtggc atctggtatc     1200
gtaatthttac gatgaactgg ctgggagaac aatgtggaca aaaaccgcgc gaataagctt     1260
aaaaaccgcg ggaataagct t                                             1281

```

<210> 19
 <211> 444
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 19

```

Met Val Lys Arg Lys Lys Lys Gly Gly Lys Asn Gly Lys Asn Arg Arg
1          5          10          15

```

Asn Arg Lys Lys Lys Asn Pro Gly Asp His Pro Gln Ala Thr Pro Ala
 20 25 30

Pro Ala Pro Asp Ala Ser Thr Glu Leu Pro Ala Ser Met Ser Gln Ala
 35 40 45

Gln His Leu Ala Ala Asn Thr Ala Thr Asp Asn Tyr Arg Ile Pro Ala
 50 55 60

Ile Thr Thr Ala Pro Asn Gly Asp Leu Leu Ile Ser Tyr Asp Glu Arg
 65 70 75 80

Pro Lys Asp Asn Gly Asn Gly Gly Ser Asp Ala Pro Asn Pro Asn His
 85 90 95

Ile Val Gln Arg Arg Ser Thr Asp Gly Gly Lys Thr Trp Ser Ala Pro
 100 105 110

Thr Tyr Ile His Gln Gly Thr Glu Thr Gly Lys Lys Val Gly Tyr Ser
 115 120 125

Asp Pro Ser Tyr Val Val Asp His Gln Thr Gly Thr Ile Phe Asn Phe
 130 135 140

His Val Lys Ser Tyr Asp Gln Gly Trp Gly Gly Ser Arg Gly Gly Thr
 145 150 155 160

Asp Pro Glu Asn Arg Gly Ile Ile Gln Ala Glu Val Ser Thr Ser Thr
 165 170 175

Asp Asn Gly Trp Thr Trp Thr His Arg Thr Ile Thr Ala Asp Ile Thr
 180 185 190

Lys Asp Lys Pro Trp Thr Ala Arg Phe Ala Ala Ser Gly Gln Gly Ile
 195 200 205

Gln Ile Gln His Gly Pro His Ala Gly Arg Leu Val Gln Gln Tyr Thr
 210 215 220

Ile Arg Thr Ala Gly Gly Ala Val Gln Ala Val Ser Val Tyr Ser Asp
 225 230 235 240

```
<210> 20
<211> 1248
<212> DNA
<213> Artificial Sequence
```

<220>

<223> Synthetic Construct

<400> 20

```
atgggagatc atccacaagc tacaccagca cctgcaccag atgctagcac tgagctgccca      60
gcaagcatgt ctcaggctca gcatcttgca gcaaatacgg ctactgataa ttatcgcatc      120
ccagcgatta caaccgctcc gaatgggtgat ttactgatta gctatgatga acggccgaag      180
gacaatggaa atgggtggttc cgatgcccct aaccggaatc atattgttca gcgtcgctcc      240
acagatggcg gtaaaacttg gagcgcgcca acctatattc atcagggtac ggagactggc      300
aagaaaagtg gatattccga cccctcttat gtggtggatc atcaaaccgg tacaatcttc      360
aattttcatg tgaaatcata cgatcagggc tggggaggta gccgtggggg aacagaccgc      420
gaaaaccgcg ggattattca ggcagagggtg tctacgagca cggataatgg atggacgtgg      480
acacatcgca ccatcacgcg ggatattacg aaagataaac cgtggaccgc gcgttttgcg      540
gcgtccggcc aaggcattca gatccagcat gggccgcatg ccggccgtct ggtgcaacag      600
tataccattc gtacggccgg tggagcgggtg caggctgtat cggtttattc cgatgatcat      660
gggaaaacgt ggcaggctgg caccocgatt gggacgggta tggatgaaaa caaagtgtga      720
gagctgtctg acggctctct gatgctgaac agtcgtgcgt cggacgggag cggctttcgt      780
aaggttgcgc atagcactga tgggtgggcag acctggtccg aaccggtttc ggacaaaaat      840
ttgccggatt cggttgataa tgcccagata attcgtgcgt ttcctaatgc tgccccgat      900
gaccgcgcgc cgaaagtact tcttctgagt cattcccaa atccacgtcc gtggtcccgg      960
gatcgtggta cgataagcat gtcatgtgat gacggggcct catggaccac ttccaaagtt     1020
tttcacgaac cgtttgtggg ctacacgact attgcagttc agagtgatgg aagcatcggc     1080
ctgctgtcgg aggacgcgca caatggcgct gattatggtg gcatctggta tcgtaatttt     1140
acgatgaact ggctgggaga acaatgtgga caaaaaccg cgaagcgcaa aaaaaaaggc     1200
ggcaaaaacg gtaaaaatcg tcgtaaccgt aagaaaaaaa atccttga      1248
```

<210> 21

<211> 415

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 21

Met Gly Asp His Pro Gln Ala Thr Pro Ala Pro Ala Pro Asp Ala Ser
 1 5 10 15
 Thr Glu Leu Pro Ala Ser Met Ser Gln Ala Gln His Leu Ala Ala Asn
 20 25 30
 Thr Ala Thr Asp Asn Tyr Arg Ile Pro Ala Ile Thr Thr Ala Pro Asn
 35 40 45
 Gly Asp Leu Leu Ile Ser Tyr Asp Glu Arg Pro Lys Asp Asn Gly Asn
 50 55 60
 Gly Gly Ser Asp Ala Pro Asn Pro Asn His Ile Val Gln Arg Arg Ser
 65 70 75 80
 Thr Asp Gly Gly Lys Thr Trp Ser Ala Pro Thr Tyr Ile His Gln Gly
 85 90 95
 Thr Glu Thr Gly Lys Lys Val Gly Tyr Ser Asp Pro Ser Tyr Val Val
 100 105 110
 Asp His Gln Thr Gly Thr Ile Phe Asn Phe His Val Lys Ser Tyr Asp
 115 120 125
 Gln Gly Trp Gly Gly Ser Arg Gly Gly Thr Asp Pro Glu Asn Arg Gly
 130 135 140
 Ile Ile Gln Ala Glu Val Ser Thr Ser Thr Asp Asn Gly Trp Thr Trp
 145 150 155 160
 Thr His Arg Thr Ile Thr Ala Asp Ile Thr Lys Asp Lys Pro Trp Thr
 165 170 175
 Ala Arg Phe Ala Ala Ser Gly Gln Gly Ile Gln Ile Gln His Gly Pro
 180 185 190
 His Ala Gly Arg Leu Val Gln Gln Tyr Thr Ile Arg Thr Ala Gly Gly
 195 200 205
 Ala Val Gln Ala Val Ser Val Tyr Ser Asp Asp His Gly Lys Thr Trp
 210 215 220

Gln Ala Gly Thr Pro Ile Gly Thr Gly Met Asp Glu Asn Lys Val Val
 225 230 235 240

Glu Leu Ser Asp Gly Ser Leu Met Leu Asn Ser Arg Ala Ser Asp Gly
 245 250 255

Ser Gly Phe Arg Lys Val Ala His Ser Thr Asp Gly Gly Gln Thr Trp
 260 265 270

Ser Glu Pro Val Ser Asp Lys Asn Leu Pro Asp Ser Val Asp Asn Ala
 275 280 285

Gln Ile Ile Arg Ala Phe Pro Asn Ala Ala Pro Asp Asp Pro Arg Ala
 290 295 300

Lys Val Leu Leu Leu Ser His Ser Pro Asn Pro Arg Pro Trp Ser Arg
 305 310 315 320

Asp Arg Gly Thr Ile Ser Met Ser Cys Asp Asp Gly Ala Ser Trp Thr
 325 330 335

Thr Ser Lys Val Phe His Glu Pro Phe Val Gly Tyr Thr Thr Ile Ala
 340 345 350

Val Gln Ser Asp Gly Ser Ile Gly Leu Leu Ser Glu Asp Ala His Asn
 355 360 365

Gly Ala Asp Tyr Gly Gly Ile Trp Tyr Arg Asn Phe Thr Met Asn Trp
 370 375 380

Leu Gly Glu Gln Cys Gly Gln Lys Pro Ala Lys Arg Lys Lys Lys Gly
 385 390 395 400

Gly Lys Asn Gly Lys Asn Arg Arg Asn Arg Lys Lys Lys Asn Pro
 405 410 415

<210> 22
 <211> 1293
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 22

```

atgggagatc atccacaagc tacaccagca cctgcaccag atgctagcac tgagctgcca      60
gcaagcatgt ctcaggctca gcatcttgca gcaaatacgg ctactgataa ttatcgcatc      120
ccagcgatta caaccgctcc gaatggtgat ttactgatta gctatgatga acggccgaag      180
gacaatggaa atggtggttc cgatgcccct aaccgaatc atattgttca gcgtcgctcc      240
acagatggcg gtaaaacttg gagcgcgcca acctatattc atcagggtac ggagactggc      300
aagaaagtgg gatattccga cccctcttat gtggtggatc atcaaaccgg tacaatcttc      360
aattttcatg tgaaatcata cgatcagggc tggggaggta gccgtggggg aacagaccgc      420
gaaaaccgcg ggattattca ggagagggtg tctacgagca cggataatgg atggacgtgg      480
acacatcgca ccatcacgcg ggatattacg aaagataaac cgtggaccgc gcgttttgcg      540
gcgtccggcc aaggcattca gatccagcat gggccgcatg ccggccgtct ggtgcaacag      600
tataccattc gtacggccgg tggagcgggtg caggctgtat cggtttattc cgatgatcat      660
gggaaaacgt ggcaggctgg caccgccgatt gggacgggta tggatgaaaa caaagttgta      720
gagctgtctg acggctctct gatgctgaac agtcgtgctg cggacgggag cggctttcgt      780
aaggttgctc atagcactga tgggtgggcg acctggtccg aaccggtttc ggacaaaaat      840
ttgccggatt cggttgataa tgcccagata attcgtgctg ttctaatgc tgccccgat      900
gaccgcgcg cgaaagtact tcttctgagt cattcccaa atccacgtcc gtggtcccgg      960
gatcgtggta cgataagcat gtcattgtgat gacggggcct catggaccac ttccaaagtt     1020
tttcacgaac cgtttgtggg ctacacgact attgcagttc agagtgatgg aagcatcggt     1080
ctgctgtcgg aggacgcgca caatggcgct gattatggtg gcatctggta tcgtaatttt     1140
acgatgaact ggctgggaga acaatgtgga caaaaaccg cggaaccgag cccagcccct     1200
agccctactg cagcaccgtc cgctgcaaag cgcaaaaaaa aaggcggcaa aaacggtaaa     1260
aatcgtcgta accgtaagaa aaaaaatcct tga                                     1293

```

```

<210> 23
<211> 430
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Synthetic Construct

```

```

<400> 23

```

```

Met Gly Asp His Pro Gln Ala Thr Pro Ala Pro Ala Pro Asp Ala Ser
1           5           10           15

```

Thr Glu Leu Pro Ala Ser Met Ser Gln Ala Gln His Leu Ala Ala Asn
 20 25 30

Thr Ala Thr Asp Asn Tyr Arg Ile Pro Ala Ile Thr Thr Ala Pro Asn
 35 40 45

Gly Asp Leu Leu Ile Ser Tyr Asp Glu Arg Pro Lys Asp Asn Gly Asn
 50 55 60

Gly Gly Ser Asp Ala Pro Asn Pro Asn His Ile Val Gln Arg Arg Ser
 65 70 75 80

Thr Asp Gly Gly Lys Thr Trp Ser Ala Pro Thr Tyr Ile His Gln Gly
 85 90 95

Thr Glu Thr Gly Lys Lys Val Gly Tyr Ser Asp Pro Ser Tyr Val Val
 100 105 110

Asp His Gln Thr Gly Thr Ile Phe Asn Phe His Val Lys Ser Tyr Asp
 115 120 125

Gln Gly Trp Gly Gly Ser Arg Gly Gly Thr Asp Pro Glu Asn Arg Gly
 130 135 140

Ile Ile Gln Ala Glu Val Ser Thr Ser Thr Asp Asn Gly Trp Thr Trp
 145 150 155 160

Thr His Arg Thr Ile Thr Ala Asp Ile Thr Lys Asp Lys Pro Trp Thr
 165 170 175

Ala Arg Phe Ala Ala Ser Gly Gln Gly Ile Gln Ile Gln His Gly Pro
 180 185 190

His Ala Gly Arg Leu Val Gln Gln Tyr Thr Ile Arg Thr Ala Gly Gly
 195 200 205

Ala Val Gln Ala Val Ser Val Tyr Ser Asp Asp His Gly Lys Thr Trp
 210 215 220

Gln Ala Gly Thr Pro Ile Gly Thr Gly Met Asp Glu Asn Lys Val Val
 225 230 235 240

Glu Leu Ser Asp Gly Ser Leu Met Leu Asn Ser Arg Ala Ser Asp Gly
245 250 255

Ser Gly Phe Arg Lys Val Ala His Ser Thr Asp Gly Gly Gln Thr Trp
260 265 270

Ser Glu Pro Val Ser Asp Lys Asn Leu Pro Asp Ser Val Asp Asn Ala
275 280 285

Gln Ile Ile Arg Ala Phe Pro Asn Ala Ala Pro Asp Asp Pro Arg Ala
290 295 300

Lys Val Leu Leu Leu Ser His Ser Pro Asn Pro Arg Pro Trp Ser Arg
305 310 315 320

Asp Arg Gly Thr Ile Ser Met Ser Cys Asp Asp Gly Ala Ser Trp Thr
325 330 335

Thr Ser Lys Val Phe His Glu Pro Phe Val Gly Tyr Thr Thr Ile Ala
340 345 350

Val Gln Ser Asp Gly Ser Ile Gly Leu Leu Ser Glu Asp Ala His Asn
355 360 365

Gly Ala Asp Tyr Gly Gly Ile Trp Tyr Arg Asn Phe Thr Met Asn Trp
370 375 380

Leu Gly Glu Gln Cys Gly Gln Lys Pro Ala Glu Pro Ser Pro Ala Pro
385 390 395 400

Ser Pro Thr Ala Ala Pro Ser Ala Ala Lys Arg Lys Lys Lys Gly Gly
405 410 415

Lys Asn Gly Lys Asn Arg Arg Asn Arg Lys Lys Lys Asn Pro
420 425 430

<210> 24
<211> 1203
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Construct

```

<400> 24
atgggagagc tggcagcaag catgtctcag gctcagcatc ttgcagcaaa tacggctact      60
gataattatc gcattccagc gattacaacc gctccgaatg gtgattttact gattagctat      120
gatgaacggc cgaaggacaa tggaaatggg gggtccgatg cccctaaccg gaatcatatt      180
gttcagcgtc gctccacaga tggcggtaaa acttggagcg cgccaacctt tattcatcag      240
gggtacggaga ctggcaagaa agtgggatat tccgaccctt cttatgtggg ggatcatcaa      300
accggtacaa tcttcaattt tcatgtgaaa tcatacgatc agggctgggg aggtagccgt      360
gggggaacag acccggaaaa ccgcgggatt attcaggcag aggtgtctac gagcacggat      420
aatggatgga cgtggacaca tcgcaccatc acccgggata ttacgaaaga taaaccgtgg      480
accgcgcgtt ttgcggcgtc cggccaaggc attcagatcc agcatggggc gcatgccggc      540
cgtctgggtc aacagtatac cattcgtacg gccgggtggg cgggtgcaggc tgtatcgggt      600
tattccgatg atcatgggaa aacgtggcag gctggcaccg cgattggggc ggggatggat      660
gaaaacaaag ttgtagagct gtctgacggc tctctgatgc tgaacagtcg tgcgtcggac      720
gggagcgggt ttcgtaaggt tgcgcatagc actgatgggt ggcagacctg gtccgaaccg      780
gtttcggaca aaaatttgcc ggattcgggt gataatgcc agataattcg tgcgtttcct      840
aatgctgccc ccgatgacct gcgcgcgaaa gtacttcttc tgagtcattc cccaaatcca      900
cgtccgtggt cccgggatcg tggtagata agcatgtcat gtgatgacgg ggcctcatgg      960
accacttcca aagtttttca cgaaccgttt gtgggctaca cgactattgc agttcagagt     1020
gatggaagca tcgggtctgt gtcggaggac gcgcacaatg gcgctgatta tgggtggcatc     1080
tggtatcgta attttacgat gaactggctg ggagaacaat gtggacaaaa acccgcaag      1140
cgcaaaaaaa aaggcggcaa aaacggtaaa aatcgctgta accgtaagaa aaaaaatcct     1200
tga                                                                    1203

```

```

<210> 25
<211> 400
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Synthetic Construct

```

```

<400> 25

```

```

Met Gly Glu Leu Pro Ala Ser Met Ser Gln Ala Gln His Leu Ala Ala
1           5           10           15

```

Asn Thr Ala Thr Asp Asn Tyr Arg Ile Pro Ala Ile Thr Thr Ala Pro
20 25 30

Asn Gly Asp Leu Leu Ile Ser Tyr Asp Glu Arg Pro Lys Asp Asn Gly
35 40 45

Asn Gly Gly Ser Asp Ala Pro Asn Pro Asn His Ile Val Gln Arg Arg
50 55 60

Ser Thr Asp Gly Gly Lys Thr Trp Ser Ala Pro Thr Tyr Ile His Gln
65 70 75 80

Gly Thr Glu Thr Gly Lys Lys Val Gly Tyr Ser Asp Pro Ser Tyr Val
85 90 95

Val Asp His Gln Thr Gly Thr Ile Phe Asn Phe His Val Lys Ser Tyr
100 105 110

Asp Gln Gly Trp Gly Gly Ser Arg Gly Gly Thr Asp Pro Glu Asn Arg
115 120 125

Gly Ile Ile Gln Ala Glu Val Ser Thr Ser Thr Asp Asn Gly Trp Thr
130 135 140

Trp Thr His Arg Thr Ile Thr Ala Asp Ile Thr Lys Asp Lys Pro Trp
145 150 155 160

Thr Ala Arg Phe Ala Ala Ser Gly Gln Gly Ile Gln Ile Gln His Gly
165 170 175

Pro His Ala Gly Arg Leu Val Gln Gln Tyr Thr Ile Arg Thr Ala Gly
180 185 190

Gly Ala Val Gln Ala Val Ser Val Tyr Ser Asp Asp His Gly Lys Thr
195 200 205

Trp Gln Ala Gly Thr Pro Ile Gly Thr Gly Met Asp Glu Asn Lys Val
210 215 220

Val Glu Leu Ser Asp Gly Ser Leu Met Leu Asn Ser Arg Ala Ser Asp
225 230 235 240

Gly Ser Gly Phe Arg Lys Val Ala His Ser Thr Asp Gly Gly Gln Thr
 245 250 255

Trp Ser Glu Pro Val Ser Asp Lys Asn Leu Pro Asp Ser Val Asp Asn
 260 265 270

Ala Gln Ile Ile Arg Ala Phe Pro Asn Ala Ala Pro Asp Asp Pro Arg
 275 280 285

Ala Lys Val Leu Leu Leu Ser His Ser Pro Asn Pro Arg Pro Trp Ser
 290 295 300

Arg Asp Arg Gly Thr Ile Ser Met Ser Cys Asp Asp Gly Ala Ser Trp
 305 310 315 320

Thr Thr Ser Lys Val Phe His Glu Pro Phe Val Gly Tyr Thr Thr Ile
 325 330 335

Ala Val Gln Ser Asp Gly Ser Ile Gly Leu Leu Ser Glu Asp Ala His
 340 345 350

Asn Gly Ala Asp Tyr Gly Gly Ile Trp Tyr Arg Asn Phe Thr Met Asn
 355 360 365

Trp Leu Gly Glu Gln Cys Gly Gln Lys Pro Ala Lys Arg Lys Lys Lys
 370 375 380

Gly Gly Lys Asn Gly Lys Asn Arg Arg Asn Arg Lys Lys Lys Asn Pro
 385 390 395 400

<210> 26

<211> 1248

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 26

atgggagagc tgccagcaag catgtctcag gctcagcatc ttgcagcaaa tacggctact 60

gataattatc gcattccagc gattacaacc gctccgaatg gtgatttact gattagctat 120

gatgaacggc cgaaggacaa tggaaatggt gggtccgatg cccctaacc gaatcatatt 180

gttcagcgtc gctccacaga tggcggtaaa acttgagcgc cgccaaccta tattcatcag 240


```

ggtacggaga ctggcaagaa agtgggatat tccgaccctt cttatgtggt ggatcatcaa 300
accggtacaa tcttcaattt tcatgtgaaa tcatacgatc agggctgggg aggtagccgt 360
gggggaacag acccggaata ccgcgggatt attcaggcag aggtgtctac gagcacggat 420
aatggatgga cgtggacaca tcgcaccatc accgcggata ttacgaaaga taaaccgtgg 480
accgcgcgtt ttgcggcgct cggccaaggc attcagatcc agcatgggcc gcatgccggc 540
cgtctgggtg aacagtatac cattcgtacg gccggtggag cgggtgcaggc tgtatcggtt 600
tattccgatg atcatgggaa aacgtggcag gctggcacc cgaattgggac gggtatggat 660
gaaaacaaag ttgtagagct gtctgacggc tctctgatgc tgaacagtcg tgcgtcggac 720
gggagcggct ttcgtaaggt tgcgcatagc actgatgggt ggcagacctg gtccgaaccg 780
gtttcggaca aaaatttgcc ggattcgggt gataatgccc agataattcg tgcgtttcct 840
aatgctgccc ccgatgaccc gcgcgcgaaa gtacttcttc tgagtcattc cccaaatcca 900
cgtccgtggt cccgggatcg tggtagata agcatgtcat gtgatgacgg ggcctcatgg 960
accacttcca aagtttttca cgaaccgttt gtgggctaca cgactattgc agttcagagt 1020
gatggaagca tcgggtctgt gtcggaggac gcgcacaatg gcgctgatta tgggtggcatc 1080
tggtagcgtg attttacgat gaactggctg ggagaacaat gtggacaaaa acccgcgga 1140
ccgagcccag cccctagccc tactgcagca ccgtccgctg caaagcgcaa aaaaaaaggc 1200
ggcaaaaacg gtaaaaatcg tcgtaaccgt aagaaaaaaa atccttga 1248

```

```

<210> 27
<211> 415
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Synthetic Construct

```

```

<400> 27

```

```

Met Gly Glu Leu Pro Ala Ser Met Ser Gln Ala Gln His Leu Ala Ala
1           5           10          15

```

```

Asn Thr Ala Thr Asp Asn Tyr Arg Ile Pro Ala Ile Thr Thr Ala Pro
20          25          30

```

```

Asn Gly Asp Leu Leu Ile Ser Tyr Asp Glu Arg Pro Lys Asp Asn Gly
35          40          45

```

Asn Gly Gly Ser Asp Ala Pro Asn Pro Asn His Ile Val Gln Arg Arg
 50 55 60

Ser Thr Asp Gly Gly Lys Thr Trp Ser Ala Pro Thr Tyr Ile His Gln
 65 70 75 80

Gly Thr Glu Thr Gly Lys Lys Val Gly Tyr Ser Asp Pro Ser Tyr Val
 85 90 95

Val Asp His Gln Thr Gly Thr Ile Phe Asn Phe His Val Lys Ser Tyr
 100 105 110

Asp Gln Gly Trp Gly Gly Ser Arg Gly Gly Thr Asp Pro Glu Asn Arg
 115 120 125

Gly Ile Ile Gln Ala Glu Val Ser Thr Ser Thr Asp Asn Gly Trp Thr
 130 135 140

Trp Thr His Arg Thr Ile Thr Ala Asp Ile Thr Lys Asp Lys Pro Trp
 145 150 155 160

Thr Ala Arg Phe Ala Ala Ser Gly Gln Gly Ile Gln Ile Gln His Gly
 165 170 175

Pro His Ala Gly Arg Leu Val Gln Gln Tyr Thr Ile Arg Thr Ala Gly
 180 185 190

Gly Ala Val Gln Ala Val Ser Val Tyr Ser Asp Asp His Gly Lys Thr
 195 200 205

Trp Gln Ala Gly Thr Pro Ile Gly Thr Gly Met Asp Glu Asn Lys Val
 210 215 220

Val Glu Leu Ser Asp Gly Ser Leu Met Leu Asn Ser Arg Ala Ser Asp
 225 230 235 240

Gly Ser Gly Phe Arg Lys Val Ala His Ser Thr Asp Gly Gly Gln Thr
 245 250 255

Trp Ser Glu Pro Val Ser Asp Lys Asn Leu Pro Asp Ser Val Asp Asn
 260 265 270

Ala Gln Ile Ile Arg Ala Phe Pro Asn Ala Ala Pro Asp Asp Pro Arg
275 280 285

Ala Lys Val Leu Leu Leu Ser His Ser Pro Asn Pro Arg Pro Trp Ser
290 295 300

Arg Asp Arg Gly Thr Ile Ser Met Ser Cys Asp Asp Gly Ala Ser Trp
305 310 315 320

Thr Thr Ser Lys Val Phe His Glu Pro Phe Val Gly Tyr Thr Thr Ile
325 330 335

Ala Val Gln Ser Asp Gly Ser Ile Gly Leu Leu Ser Glu Asp Ala His
340 345 350

Asn Gly Ala Asp Tyr Gly Gly Ile Trp Tyr Arg Asn Phe Thr Met Asn
355 360 365

Trp Leu Gly Glu Gln Cys Gly Gln Lys Pro Ala Glu Pro Ser Pro Ala
370 375 380

Pro Ser Pro Thr Ala Ala Pro Ser Ala Ala Lys Arg Lys Lys Lys Gly
385 390 395 400

Gly Lys Asn Gly Lys Asn Arg Arg Asn Arg Lys Lys Lys Asn Pro
405 410 415

<210> 28
<211> 1221
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<400> 28
ccatggggca tcaccatcac catcatctag agggagatca tccacaagct acaccagcac 60
ctgcaccaga tgctagcact gagctgccag caagcatgtc tcaggctcag catcttgcag 120
caaatacggc tactgataat tatcgcatte cagcgattac aaccgctccg aatgggtgatt 180
tactgattag ctatgatgaa cggccgaagg acaatggaaa tgggtggttcc gatgcccccta 240
acccgaatca tattgttcag cgctcgctcca cagatggcgg taaaacttgg agcgcgccaa 300
cctatattca tcagggtacg gagactggca agaaagtggg atattccgac ccctcttatg 360

```

tgggtggatca tcaaaccggt acaatcttca attttcatgt gaaatcatac gatcaggggt 420
ggggaggttag ccgtggggga acagaccggg aaaaccgcgg gattattcag gcagaggtgt 480
ctacgagcac ggataatgga tggacgtgga cacatcgcac catcaccgcg gatattacga 540
aagataaacc gtggaccgcg cgttttgcg cgtccggcca aggcattcag atccagcatg 600
ggccgcgatgc cggccgtctg gtgcaacagt ataccattcg tacggccggt ggagcgggtgc 660
aggctgtatc ggtttattcc gatgatcatg ggaaaacgtg gcaggctggc accccgattg 720
ggacgggtat ggatgaaaac aaagttgtag agctgtctga cggctctctg atgctgaaca 780
gtcgtgcgtc ggacgggagc ggctttcgta aggttgcgca tagcactgat ggtgggcaga 840
cctgggtccga accggtttcg gacaaaaatt tgccggattc ggttgataat gcccagataa 900
ttcgtgcgtt tctaattgct gccccgatg acccgcgcgc gaaagtactt cttctgagtc 960
attccccaaa tccacgtccg tgggtcccggg atcgtggtac gataagcatg tcatgtgatg 1020
acggggcctc atggaccact tccaaagttt ttcacgaacc gtttgtgggc tacacgacta 1080
ttgcagttca gagtgatgga agcatcggtc tgctgtcgga ggacgcgcac aatggcgctg 1140
attatgggtg catctggtat cgtaatttta cgatgaactg gctgggagaa caatgtggac 1200
aaaaaccgcg ggaataagct t 1221

```

<210> 29
 <211> 404
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 29

```

Met Gly His His His His His His Leu Glu Gly Asp His Pro Gln Ala
1           5           10           15

```

```

Thr Pro Ala Pro Ala Pro Asp Ala Ser Thr Glu Leu Pro Ala Ser Met
20           25           30

```

```

Ser Gln Ala Gln His Leu Ala Ala Asn Thr Ala Thr Asp Asn Tyr Arg
35           40           45

```

```

Ile Pro Ala Ile Thr Thr Ala Pro Asn Gly Asp Leu Leu Ile Ser Tyr
50           55           60

```

Asp Glu Arg Pro Lys Asp Asn Gly Asn Gly Gly Ser Asp Ala Pro Asn
65 70 75 80

Pro Asn His Ile Val Gln Arg Arg Ser Thr Asp Gly Gly Lys Thr Trp
85 90 95

Ser Ala Pro Thr Tyr Ile His Gln Gly Thr Glu Thr Gly Lys Lys Val
100 105 110

Gly Tyr Ser Asp Pro Ser Tyr Val Val Asp His Gln Thr Gly Thr Ile
115 120 125

Phe Asn Phe His Val Lys Ser Tyr Asp Gln Gly Trp Gly Gly Ser Arg
130 135 140

Gly Gly Thr Asp Pro Glu Asn Arg Gly Ile Ile Gln Ala Glu Val Ser
145 150 155 160

Thr Ser Thr Asp Asn Gly Trp Thr Trp Thr His Arg Thr Ile Thr Ala
165 170 175

Asp Ile Thr Lys Asp Lys Pro Trp Thr Ala Arg Phe Ala Ala Ser Gly
180 185 190

Gln Gly Ile Gln Ile Gln His Gly Pro His Ala Gly Arg Leu Val Gln
195 200 205

Gln Tyr Thr Ile Arg Thr Ala Gly Gly Ala Val Gln Ala Val Ser Val
210 215 220

Tyr Ser Asp Asp His Gly Lys Thr Trp Gln Ala Gly Thr Pro Ile Gly
225 230 235 240

Thr Gly Met Asp Glu Asn Lys Val Val Glu Leu Ser Asp Gly Ser Leu
245 250 255

Met Leu Asn Ser Arg Ala Ser Asp Gly Ser Gly Phe Arg Lys Val Ala
260 265 270

His Ser Thr Asp Gly Gly Gln Thr Trp Ser Glu Pro Val Ser Asp Lys
275 280 285

Asn Leu Pro Asp Ser Val Asp Asn Ala Gln Ile Ile Arg Ala Phe Pro
 290 295 300

Asn Ala Ala Pro Asp Asp Pro Arg Ala Lys Val Leu Leu Leu Ser His
 305 310 315 320

Ser Pro Asn Pro Arg Pro Trp Ser Arg Asp Arg Gly Thr Ile Ser Met
 325 330 335

Ser Cys Asp Asp Gly Ala Ser Trp Thr Thr Ser Lys Val Phe His Glu
 340 345 350

Pro Phe Val Gly Tyr Thr Thr Ile Ala Val Gln Ser Asp Gly Ser Ile
 355 360 365

Gly Leu Leu Ser Glu Asp Ala His Asn Gly Ala Asp Tyr Gly Gly Ile
 370 375 380

Trp Tyr Arg Asn Phe Thr Met Asn Trp Leu Gly Glu Gln Cys Gly Gln
 385 390 395 400

Lys Pro Ala Glu

<210> 30
 <211> 1257
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 30
 ccatgaagcg caaaaaaaaaa ggcggcaaaa acggtaaaaa tcgtcgtaac cgtaagaaaa 60
 aaaatcctgg agatcatcca caagctacac cagcacctgc accagatgct agcactgagc 120
 tgccagcaag catgtctcag gctcagcatc ttgcagcaaa tacggctact gataattatc 180
 gcattccagc gattacaacc gctccgaatg gtgatttact gattagctat gatgaacggc 240
 cgaaggacaa tggaaatggg gggtccgatg cccctaaccg gaatcatatt gttcagcgtc 300
 gctccacaga tggcggtaaa acttggagcg cgccaacctt tattcatcag ggtacggaga 360
 ctggcaagaa agtgggatat tccgaccctt cttatgtggg ggatcatcaa accggtacaa 420
 tcttcaattt tcatgtgaaa tcatacgatc agggctgggg aggtagccgt gggggaacag 480

acccggaataa ccgcgggatt attcaggcag aggtgtctac gagcacggat aatggatgga 540
 cgtggacaca tcgcaccatc accgcggata ttacgaaaga taaaccgtgg accgcgcgtt 600
 ttgcggcgtc cggccaaggc attcagatcc agcatgggcc gcatgccggc cgtctggtgc 660
 aacagtatac cattcgtacg gccgggtggag cgggtgcaggc tgtatcggtt tattccgatg 720
 atcatgggaa aacgtggcag gctggcaccc cgattgggac gggatatggat gaaaacaaag 780
 ttgtagagct gtctgacggc tctctgatgc tgaacagtcg tgcgtcggac gggagcggct 840
 ttcgtaaggt tgcgcatagc actgatgggtg ggcagacctg gtccgaaccg gtttcggaca 900
 aaaatttgcc ggattcgggtt gataatgccc agataattcg tgcgtttcct aatgctgccc 960
 ccgatgaccc gcgcgcgaaa gtactttctt tgagtcattc cccaaatcca cgtccgtggt 1020
 cccgggatcg tggtacgata agcatgtcat gtgatgacgg ggcctcatgg accacttcca 1080
 aagtttttca cgaaccgttt gtgggctaca cgactattgc agttcagagt gatggaagca 1140
 tcggtctgct gtcggaggac gcgcacaatg gcgctgatta tgggtggcatc tggtatcgta 1200
 attttacgat gaactggctg ggagaacaat gtggacaaaa acccgcgga taagctt 1257

<210> 31
 <211> 416
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 31

Met Lys Arg Lys Lys Lys Gly Gly Lys Asn Gly Lys Asn Arg Arg Asn
 1 5 10 15

Arg Lys Lys Lys Asn Pro Gly Asp His Pro Gln Ala Thr Pro Ala Pro
 20 25 30

Ala Pro Asp Ala Ser Thr Glu Leu Pro Ala Ser Met Ser Gln Ala Gln
 35 40 45

His Leu Ala Ala Asn Thr Ala Thr Asp Asn Tyr Arg Ile Pro Ala Ile
 50 55 60

Thr Thr Ala Pro Asn Gly Asp Leu Leu Ile Ser Tyr Asp Glu Arg Pro
 65 70 75 80

Lys Asp Asn Gly Asn Gly Gly Ser Asp Ala Pro Asn Pro Asn His Ile
 85 90 95

Val Gln Arg Arg Ser Thr Asp Gly Gly Lys Thr Trp Ser Ala Pro Thr
 100 105 110

Tyr Ile His Gln Gly Thr Glu Thr Gly Lys Lys Val Gly Tyr Ser Asp
 115 120 125

Pro Ser Tyr Val Val Asp His Gln Thr Gly Thr Ile Phe Asn Phe His
 130 135 140

Val Lys Ser Tyr Asp Gln Gly Trp Gly Gly Ser Arg Gly Gly Thr Asp
 145 150 155 160

Pro Glu Asn Arg Gly Ile Ile Gln Ala Glu Val Ser Thr Ser Thr Asp
 165 170 175

Asn Gly Trp Thr Trp Thr His Arg Thr Ile Thr Ala Asp Ile Thr Lys
 180 185 190

Asp Lys Pro Trp Thr Ala Arg Phe Ala Ala Ser Gly Gln Gly Ile Gln
 195 200 205

Ile Gln His Gly Pro His Ala Gly Arg Leu Val Gln Gln Tyr Thr Ile
 210 215 220

Arg Thr Ala Gly Gly Ala Val Gln Ala Val Ser Val Tyr Ser Asp Asp
 225 230 235 240

His Gly Lys Thr Trp Gln Ala Gly Thr Pro Ile Gly Thr Gly Met Asp
 245 250 255

Glu Asn Lys Val Val Glu Leu Ser Asp Gly Ser Leu Met Leu Asn Ser
 260 265 270

Arg Ala Ser Asp Gly Ser Gly Phe Arg Lys Val Ala His Ser Thr Asp
 275 280 285

Gly Gly Gln Thr Trp Ser Glu Pro Val Ser Asp Lys Asn Leu Pro Asp
 290 295 300

Ser Val Asp Asn Ala Gln Ile Ile Arg Ala Phe Pro Asn Ala Ala Pro
305 310 315 320

Asp Asp Pro Arg Ala Lys Val Leu Leu Leu Ser His Ser Pro Asn Pro
325 330 335

Arg Pro Trp Ser Arg Asp Arg Gly Thr Ile Ser Met Ser Cys Asp Asp
340 345 350

Gly Ala Ser Trp Thr Thr Ser Lys Val Phe His Glu Pro Phe Val Gly
355 360 365

Tyr Thr Thr Ile Ala Val Gln Ser Asp Gly Ser Ile Gly Leu Leu Ser
370 375 380

Glu Asp Ala His Asn Gly Ala Asp Tyr Gly Gly Ile Trp Tyr Arg Asn
385 390 395 400

Phe Thr Met Asn Trp Leu Gly Glu Gln Cys Gly Gln Lys Pro Ala Glu
405 410 415

<210> 32
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<220>
<221> misc_feature
<222> (17)..(18)
<223> n is a, c, g, or t

<220>
<221> misc_feature
<222> (20)..(21)
<223> n is a, c, g, or t

<400> 32
ttttcgtctc ccatgvnnvn naagcgcaaa aaaaaaggcg gca

43

<210> 33
<211> 10
<212> PRT
<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> misc_feature

<222> (2)..(3)

<223> Xaa can be any naturally occurring amino acid

<400> 33

Met Xaa Xaa Lys Arg Lys Lys Lys Gly Gly
1 5 10

<210> 34

<211> 1272

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 34

ccatgaagcg	caaaaaaaaa	ggcggcaaaa	acggtaaaaa	tcgtcgtaac	cgtaagaaaa	60
aaaatcctgg	tgggtggtgt	tctggagatc	atccacaagc	tacaccagca	cctgcaccag	120
atgctagcac	tgagctgcc	gcaagcatgt	ctcaggctca	gcatcttgca	gcaaatacgg	180
ctactgataa	ttatcgcatt	ccagcgatta	caaccgctcc	gaatggatgat	ttactgatta	240
gctatgatga	acggccgaag	gacaatggaa	atgggtggttc	cgatgcccct	aacccgaatc	300
atattgttca	gcgtcgctcc	acagatggcg	gtaaaacttg	gagcgcgcca	acctatattc	360
atcaggggtac	ggagactggc	aagaaagtgg	gatattccga	cccctcttat	gtggtggatc	420
atcaaaccgg	tacaatcttc	aattttcatg	tgaaatcata	cgatcagggc	tggggaggta	480
gccgtggggg	aacagacccg	gaaaaccgcg	ggattattca	ggcagagggtg	tctacgagca	540
cggataatgg	atggacgtgg	acacatcgca	ccatcaccgc	ggatattacg	aaagataaac	600
cgtggaccgc	gcgttttgcg	gcgtccggcc	aaggcattca	gatccagcat	gggccgcatg	660
ccggccgctct	ggtgcaacag	tataccattc	gtacggccgg	tggagcgggtg	caggctgtat	720
cggttttattc	cgatgatcat	gggaaaacgt	ggcaggctgg	caccccgatt	gggacgggta	780
tggatgaaaa	caaagtgtga	gagctgtctg	acggctctct	gatgctgaac	agtcgtgcgt	840
cggacggggag	cggcttttcgt	aaggttgcgc	atagcactga	tgggtgggcag	acctggtccg	900
aaccggtttc	ggacaaaaat	ttgccggatt	cggttgataa	tgcccagata	attcgtgcgt	960
ttcctaatagc	tgcccccgat	gaccgcgcgc	cgaaagtact	tcttctgagt	cattccccaa	1020

atccacgtcc gtgggtcccg gatcgtggta cgataagcat gtcatgtgat gacggggcct 1080
 catggaccac ttccaaagtt tttcacgaac cgtttgtggg ctacacgact attgcagttc 1140
 agagtgatgg aagcatcggt ctgctgtcgg aggacgcgca caatggcgct gattatggtg 1200
 gcatctggta tcgtaatttt acgatgaact ggctgggaga acaatgtgga caaaaacccg 1260
 cggaataagc tt 1272

<210> 35
 <211> 421
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 35

Met Lys Arg Lys Lys Lys Gly Gly Lys Asn Gly Lys Asn Arg Arg Asn
 1 5 10 15

Arg Lys Lys Lys Asn Pro Gly Gly Gly Gly Ser Gly Asp His Pro Gln
 20 25 30

Ala Thr Pro Ala Pro Ala Pro Asp Ala Ser Thr Glu Leu Pro Ala Ser
 35 40 45

Met Ser Gln Ala Gln His Leu Ala Ala Asn Thr Ala Thr Asp Asn Tyr
 50 55 60

Arg Ile Pro Ala Ile Thr Thr Ala Pro Asn Gly Asp Leu Leu Ile Ser
 65 70 75 80

Tyr Asp Glu Arg Pro Lys Asp Asn Gly Asn Gly Gly Ser Asp Ala Pro
 85 90 95

Asn Pro Asn His Ile Val Gln Arg Arg Ser Thr Asp Gly Gly Lys Thr
 100 105 110

Trp Ser Ala Pro Thr Tyr Ile His Gln Gly Thr Glu Thr Gly Lys Lys
 115 120 125

Val Gly Tyr Ser Asp Pro Ser Tyr Val Val Asp His Gln Thr Gly Thr
 130 135 140

Ile Phe Asn Phe His Val Lys Ser Tyr Asp Gln Gly Trp Gly Gly Ser
145 150 155 160

Arg Gly Gly Thr Asp Pro Glu Asn Arg Gly Ile Ile Gln Ala Glu Val
165 170 175

Ser Thr Ser Thr Asp Asn Gly Trp Thr Trp Thr His Arg Thr Ile Thr
180 185 190

Ala Asp Ile Thr Lys Asp Lys Pro Trp Thr Ala Arg Phe Ala Ala Ser
195 200 205

Gly Gln Gly Ile Gln Ile Gln His Gly Pro His Ala Gly Arg Leu Val
210 215 220

Gln Gln Tyr Thr Ile Arg Thr Ala Gly Gly Ala Val Gln Ala Val Ser
225 230 235 240

Val Tyr Ser Asp Asp His Gly Lys Thr Trp Gln Ala Gly Thr Pro Ile
245 250 255

Gly Thr Gly Met Asp Glu Asn Lys Val Val Glu Leu Ser Asp Gly Ser
260 265 270

Leu Met Leu Asn Ser Arg Ala Ser Asp Gly Ser Gly Phe Arg Lys Val
275 280 285

Ala His Ser Thr Asp Gly Gly Gln Thr Trp Ser Glu Pro Val Ser Asp
290 295 300

Lys Asn Leu Pro Asp Ser Val Asp Asn Ala Gln Ile Ile Arg Ala Phe
305 310 315 320

Pro Asn Ala Ala Pro Asp Asp Pro Arg Ala Lys Val Leu Leu Leu Ser
325 330 335

His Ser Pro Asn Pro Arg Pro Trp Ser Arg Asp Arg Gly Thr Ile Ser
340 345 350

Met Ser Cys Asp Asp Gly Ala Ser Trp Thr Thr Ser Lys Val Phe His
355 360 365

Glu Pro Phe Val Gly Tyr Thr Thr Ile Ala Val Gln Ser Asp Gly Ser
 370 375 380

Ile Gly Leu Leu Ser Glu Asp Ala His Asn Gly Ala Asp Tyr Gly Gly
 385 390 395 400

Ile Trp Tyr Arg Asn Phe Thr Met Asn Trp Leu Gly Glu Gln Cys Gly
 405 410 415

Gln Lys Pro Ala Glu
 420

<210> 36
 <211> 1275
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 36
 ccatgggttaa gcgcaaaaaa aaaggcggca aaaacggtaa aaatcgtcgt aaccgtaaga 60
 aaaaaaatcc tgggtggtggt ggttctggag atcatccaca agctacacca gcacctgcac 120
 cagatgctag cactgagctg ccagcaagca tgtctcaggc tcagcatctt gcagcaaata 180
 cggctactga taattatcgc attccagcga ttacaaccgc tccgaatggt gatttactga 240
 ttagctatga tgaacggccg aaggacaatg gaaatggtgg ttccgatgcc cctaaccgga 300
 atcatattgt tcagcgtcgc tccacagatg gcggtaaaac ttggagcgcg ccaacctata 360
 ttcacagggg tacggagact ggcaagaaag tgggatattc cgaccctct tatgtggtgg 420
 atcatcaaac cggtaaatc ttcaattttc atgtgaaatc atacgatcag ggctggggag 480
 gtagccgtgg gggacagac ccggaaaacc gcgggattat tcaggcagag gtgtctacga 540
 gcacggataa tggatggacg tggacacatc gcacatcac cgcggatatt acgaaagata 600
 aaccgtggac cgcgcgtttt gcggcgtccg gccaaaggcat tcagatccag catgggccgc 660
 atgccggccg tctggtgcaa cagtatacca ttcgtacggc cgggtggagcg gtgcaggctg 720
 tatcggttta ttccgatgat catgggaaaa cgtggcaggc tggcaccctg attgggacgg 780
 gtaggatga aaacaaagt gtgagctgt ctgacggctc tctgatgctg aacagtcgtg 840
 cgtcggacgg gagcggcttt cgtaagggtg cgcatagcac tgatgggtggg cagacctggt 900
 ccgaaccggt ttcggacaaa aatttgccgg attcggttga taatgccag ataattcgtg 960

cgtttcctaa tgctgcccc gatgaccgc gcgcgaaagt acttcttctg agtcattccc 1020
 caaatccacg tccgtggtcc cgggatcgtg gtacgataag catgtcatgt gatgacgggg 1080
 cctcatggac cacttccaaa gtttttcacg aaccgtttgt gggctacacg actattgcag 1140
 ttcagagtga tggaagcatc ggtctgctgt cggaggacgc gcacaatggc gctgattatg 1200
 gtggcatctg gtatcgtaat tttacgatga actggctggg agaacaatgt ggacaaaaac 1260
 ccgcggaata agctt 1275

<210> 37
 <211> 422
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 37

Met Val Lys Arg Lys Lys Lys Gly Gly Lys Asn Gly Lys Asn Arg Arg
 1 5 10 15

Asn Arg Lys Lys Lys Asn Pro Gly Gly Gly Ser Gly Asp His Pro
 20 25 30

Gln Ala Thr Pro Ala Pro Ala Pro Asp Ala Ser Thr Glu Leu Pro Ala
 35 40 45

Ser Met Ser Gln Ala Gln His Leu Ala Ala Asn Thr Ala Thr Asp Asn
 50 55 60

Tyr Arg Ile Pro Ala Ile Thr Thr Ala Pro Asn Gly Asp Leu Leu Ile
 65 70 75 80

Ser Tyr Asp Glu Arg Pro Lys Asp Asn Gly Asn Gly Gly Ser Asp Ala
 85 90 95

Pro Asn Pro Asn His Ile Val Gln Arg Arg Ser Thr Asp Gly Gly Lys
 100 105 110

Thr Trp Ser Ala Pro Thr Tyr Ile His Gln Gly Thr Glu Thr Gly Lys
 115 120 125

Lys Val Gly Tyr Ser Asp Pro Ser Tyr Val Val Asp His Gln Thr Gly
 130 135 140

Thr Ile Phe Asn Phe His Val Lys Ser Tyr Asp Gln Gly Trp Gly Gly
145 150 155 160

Ser Arg Gly Gly Thr Asp Pro Glu Asn Arg Gly Ile Ile Gln Ala Glu
165 170 175

Val Ser Thr Ser Thr Asp Asn Gly Trp Thr Trp Thr His Arg Thr Ile
180 185 190

Thr Ala Asp Ile Thr Lys Asp Lys Pro Trp Thr Ala Arg Phe Ala Ala
195 200 205

Ser Gly Gln Gly Ile Gln Ile Gln His Gly Pro His Ala Gly Arg Leu
210 215 220

Val Gln Gln Tyr Thr Ile Arg Thr Ala Gly Gly Ala Val Gln Ala Val
225 230 235 240

Ser Val Tyr Ser Asp Asp His Gly Lys Thr Trp Gln Ala Gly Thr Pro
245 250 255

Ile Gly Thr Gly Met Asp Glu Asn Lys Val Val Glu Leu Ser Asp Gly
260 265 270

Ser Leu Met Leu Asn Ser Arg Ala Ser Asp Gly Ser Gly Phe Arg Lys
275 280 285

Val Ala His Ser Thr Asp Gly Gly Gln Thr Trp Ser Glu Pro Val Ser
290 295 300

Asp Lys Asn Leu Pro Asp Ser Val Asp Asn Ala Gln Ile Ile Arg Ala
305 310 315 320

Phe Pro Asn Ala Ala Pro Asp Asp Pro Arg Ala Lys Val Leu Leu Leu
325 330 335

Ser His Ser Pro Asn Pro Arg Pro Trp Ser Arg Asp Arg Gly Thr Ile
340 345 350

Ser Met Ser Cys Asp Asp Gly Ala Ser Trp Thr Thr Ser Lys Val Phe
355 360 365

His Glu Pro Phe Val Gly Tyr Thr Thr Ile Ala Val Gln Ser Asp Gly
370 375 380

Ser Ile Gly Leu Leu Ser Glu Asp Ala His Asn Gly Ala Asp Tyr Gly
385 390 395 400

Gly Ile Trp Tyr Arg Asn Phe Thr Met Asn Trp Leu Gly Glu Gln Cys
405 410 415

Gly Gln Lys Pro Ala Glu
420

<210> 38
<211> 416
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<400> 38

Met Lys Arg Lys Lys Lys Gly Gly Lys Asn Gly Lys Asn Arg Arg Asn
1 5 10 15

Arg Lys Lys Lys Asn Pro Gly Asp His Pro Gln Ala Thr Pro Ala Pro
20 25 30

Ala Pro Asp Ala Ser Thr Glu Leu Pro Ala Ser Met Ser Gln Ala Gln
35 40 45

His Leu Ala Ala Asn Thr Ala Thr Asp Asn Tyr Arg Ile Pro Ala Ile
50 55 60

Thr Thr Ala Pro Asn Gly Asp Leu Leu Ile Ser Tyr Asp Glu Arg Pro
65 70 75 80

Lys Asp Asn Gly Asn Gly Gly Ser Asp Ala Pro Asn Pro Asn His Ile
85 90 95

Val Gln Arg Arg Ser Thr Asp Gly Gly Lys Thr Trp Ser Ala Pro Thr
100 105 110

Tyr Ile His Gln Gly Thr Glu Thr Gly Lys Lys Val Gly Tyr Ser Asp
115 120 125

Pro Ser Tyr Val Val Asp His Gln Thr Gly Thr Ile Phe Asn Phe His
130 135 140

Val Lys Ser Tyr Asp Gln Gly Trp Gly Gly Ser Arg Gly Gly Thr Asp
145 150 155 160

Pro Glu Asn Arg Gly Ile Ile Gln Ala Glu Val Ser Thr Ser Thr Asp
165 170 175

Asn Gly Trp Thr Trp Thr His Arg Thr Ile Thr Ala Asp Ile Thr Lys
180 185 190

Asp Lys Pro Trp Thr Ala Arg Phe Ala Ala Ser Gly Gln Gly Ile Gln
195 200 205

Ile Gln His Gly Pro His Ala Gly Arg Leu Val Gln Gln Tyr Thr Ile
210 215 220

Arg Thr Ala Gly Gly Ala Val Gln Ala Val Ser Val Tyr Ser Asp Asp
225 230 235 240

His Gly Lys Thr Trp Gln Ala Gly Thr Pro Ile Gly Thr Gly Met Asp
245 250 255

Glu Asn Lys Val Val Glu Leu Ser Asp Gly Ser Leu Met Leu Asn Ser
260 265 270

Arg Ala Ser Asp Gly Ser Gly Phe Arg Lys Val Ala His Ser Thr Asp
275 280 285

Gly Gly Gln Thr Trp Ser Glu Pro Val Ser Asp Lys Asn Leu Pro Asp
290 295 300

Ser Val Asp Asn Ala Gln Ile Ile Arg Ala Phe Pro Asn Ala Ala Pro
305 310 315 320

Asp Asp Pro Arg Ala Lys Val Leu Leu Leu Ser His Ser Pro Asn Pro
325 330 335

Arg Pro Trp Ser Arg Asp Arg Gly Thr Ile Ser Met Ser Cys Asp Asp
340 345 350

Gly Ala Ser Trp Thr Thr Ser Lys Val Phe His Glu Pro Phe Val Gly
 355 360 365

Tyr Thr Thr Ile Ala Val Gln Ser Asp Gly Ser Ile Gly Leu Leu Ser
 370 375 380

Glu Asp Ala His Asn Gly Ala Asp Tyr Gly Gly Ile Trp Tyr Arg Asn
 385 390 395 400

Phe Thr Met Asn Trp Leu Gly Glu Gln Cys Gly Gln Lys Pro Ala Glu
 405 410 415

<210> 39
 <211> 421
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 39

Val Lys Arg Lys Lys Lys Gly Gly Lys Asn Gly Lys Asn Arg Arg Asn
 1 5 10 15

Arg Lys Lys Lys Asn Pro Gly Gly Gly Gly Ser Gly Asp His Pro Gln
 20 25 30

Ala Thr Pro Ala Pro Ala Pro Asp Ala Ser Thr Glu Leu Pro Ala Ser
 35 40 45

Met Ser Gln Ala Gln His Leu Ala Ala Asn Thr Ala Thr Asp Asn Tyr
 50 55 60

Arg Ile Pro Ala Ile Thr Thr Ala Pro Asn Gly Asp Leu Leu Ile Ser
 65 70 75 80

Tyr Asp Glu Arg Pro Lys Asp Asn Gly Asn Gly Gly Ser Asp Ala Pro
 85 90 95

Asn Pro Asn His Ile Val Gln Arg Arg Ser Thr Asp Gly Gly Lys Thr
 100 105 110

Trp Ser Ala Pro Thr Tyr Ile His Gln Gly Thr Glu Thr Gly Lys Lys
 115 120 125

Val Gly Tyr Ser Asp Pro Ser Tyr Val Val Asp His Gln Thr Gly Thr
130 135 140

Ile Phe Asn Phe His Val Lys Ser Tyr Asp Gln Gly Trp Gly Gly Ser
145 150 155 160

Arg Gly Gly Thr Asp Pro Glu Asn Arg Gly Ile Ile Gln Ala Glu Val
165 170 175

Ser Thr Ser Thr Asp Asn Gly Trp Thr Trp Thr His Arg Thr Ile Thr
180 185 190

Ala Asp Ile Thr Lys Asp Lys Pro Trp Thr Ala Arg Phe Ala Ala Ser
195 200 205

Gly Gln Gly Ile Gln Ile Gln His Gly Pro His Ala Gly Arg Leu Val
210 215 220

Gln Gln Tyr Thr Ile Arg Thr Ala Gly Gly Ala Val Gln Ala Val Ser
225 230 235 240

Val Tyr Ser Asp Asp His Gly Lys Thr Trp Gln Ala Gly Thr Pro Ile
245 250 255

Gly Thr Gly Met Asp Glu Asn Lys Val Val Glu Leu Ser Asp Gly Ser
260 265 270

Leu Met Leu Asn Ser Arg Ala Ser Asp Gly Ser Gly Phe Arg Lys Val
275 280 285

Ala His Ser Thr Asp Gly Gly Gln Thr Trp Ser Glu Pro Val Ser Asp
290 295 300

Lys Asn Leu Pro Asp Ser Val Asp Asn Ala Gln Ile Ile Arg Ala Phe
305 310 315 320

Pro Asn Ala Ala Pro Asp Asp Pro Arg Ala Lys Val Leu Leu Leu Ser
325 330 335

His Ser Pro Asn Pro Arg Pro Trp Ser Arg Asp Arg Gly Thr Ile Ser
340 345 350

Met Ser Cys Asp Asp Gly Ala Ser Trp Thr Thr Ser Lys Val Phe His
355 360 365

Glu Pro Phe Val Gly Tyr Thr Thr Ile Ala Val Gln Ser Asp Gly Ser
370 375 380

Ile Gly Leu Leu Ser Glu Asp Ala His Asn Gly Ala Asp Tyr Gly Gly
385 390 395 400

Ile Trp Tyr Arg Asn Phe Thr Met Asn Trp Leu Gly Glu Gln Cys Gly
405 410 415

Gln Lys Pro Ala Glu
420